Acta Mathematica Academiae Paedagogicae Nyíregyháziensis
33 (2017), 221–231
www.emis.de/journals
ISSN 1786-0091

ON RIESZ ALMOST LACUNARY CESÀRO $[C, 1, 1, 1]$ STATISTICAL CONVERGENCE IN PROBABILISTIC SPACE OF $\chi_f^{3\Delta}$

SHYAMAL DEBNATH AND N. SUBRAMANIAN

Abstract. In this paper we study the concept of almost lacunary statistical Cesàro of χ^3 over probabilistic space P is defined by Musielak Orlicz function. Since the study of convergence in Probabilistic space P is fundamental to probabilistic functional analysis, we feel that the concept of almost lacunary statistical Cesàro of χ^3 over probabilistic space P is defined by Musielak in a probabilistic space P would provide a more general framework for the subject.

1. Introduction

A triple sequence (real or complex) can be defined as a function $x \colon \mathbb{N} \times \mathbb{N} \times \mathbb{N} \to \mathbb{R}\,(\mathbb{C})$, where $\mathbb{N}, \mathbb{R}$ and $\mathbb{C}$ denote the set of natural numbers, real numbers and complex numbers respectively. The different types of notions of triple sequence was introduced and investigated at the initial by *Sahiner et al.* [10, 11], *Esi et al.* [1, 2, 3], *Datta et al.* [4], *Subramanian et al.* [12], *Debnath et al.* [5] and many others.

A triple sequence $x = (x_{mnk})$ is said to be triple analytic if

$$\sup_{m,n,k} |x_{mnk}|^{\frac{1}{m+n+k}} < \infty.$$

The space of all triple analytic sequences are usually denoted by Λ^3. A triple sequence $x = (x_{mnk})$ is called triple entire sequence if

$$|x_{mnk}|^{\frac{1}{m+n+k}} \to 0 \text{ as } m, n, k \to \infty.$$

The space of all triple entire sequences are usually denoted by Γ^3. The space Λ^3 and Γ^3 is a metric space with the metric

$$d(x, y) = \sup_{m,n,k} \left\{ |x_{mnk} - y_{mnk}|^{\frac{1}{m+n+k}} : m, n, k : 1, 2, 3, \dots \right\}, \tag{1.1}$$

2010 *Mathematics Subject Classification.* 40F05, 40J05, 40G05.
Key words and phrases. analytic sequence, Orlicz function, triple sequences, chi sequence, Riesz space, statistical convergence,Cesàro $C_{1,1,1}-$ statistical convergence.

for all $x = \{x_{mnk}\}$ and $y = \{y_{mnk}\} in \Gamma^3$.

A complex sequence, whose k^{th} term x_k is denoted by $\{x_k\}$ or simply x. Let w be the set of all sequences $x = (x_k)$ and ϕ be the set of all finite sequences. Let ℓ_∞, c, c_0 be the sequence spaces of bounded,convergent and null sequences $x = (x_k)$ respectively. In respect of ℓ_∞, c, c_0 we have $\|x\| = \sup_k |x_k|$, where $x = (x_k) \in c_0 \subset c \subset \ell_\infty$.

The notion of difference sequence spaces (for single sequences) was introduced by Kizmaz [6] as follows

$$Z(\Delta) = \{x = (x_k) \in w : (\Delta x_k) \in Z\}$$

for $Z = c, c_0$ and ℓ_∞, where $\Delta x_k = x_k - x_{k+1}$ for all $k \in \mathbb{N}$.

The difference triple sequence space was introduced by Debnath et al. (see [5]) and is defined as

$\Delta x_{mnk} = x_{mnk} - x_{m,n+1,k} - x_{m,n,k+1} + x_{m,n+1,k+1} - x_{m+1,n,k} + x_{m+1,n+1,k} + x_{m+1,n,k+1} - x_{m+1,n+1,k+1}$ and $\Delta^0 x_{mnk} = \langle x_{mnk} \rangle$.

2. Definitions and Preliminaries

Throughout the article $w^3, \chi^3(\Delta), \Lambda^3(\Delta)$ denote the spaces of all, triple gai difference sequence spaces and triple analytic difference sequence spaces respectively.

Subramanian et al. (see [12]) introduced by a triple entire sequence spaces, triple analytic sequences spaces and triple gai sequence spaces. The triple sequence spaces of $\chi^3(\Delta), \Lambda^3(\Delta)$ are defined as follows:

$\chi^3(\Delta) = \left\{x \in w^3 : ((m+n+k)! \, |\Delta x_{mnk}|)^{1/m+n+k} \to 0 \text{ as } m, n, k \to \infty\right\}$,

$\Lambda^3(\Delta) = \left\{x \in w^3 : \sup_{m,n,k} |\Delta x_{mnk}|^{1/m+n+k} < \infty\right\}$.

Definition 2.1. An Orlicz function (see [7]) is a function $M\colon [0,\infty) \to [0,\infty)$ which is continuous, non-decreasing and convex with $M(0) = 0$, $M(x) > 0$, for $x > 0$ and $M(x) \to \infty$ as $x \to \infty$. If convexity of Orlicz function M is replaced by $M(x+y) \le M(x) + M(y)$, then this function is called modulus function.

Lindenstrauss and Tzafriri ([8]) used the idea of Orlicz function to construct Orlicz sequence space.

A sequence $g = (g_{mn})$ defined by

$$g_{mn}(v) = \sup\{|v| u - (f_{mnk})(u) : u \ge 0\}, m, n, k = 1, 2, \ldots$$

is called the complementary function of a Musielak-Orlicz function f. For a given Musielak-Orlicz function f, (see [9]) the Musielak-Orlicz sequence space t_f is defined as follows

$$t_f = \left\{x \in w^3 : I_f(|x_{mnk}|)^{1/m+n+k} \to 0 \text{ as } m, n, k \to \infty\right\},$$

where I_f is a convex modular defined by

$$I_f(x) = \sum_{m=1}^{\infty}\sum_{n=1}^{\infty}\sum_{k=1}^{\infty} f_{mnk}\left(|x_{mnk}|\right)^{1/m+n+k}, \quad x = (x_{mnk}) \in t_f.$$

We consider t_f equipped with the Luxemburg metric

$$d(x,y) = \sum_{m=1}^{\infty}\sum_{n=1}^{\infty}\sum_{k=1}^{\infty} f_{mnk}\left(\frac{|x_{mnk}|^{1/m+n+k}}{mnk}\right)$$

is an extended real number.

3. Further Definitions and Preliminaries

A triple sequence $x = (x_{mnk})$ has limit 0 (denoted by $P - limx = 0$) (i.e) $P - ((m+n+k)!\,|x_{mnk}|)^{1/m+n+k} \to 0$ as $m,n,k \to \infty$. We shall write more briefly as *P-convergent to 0.*

Definition 3.1. A triple sequence spaces of $x = (x_{mnk})$ of real numbers is called almost $P-$convergent to a limit 0 if

$$P - \lim_{p,q,u\to\infty} \sup_{r,s,t\geq 0} \frac{1}{pqu} \sum_{m=r}^{r+p-1}\sum_{n=s}^{s+q-1}\sum_{k=t}^{t+u-1} \left((m+n+k)!\,|x_{mnk}|\right)^{1/m+n+k} \to 0.$$

that is, the average value of (x_{mnk}) taken over any rectangle

$\{(m,n,k) : r \leq m \leq r+p-1, s \leq n \leq s+q-1, t \leq k \leq t+u-1\}$ tends to 0 as both p,q and u to ∞, and this $P-$convergence is uniform in i,ℓ and j. Let denote the set of sequences with this property as $\left[\widehat{\chi^3}\right]$.

Definition 3.2. Let $(q_m), (\overline{q_n}), (\overline{\overline{q_k}})$ be sequences of positive numbers and

$$Q_r = \begin{bmatrix} q_{11} & q_{12} & \cdots & q_{1s} & 0\ldots \\ q_{21} & q_{22} & \cdots & q_{2s} & 0\ldots \\ . & & & & \\ . & & & & \\ . & & & & \\ q_{r1} & q_{r2} & \cdots & q_{rs} & 0\ldots \\ 0 & 0 & \ldots 0 & 0 & 0\ldots \\ . & . & \ldots. & . & \ldots. \end{bmatrix} = q_{11} + q_{12} + \ldots + q_{rs} + \cdots \neq 0,$$

$$\overline{Q}_s = \begin{bmatrix} \overline{q}_{11} & \overline{q}_{12} & \cdots & \overline{q}_{1s} & 0\ldots \\ \overline{q}_{21} & \overline{q}_{22} & \cdots & \overline{q}_{2s} & 0\ldots \\ . & & & & \\ . & & & & \\ . & & & & \\ \overline{q}_{r1} & \overline{q}_{r2} & \cdots & \overline{q}_{rs} & 0\ldots \\ 0 & 0 & \ldots 0 & 0 & 0\ldots \\ . & . & \ldots. & . & \ldots. \end{bmatrix} = \overline{q}_{11} + \overline{q}_{12} + \ldots + \overline{q}_{rs} + \cdots \neq 0,$$

$$\overline{\overline{Q}}_t = \begin{bmatrix} \overline{\overline{q}}_{11} & \overline{\overline{q}}_{12} & \ldots & \overline{\overline{q}}_{1s} & 0\ldots \\ \overline{\overline{q}}_{21} & \overline{\overline{q}}_{22} & \ldots & \overline{\overline{q}}_{2s} & 0\ldots \\ . & & & & \\ . & & & & \\ . & & & & \\ \overline{\overline{q}}_{r1} & \overline{\overline{q}}_{r2} & \ldots & \overline{\overline{q}}_{rs} & 0\ldots \\ 0 & 0 & \ldots 0 & 0 & 0\ldots \\ . & . & \ldots. & . & \ldots. \end{bmatrix} = \overline{\overline{q}}_{11} + \overline{\overline{q}}_{12} + \cdots + \overline{\overline{q}}_{rs} + \cdots \neq 0.$$

Then the transformation is given by

$$T_{rst} = \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r} \sum_{n=1}^{s} \sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left((m+n+k)! \, |x_{mnk}|\right)^{1/m+n+k}$$

is called the Riesz mean of triple sequence spaces of $x = (x_{mnk})$. If $P - \lim_{rst} T_{rst}(x) = 0, 0 \in \mathbb{R}$, then the triple sequence spaces of $x = (x_{mnk})$ is said to be Riesz convergent to 0. If the triple sequence spaces of $x = (x_{mnk})$ is Riesz convergent to 0, then we write $P_R - \lim x = 0$.

Definition 3.3. The triple sequence $\theta_{i,\ell,j} = \{(m_i, n_\ell, k_j)\}$ is called triple lacunary if there exist three increasing sequences of integers such that

$$m_0 = 0, h_i = m_i - m_{r-1} \to \infty \text{ as } i \to \infty,$$

$$n_0 = 0, \overline{h_\ell} = n_\ell - n_{\ell-1} \to \infty \text{ as } \ell \to \infty,$$

$$k_0 = 0, \overline{h_j} = k_j - k_{j-1} \to \infty \text{ as } j \to \infty.$$

Let $m_{i,\ell,j} = m_i n_\ell k_j, h_{i,\ell,j} = h_i \overline{h_\ell h_j}$, and $\theta_{i,\ell,j}$ is determined by $I_{i,\ell,j} = \{(m,n,k) : m_{i-1} < m < m_i \text{ and } n_{\ell-1} < n \leq n_\ell \text{ and } k_{j-1} < k \leq k_j\}$, $q_k = \frac{m_k}{m_{k-1}}, \overline{q_\ell} = \frac{n_\ell}{n_{\ell-1}}, \overline{q_j} = \frac{k_j}{k_{j-1}}$.

Using the notations of lacunary sequence and Riesz mean for triple sequence spaces. $\theta_{i,\ell,j} = \{(m_i, n_\ell, k_j)\}$ be a triple lacunary sequence and $q_m \overline{q}_n \overline{\overline{q}}_k$ be sequences of positive real numbers such that $Q_{m_i} = \sum_{m \in (0,m_i]} p_{m_i}, Q_{n_\ell} = \sum_{n \in (0,n_\ell]} p_{n_\ell}, Q_{n_j} = \sum_{k \in (0,k_j]} p_{k_j}$ and $H_i = \sum_{m \in (0,m_i]} p_{m_i}, \overline{H} = \sum_{n \in (0,n_\ell]} p_{n_\ell}, \overline{\overline{H}} = \sum_{k \in (0,k_j]} p_{k_j}$. Clearly, $H_i = Q_{m_i} - Q_{m_{i-1}}, \overline{H}_\ell = Q_{n_\ell} - Q_{n_{\ell-1}}, \overline{\overline{H}}_j = Q_{k_j} - Q_{k_{j-1}}$. If the Riesz transformation of triple sequences is RH-regular, and $H_i = Q_{m_i} - Q_{m_{i-1}} \to \infty$ as $i \to \infty, \overline{H} = \sum_{n \in (0,n_\ell]} p_{n_\ell} \to \infty$ as $\ell \to \infty, \overline{\overline{H}} = \sum_{k \in (0,k_j]} p_{k_j} \to \infty$ as $j \to \infty$, then $\theta'_{i,\ell,j} = \{(m_i, n_\ell, k_j)\} = \left\{\left(Q_{m_i} Q_{n_j} Q_{k_k}\right)\right\}$ is a triple lacunary sequence. If the assumptions $Q_r \to \infty$ as $r \to \infty$, $\overline{Q}_s \to \infty$ as $s \to \infty$ and $\overline{\overline{Q}}_t \to \infty$ as $t \to \infty$ may be not enough to obtain the conditions $H_i \to \infty$ as $i \to \infty, \overline{H}_\ell \to \infty$ as $\ell \to \infty$ and $\overline{\overline{H}}_j \to \infty$ as $j \to \infty$ respectively. For any lacunary sequences $(m_i), (n_\ell)$ and (k_j) are integers.

Throughout the paper, we assume that $Q_r = q_{11} + q_{12} + \ldots + q_{rs} + \cdots \to \infty \, (r \to \infty), \overline{Q}_s = \overline{q}_{11} + \overline{q}_{12} + \ldots + \overline{q}_{rs} + \cdots \to \infty \, (s \to \infty), \overline{\overline{Q}}_t = \overline{\overline{q}}_{11} + \overline{\overline{q}}_{12} +$

$\ldots + \overline{\overline{q}}_{rs} + \cdots \to \infty \, (t \to \infty)$, such that $H_i = Q_{m_i} - Q_{m_{i-1}} \to \infty$ as $i \to \infty, \overline{H}_\ell = Q_{n_\ell} - Q_{n_{\ell-1}} \to \infty$ as $\ell \to \infty$ and $\overline{\overline{H}}_j = Q_{k_j} - Q_{k_{j-1}} \to \infty$ as $j \to \infty$.

Let $Q_{m_i,n_\ell,k_j} = Q_{m_i}\overline{Q}_{n_\ell}\overline{\overline{Q}}_{k_j}, H_{i\ell j} = H_i\overline{H}_\ell\overline{\overline{H}}_j$,

$I'_{i\ell j} = \left\{(m,n,k) : Q_{m_{i-1}} < m < Q_{m_i}, \overline{Q}_{n_{\ell-1}} < n < Q_{n_\ell} \text{ and } \overline{Q}_{k_{j-1}} < k < \overline{Q}_{k_j}\right\}$,

$V_i = \frac{Q_{m_i}}{Q_{m_{i-1}}}, \overline{V}_\ell = \frac{Q_{n_\ell}}{Q_{n_{\ell-1}}}$ and $\overline{\overline{V}}_j = \frac{Q_{k_j}}{Q_{k_{j-1}}}$, and $V_{i\ell j} = V_i\overline{V}_\ell\overline{\overline{V}}_j$.

If we take $q_m = 1, \overline{q}_n = 1$ and $\overline{\overline{q}}_k = 1$ for all m, n and k then $H_{i\ell j}, Q_{i\ell j}, V_{i\ell j}$ and $I'_{i\ell j}$ reduce to $h_{i\ell j}, q_{i\ell j}, v_{i\ell j}$ and $I_{i\ell j}$.

4. Almost Lacunary Cesàro $[C,1,1,1]$−statistical convergence of probabilistic space P with triple sequence spaces of χ^3

Let $A = [a^{pqr}_{mnk}]^\infty_{m,n,k=0}$ be a triple infinite matrix of real number for $p, q, r = 1, 2, \ldots$ forming the sum

$$\mu_{pqr}(X) = \sum_{m=0}^\infty \sum_{n=0}^\infty \sum_{k=0}^\infty a^{pqr}_{mnk} \left(\left((m+n+k)! \left(\frac{x_{mnk}}{y_{mnk}}\right)\right)^{1/m+n+k}, \bar{0}\right) \tag{4.1}$$

called the A means of the triple sequence X yielded a method of summability. We say that a sequence X is A summable to the limit 0 of the A mean exist for all $p, q, r = 0, 1, \ldots$ and converges.

$$\lim_{uvw\to\infty} \sum_m^u \sum_n^v \sum_k^w a^{pqr}_{mnk} \left((m+n+k)! \left(\frac{x_{mnk}}{y_{mnk}}\right)\right)^{1/m+n+k} = \mu_{pqr}$$

and

$$\lim_{pqr\to\infty} \mu_{pqr} = 0$$

Define the means

$$\sigma^X_{pqr} = \frac{1}{pqr} \sum_{m=0}^p \sum_{n=0}^q \sum_{k=0}^r \left((m+n+k)! \left(\frac{x_{mnk}}{y_{mnk}}\right)\right)^{1/m+n+k}$$

and

$$A\sigma^X_{pqr} = \frac{1}{pqr} \sum_{m=0}^p \sum_{n=0}^q \sum_{k=0}^r a^{pqr}_{mnk} \left(\left((m+n+k)! \left(\frac{x_{mnk}}{y_{mnk}}\right)\right)^{1/m+n+k}, \bar{0}\right).$$

We say that $\left(\frac{x_{mnk}}{y_{mnk}}\right)$ is statistically lacunary equivalent summable $[C,1,1,1]$ to 0, if the sequence $\sigma = (\sigma^x_{mnk})$ is statistically convergent to $\bar{0}$, that is, $st_3 - \lim_{pqr} \sigma^x_{pqr} = 0$. It is denoted by $[C,1,1,1]\,(st_3)$, the set of all triple sequence which one statistically lacunary equivalent to summable $[C,1,1,1]$.

Let $q_m, \overline{q}_n$ and $\overline{\overline{q}}_k$ be sequences of positive numbers and $Q_r = q_{11} + \cdots q_{rs}$, $\overline{Q}_s = \overline{q}_{11} \cdots \overline{q}_{rs}$ and $\overline{\overline{Q}}_t = \overline{\overline{q}}_{11} \cdots \overline{\overline{q}}_{rs}$. If we choose $q_m = 1, \overline{q}_n = 1$ and $\overline{\overline{q}}_k = 1$ for all m, n and k.

Definition 4.1. A triple $(X, P, *)$ be a probabilistic space. Then a triple sequence spaces $x = (x_{mnk})$ is said to statistically convergent to $\bar{0}$ with respect to the probabilistic, $P-$ provided that for every $\epsilon > 0$ and $\gamma \in (0, 1)$

$$\delta\left(\left\{m, n, k \in \mathbb{N} : P - \lim_{r,s,t\to\infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon)\right] \leq \right.\right.$$

$$\left.\left. \leq 1 - \gamma \right\}\right) = 0$$

or equivalently $\lim_{k\ell v} \frac{1}{k\ell v} m \leq k, n \leq \ell, k \leq v :$

$$P - \lim_{r,s,t\to\infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon)\right] \leq 1 - \gamma = 0$$

In this case we write $St_P - \lim_x = \bar{0}$.

Definition 4.2. A triple sequence spaces of $(X, P, *)$ be a probabilistic space P. The two non-negative triple sequence spaces of $x = (x_{mnk})$ and $y = (y_{mnk})$ are said to be almost asymptotically statistical equivalent of multiple $\bar{0}$ in probabilistic space X if for every $\epsilon > 0$ and $\gamma \in (0, 1)$.

$$\delta\left(\left\{ m, n \in \mathbb{N} : \right.\right.$$

$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] \leq 1 - \gamma \right\}\right) = 0$$

or equivalently

$$\lim_{k\ell} \frac{1}{k\ell} \left|\left\{ m \leq k, n \leq \ell : \right.\right.$$

$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] \leq 1 - \gamma \right\}\right| = 0.$$

In this case we write $x \overset{\widehat{S}(P)}{\equiv} y$.

Definition 4.3. A triple sequence spaces of $(X, P, *)$ be a probabilistic space P and $\theta = (m_r n_s k_t)$ be a triple lacunary sequence spaces are said to be a almost asymptotically lacunary statistical equivalent of multiple $\bar{0}$ in probabilistic space X if for every $\epsilon > 0$ and $\gamma \in (0, 1)$

$$\delta_\theta\left(\left\{ m, n, k \in I_{rst} : \right.\right. \tag{4.2}$$

$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] \leq 1 - \gamma \right\}\right) = 0$$

or equivalently

$$\lim_{rst} \frac{1}{h_{rst}} \Bigg| \Bigg\{ \quad m, n \in I_{rst} :$$

$$P - \lim_{r,s,t \to \infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r} \sum_{n=1}^{s} \sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f \left(A\sigma_{pqr}^{x} \right) (\epsilon), \bar{0} \right] \leq 1 - \gamma \Bigg\} \Bigg| = 0.$$

In this case we write $x \overset{\widehat{S_\theta}(P)}{\equiv} y$.

Lemma 4.4. *A triple sequence spaces of $(X, P, *)$ be a probabilistic space P. Then for every $\epsilon > 0$ and $\gamma \in (0, 1)$, the following statements are equivalent:*

(1) $$\lim_{rst} \frac{1}{h_{rst}} \Bigg| \Bigg\{ \quad m, n, k \in I_{rst} :$$

$$P - \lim_{r,s,t \to \infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r} \sum_{n=1}^{s} \sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f \left(A\sigma_{pqr}^{x} \right) (\epsilon), \bar{0} \right] \leq 1 - \gamma \Bigg\} \Bigg| = 0,$$

(2) $$\delta_\theta \Bigg(\Bigg\{ \quad m, n, k \in I_{rst} :$$

$$P - \lim_{r,s,t \to \infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r} \sum_{n=1}^{s} \sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f \left(A\sigma_{pqr}^{x} \right) (\epsilon), \bar{0} \right] \leq 1 - \gamma \Bigg\} \Bigg) = 0,$$

(3) $$\delta_\theta \Bigg(\Bigg\{ \quad m, n, k \in I_{rst} :$$

$$P - \lim_{r,s,t \to \infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r} \sum_{n=1}^{s} \sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f \left(A\sigma_{pqr}^{x} \right) (\epsilon), \bar{0} \right] \leq 1 - \gamma \Bigg\} \Bigg) = 1,$$

(4) $$\lim_{rst} \frac{1}{h_{rst}} \Bigg| \Bigg\{ \quad m, n, k \in I_{rst} :$$

$$P - \lim_{r,s,t \to \infty} \frac{1}{Q_r \overline{Q}_s \overline{\overline{Q}}_t} \sum_{m=1}^{r} \sum_{n=1}^{s} \sum_{k=1}^{t} q_m \overline{q}_n \overline{\overline{q}}_k \left[f \left(A\sigma_{pqr}^{x} \right) (\epsilon), \bar{0} \right] \leq 1 - \gamma \Bigg\} \Bigg| = 1.$$

5. Main Results

Theorem 5.1. *A triple sequence spaces of $(X, P, *)$ be a probabilistic space P. If two non-negative triple sequence spaces of $x = (x_{mnk})$ and $y = (y_{mnk})$ are almost asympototically lacunary statistical equivalent of multiple $\bar{0}$ with respect to the probabilistic P, then $\bar{0}$ is unique sequence.*

Proof. Assume that $x \overset{\widehat{S_{\theta}^{\bar{0}}}(P)}{\equiv} y$. For a given $\lambda > 0$ choose $\gamma \in (0,1)$ such that $(1-\gamma) > 1-\lambda$. Then, for any $\epsilon > 0$, define the following set:

$$K = \Bigg\{ \quad m,n \in I_{r,s} :$$

$$P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] \leq 1-\gamma \Bigg\}$$

Then, clearly

$$\lim_{rst} \frac{K \bigcap \bar{0}}{h_{rst}} = 1,$$

so K is non-empty set, since $x \overset{\widehat{S_{\theta}^{\bar{0}}}(P)}{\equiv} y, \delta_\theta(K) = 0$ for all $\epsilon > 0$, which implies $\delta_\theta(\mathbb{N} - K) = 1$. If $m,n,k \in \mathbb{N} - K$, then we have

$$\begin{aligned} P_{\bar{0}}(\epsilon) &= P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] \\ &> (1-\gamma) \geq 1-\lambda \end{aligned}$$

since λ is arbitrary, we get $P_{\bar{0}}(\epsilon) = 1$.

This completes the proof. □

Theorem 5.2. *A triple sequence spaces of $(X,P,*)$ be a probabilistic space. For any lacunary sequence $\theta = (m_r n_s k_t)$, $\widehat{S}_\theta(P) \subset \widehat{S}(P)$ if $\limsup_{rst} q_{rst} < \infty$.*

Proof. If $\limsup_{rst} q_{rst} < \infty$. then there exists a $B > 0$ such that $q_{rst} < B$ for all $r,s,t \geq 1$. Let $x \overset{\widehat{S_{\theta}}(P)}{\equiv} y$ and $\epsilon > 0$. Now we have to prove $\widehat{S}(P)$. Set

$$K_{rst} = \Bigg|\Bigg\{ \quad m,n,k \in I_{rst} :$$

$$P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] 1-\gamma \Bigg\}\Bigg|.$$

Then by definition, for given $\epsilon > 0$, there exists $r_0 s_0 t_0 \in \mathbb{N}$ such that

$$\frac{K_{rst}}{h_{rst}} < \frac{\epsilon}{2B} \text{ for all } r > r_0,\ s > s_0 \text{ and } t > t_0.$$

Let

$$M = \max\{K_{rs} :\ 1 \leq r \leq r_0,\ 1 \leq s \leq s_0,\ 1 \leq t \leq t_0\}$$

and let uvw be any positive integer with $m_{r-1} < u \leq m_r, n_{s-1} < v \leq n_s$ and $k_{t-1} < w \leq k_t$. Then

$$\frac{1}{uvw}\left|\left\{ m \leq u, n \leq v, k \leq w : \right.\right.$$
$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] > 1-\gamma\right\}\right|$$
$$\leq \frac{1}{m_{r-1}n_{s-1}k_{t-1}}\left|\left\{ m \leq m_r, n \leq n_s, k \leq k_t : \right.\right.$$
$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] > 1-\gamma\right\}\right|$$
$$= \frac{1}{m_{r-1}n_{s-1}k_{t-1}}\left\{K_{111} + \ldots + K_{rst}\right\}$$
$$\leq \frac{M}{m_{r-1}n_{s-1}k_{t-1}} r_0s_0t_0 + \frac{\epsilon}{2B}q_{rst} \leq \frac{M}{m_{r-1}n_{s-1}k_{t-1}} r_0s_0t_0 + \frac{\epsilon}{2}.$$

This completes the proof. □

Theorem 5.3. *A triple sequence spaces of* $(X, P, *)$ *be a probabilistic space* P. *For any lacunary sequence* $\theta = (m_rn_sk_t)$, $\widehat{S}(P) \subset \widehat{S}_\theta(P)$ *if* $\liminf_{rst} q_{rst} > 1$.

Proof. If $\liminf_{rst} q_{rst} > 1$, then there exists a $\beta > 0$ such that $q_{rst} > 1+\beta$ for sufficiently large rst, which implies

$$\frac{h_{rst}}{K_{rst}} \geq \frac{\beta}{1+\beta}.$$

Let $x \overset{\widehat{S^{\bar{0}}}(p)}{\equiv} y$, then for every $\epsilon > 0$ and for sufficiently large r, s, t we have

$$\frac{1}{m_rn_sk_t}\left|\left\{ m \leq m_r, n \leq n_s, k \leq k_t : \right.\right.$$
$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] > 1-\gamma\right\}\right|$$
$$\geq \frac{1}{m_rn_sk_t}\left|\left\{ m, n, k \in I_{rst} : \right.\right.$$
$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] > 1-\gamma\right\}\right|$$
$$\geq \frac{\beta}{1+\beta}\frac{1}{h_{rst}}\left|\left\{ m, n, k \in I_{rst} : \right.\right.$$
$$\left.\left. P - \lim_{r,s,t\to\infty} \frac{1}{Q_r\overline{Q}_s\overline{\overline{Q}}_t} \sum_{m=1}^{r}\sum_{n=1}^{s}\sum_{k=1}^{t} q_m\overline{q}_n\overline{\overline{q}}_k \left[f\left(A\sigma^x_{pqr}\right)(\epsilon), \bar{0}\right] > 1-\gamma\right\}\right|.$$

Therefore $x \overset{\widehat{S^{0}_{\theta}}(p)}{\equiv} y$. This completes the proof. □

Corollary 5.4. *A triple sequence spaces of* $(X, P, *)$ *be a probabilistic space* P *. For any lacunary sequence* $\theta = (m_r n_s k_t)$, *with* $1 < \liminf_{rst} q_{rst} \leq \limsup_{rst} q_{rst} < \infty$, *then* $\widehat{S}(P) = \widehat{S}_{\theta}(P)$.

Proof. The result clearly follows from Theorem 4.2 and Theorem 4.3. □

6. Acknowledgements

The authors are extremely grateful to the anonymous learned referee(s) for their keen reading, valuable suggestion and constructive comments for the improvement of the manuscript. The authors are thankful to the editor(s) and reviewers of Acta Mathematica Academiae Paedagogicae Nyíregyháziensis.

The present paper was completed during a visit by Professor N. Subramanian to Tripura (A central) University (May-June,2016). The second author is very grateful to the Tripura (A Central) University for providing him hospitality. The research was supported by INSA (Indian National Science Academy visiting fellowship) while the second author was visiting Tripura (A central) University under the INSA visiting fellowship.

References

[1] A. Esi , On some triple almost lacunary sequence spaces defined by Orlicz functions, *Research and Reviews:Discrete Mathematical Structures*, **1(2)**, (2014), 16-25.
[2] A. Esi and M. Necdet Catalbas,Almost convergence of triple sequences, *Global Journal of Mathematical Analysis*, **2(1)**, (2014), 6-10.
[3] A. Esi and E. Savas, On lacunary statistically convergent triple sequences in probabilistic normed space,*Appl.Math.and Inf.Sci.*, **9 (5)** , (2015), 2529-2534.
[4] A. J. Datta A. Esi and B.C. Tripathy,Statistically convergent triple sequence spaces defined by Orlicz function , *Journal of Mathematical Analysis*, **4(2)**, (2013), 16-22.
[5] S. Debnath, B. Sarma and B.C. Das ,Some generalized triple sequence spa5ces of real numbers , *Journal of nonlinear analysis and optimization*, **Vol. 6, No. 1** (2015), 71-79.
[6] *H. Kizmaz*, On certain sequence spaces, *Canadian Mathematical Bulletin,* **24(2)**(1981), 169-176.
[7] P.K. Kamthan and M. Gupta, Sequence spaces and series, Lecture notes, Pure and Applied Mathematics, *65 Marcel Dekker, In c. New York* , 1981.
[8] J. Lindenstrauss and L. Tzafriri, On Orlicz sequence spaces, *Israel J. Math.*, **10** (1971), 379-390.
[9] J. Musielak, Orlicz Spaces,Lectures Notes in Math.,1034, *Springer-Verlag*, **1983**.
[10] A. Sahiner, M. Gurdal and F.K. Duden, Triple sequences and their statistical convergence, *Selcuk J. Appl. Math.* , **8 No. (2)**(2007), 49-55.
[11] A. Sahiner, B.C. Tripathy , Some I related properties of triple sequences, *Selcuk J. Appl. Math. ,* **9 No. (2)**(2008), 9-18.
[12] N. Subramanian and A. Esi, The generalized tripled difference of χ^3 sequence spaces, *Global Journal of Mathematical Analysis*, **3 (2)** (2015), 54-60.

Received May 22, 2016.

Shyamal Debnath,
Department of Mathematics,
Tripura University (A Central University),
Suryamaninagar, Agartala-799022,West Tripura,
India
E-mail address: debnathshyamal@tripurauniv.in, shyamalnitamath@gmail.com

N. Subramanian,
Department of Mathematics,
SASTRA University,
Thanjavur-613 401,
India
E-mail address: nsmaths@yahoo.com