surfaces of constant mean curvature 1 in $\mathbb{H}^3$. Thus we have a one-parameter family of surfaces of constant mean curvature 1 in $\mathbb{H}^3$ with parameter c. The deformed surfaces might not have finite total curvature, but they will have the same topological type and the same reflectional symmetries as the minimal surface. (See Section 4.)

Regarding item 3 above, Theorem 1.2 shows that the converse of the [Rossman et al. 1997] result does not hold.

3. THE EMBEDDED CASE

The proof of Theorem 1.1 uses the maximum principle and Alexandrov reflection. Before stating the maximum principle, we define some terms. If Σ_1 and Σ_2 are two smooth oriented complete hypersurfaces of $\mathbb{H}^n$ that are tangent at a point p and have the same oriented normal at p, we say that p is a *point of common tangency* for Σ_1 and Σ_2. Let the common tangent geodesic hyperplane $\mathscr{P}$ through p have the same orientation as Σ_1 and Σ_2 at p. Then, near p, expressing Σ_1 and Σ_2 as graphs $g_1(x)$ and $g_2(x)$ over points $x \in \mathscr{P}$ (the term *graph* in this context is defined in [do Carmo and Lawson 1983]), we say that Σ_1 *lies above* Σ_2 near p if $g_1 \geq g_2$.

Proposition 3.1 (Maximum Principle). *Suppose that Σ_1 and Σ_2 are closed oriented hypersurfaces in $\mathbb{H}^n$ with the same constant mean curvature c and the same smooth boundary $\partial\Sigma_1 = \partial\Sigma_2$. Suppose that Σ_1 and Σ_2 have a point p of common tangency, and that Σ_1 lies above Σ_2 near p. (The point p can be either an interior point of both Σ_1 and Σ_2 or a boundary point of both Σ_1 and Σ_2.) Then $\Sigma_1 = \Sigma_2$.*

This proposition is well known, and proofs can be found in [do Carmo and Lawson 1983; Korevaar et al. 1992], and references therein.

In the Poincaré model for $\mathbb{H}^3$, which as we recall is the open unit ball in $\mathbb{R}^3$ with the metric $ds^2 = 4|dx|^2/\big(1-|x|^2\big)^2$, the totally geodesic planes are the intersections of B^3 with spheres and planes in $\mathbb{R}^3$ that meet ∂B^3 orthogonally. We shall use the Poincaré model and these totally geodesic planes in the proof of Theorem 1.1.

Proof of Theorem 1.1. We consider a complete properly embedded CMC surface M in $\mathbb{H}^3$. First we suppose that M has asymptotic boundary consisting of exactly two points. Applying an isometry of $\mathbb{H}^3$ if necessary, we may assume that these two asymptotic points are at the north and south poles $(0,0,\pm 1)$.

Let $\vec{v}$ be a horizontal unit vector in $\mathbb{R}^3$. For $t \in (-1,1)$, let P_t be the totally geodesic plane containing the point $t\vec{v}$ and perpendicular to the line through $\vec{v}$. The plane P_t separates $\mathbb{H}^3$ into two regions: let A_t be the region containing the points $s\vec{v}, s \in (-1,t)$, and let B_t be the region containing the points $s\vec{v}$, for $s \in (t,1)$. Let $(M \cap A_t)'$ be the isometric reflection of $M \cap A_t$ across P_t.

Let t_0 be the largest value t_0 such that for all t less than t_0, $\mathrm{Int}\big((M \cap A_t)'\big)$ and $\mathrm{Int}(M \cap B_t)$ are disjoint. When t is close to -1 or 1, $P_t \cap M$ is empty, so it follows that such a t_0 exists and that $t_0 \in (-1,1)$. It then follows (since M is properly embedded) that there exists a finite point of common tangency between $(M \cap A_{t_0})'$and $M \cap B_{t_0}$, and that one surface lies above the other in a neighborhood of this point of common tangency. The maximum principle implies that $M \cap B_{t_0} = (M \cap A_{t_0})'$. (This is the Alexandrov reflection principle.) Since M has only two ends at the north and south poles, it must be that $t_0 = 0$. Since $\vec{v}$ was an arbitrary horizontal vector, it follows that M is symmetric with respect to any geodesic plane through the north and south poles. Thus it is a surface of revolution.

If the surface M has no ends or only one point in its asymptotic boundary, one can similarly conclude that M is a surface of revolution (sphere or horosphere). □

4. AN IMMERSED COUNTEREXAMPLE

In Sections 5 and 6, we give a rigorous proof of existence of the genus-one catenoid cousin. However, since the proof itself does not enlighten the reader as to why it should exist, we give a motivation in

this section for why we should expect this surface to exist.

As noted in Section 2, it was shown in [Rossman et al. 1997] that for any complete finite total curvature minimal surface in $\mathbb{R}^3$ which satisfies certain conditions, there exists a corresponding one-parameter family of surfaces of constant mean curvature 1 in $\mathbb{H}^3$. So, in this sense, the set of surfaces of constant mean curvature 1 in $\mathbb{H}^3$ is a larger set than the set of minimal surfaces in $\mathbb{R}^3$. We now briefly sketch the ideas behind the result in [Rossman et al. 1997]. We do not describe the method in detail, as the reader can refer to [Rossman et al. 1997].

We start with a given minimal surface in $\mathbb{R}^3$, and thus have a Riemann surface Σ and meromorphic functions f and g given to us by the Weierstrass representation for minimal surfaces in $\mathbb{R}^3$ (Lemma 2.1). We can use this same Σ and f and g in the hyperbolic Weierstrass representation (Lemma 2.2) to produce a surface of constant mean curvature c in $\mathbb{H}^3(-c^2)$ for each real number c.

As $c \to 0$, the Poincaré model for $\mathbb{H}^3(-c^2)$ (a ball in $\mathbb{R}^3$ with Euclidean radius $1/|c|$) converges to Euclidean space $\mathbb{R}^3$, and these surfaces of constant mean curvature c in $\mathbb{H}^3(-c^2)$ converge to the given minimal surface (in the sense of C^∞ uniform convergence on compact sets). Thus, for c close to zero, we can think of compact regions of the surfaces of constant mean curvature c in $\mathbb{H}^3(-c^2)$ as small deformations of compact regions of the given minimal surface in $\mathbb{R}^3$.

If the given minimal surface in $\mathbb{R}^3$ is not simply connected, there is a question about whether the deformed surfaces of constant mean curvature c in $\mathbb{H}^3(-c^2)$ are well-defined. This is the period problem. (The period problem being solvable essentially means that a certain set of equations

$$\mathrm{Per}_j(\lambda_i) = 0$$

can be solved with respect to certain parameters λ_i of the surface. This will be explained in detail in terms of an SU(2) condition in the next section.) The minimal surface is assumed to have a "nondegeneracy" property, as defined in [Rossman et al. 1997]. Since the period problem is nondegenerate and solvable on the minimal surface, and since the period problem changes continuously with respect to c, it can still be solved when c is sufficiently close to 0. Thus, for c sufficiently close to 0, the surfaces of constant mean curvature c in $\mathbb{H}^3(-c^2)$ are well-defined.

Dilating the Poincaré model for $\mathbb{H}^3(-c^2)$ by a factor of $|c|$, as described in Section 2, we produce a one-parameter family of surfaces of constant mean curvature 1 in $\mathbb{H}^3$, with parameter c. This is the method used in [Rossman et al. 1997] to create well-defined non-simply connected surfaces of constant mean curvature 1 in $\mathbb{H}^3$ from non-simply connected minimal surfaces in $\mathbb{R}^3$.

As an example, consider the minimal genus-one trinoid in $\mathbb{R}^3$. As discussed in [Berglund and Rossman 1995], there is a single real parameter λ in the Weierstrass data that can be adjusted to solve the period problem. The period problem is represented by a map $\lambda \in \mathbb{R} \to \mathrm{Per}(\lambda) \in \mathbb{R}$, and to solve the period problem we must show that there exists a value of λ so that $\mathrm{Per}(\lambda) = 0$. We note that the function $\mathrm{Per}(\lambda)$ changes continuously in c. Since the period problem for the minimal genus-one trinoid in $\mathbb{R}^3$ (when c is 0) is solvable and nondegenerate, there exists an interval $(a, b) \in \mathbb{R}$ whose image under the map Per contains an interval about 0. By continuity, if we perturb c slightly away from 0, we still have $0 \in \mathrm{Per}(a, b)$. Thus, for c sufficiently close to zero, there exists a CMC c genus-one trinoid cousin in $\mathbb{H}^3(-c^2)$. Then, by dilating the Poincaré model, we produce a genus-one trinoid cousin of constant mean curvature 1 in $\mathbb{H}^3(-1)$: see Figures 4 and 5.

Now we consider Weierstrass data that would produce a minimal genus-one catenoid is $\mathbb{R}^3$. Again there is a single real parameter λ in the Weierstrass data that can be adjusted, and again the period problem is represented by a map $\lambda \in \mathbb{R} \to \mathrm{Per}(\lambda) \in \mathbb{R}$, and to solve the period problem we must again show that there exists a value of λ so



FIGURE 4. View of an embedded genus-one trinoid cousin in the Poincaré model of hyperbolic space.

that $\mathrm{Per}(\lambda) = 0$. The Weierstrass data is described in the next section. In this case the period problem cannot be solved, since the function Per is always positive. But Per can get arbitrarily close to 0, so the interval $(0, \varepsilon)$ is contained in the image of Per, for some $\varepsilon > 0$. If we perturb c slightly, we expect that the image interval $(0, \varepsilon)$ gets perturbed continuously. We have found by numerical experiment that when c becomes slightly negative, the image interval $(0, \varepsilon)$ changes to an image interval of the form $(\varepsilon_1, \varepsilon_2)$, where $0 < \varepsilon_1 < \varepsilon_2$. Thus, as c becomes negative, the lower endpoint of the image interval moves in a positive direction. If this behavior were nondegenerate at $c = 0$, one would expect that as c is perturbed slightly in a positive direction, the image interval would become of the form $(\varepsilon_1, \varepsilon_2)$, where $\varepsilon_1 < 0 < \varepsilon_2$. We have found by numerical experiment that this is indeed the case. Thus, for slightly positive values of c, we can adjust a real parameter in the Weierstrass data so that the period function Per becomes zero. Then, after dilating the Poincaré model, we have existence of a genus-one catenoid cousin of constant mean curvature 1 in $\mathbb{H}^3(-1)$.

The behavior of the genus-one catenoid as c is perturbed is similar to the behavior of the genus-one trinoid as c is perturbed. As c becomes negative, the genus-one trinoid cousin eventually becomes embedded. If the period problem could be solved for the genus-one catenoid cousin when c is negative, the resulting surface would be embedded; as we know by Theorem 1.1, such a surface cannot exist. But when solving the period problem for $c > 0$, the genus-one catenoid cousin is not embedded, in the same way that the genus-one trinoid



FIGURE 5. Slices of genus-one trinoid cousins along a plane of reflective symmetry. The picture on the left corresponds to the embedded genus-one trinoid cousin of Figure 4, produced by using a negative value for c. The one on the right is an immersed genus-one trinoid cousin, produced by using a positive value for c.

cousin is not embedded for $c > 0$. So this numerical experiment is consistent with Theorem 1.1, and furthermore shows that Theorem 1.1 holds only for embedded surfaces.

In the next sections, we will prove Theorem 1.2. First we describe the Weierstrass representation and the period problem. Then we give the proof. Initially the period problem is three-dimensional. We reduce it by algebraic arguments to a one-dimensional problem, then we show by a numerical calculation and the intermediate value theorem that this reduced problem can be solved. The remainder of the proof is essentially a mathematically rigorous verification that the numerical experiment we conducted is correct. We must give rigorous bounds for both computer round-off error and for error introduced by discretizing the problem. We will show that the errors are sufficiently small to ensure the existence of a solution to the period problem.

5. THE PERIOD PROBLEM FOR THE GENUS 1 CATENOID COUSIN

Consider the Riemann surface $\mathscr{M}_a^2 \subset \mathbb{C}\times(\mathbb{C}\cup\{\infty\})$ defined by the equation

$$(z-1)(z+a)w^2 = (z+1)(z-a),$$

where $a > 1$. Thus $\mathscr{M}_a^2$ is a twice punctured torus. Let $g = w$ and let $f = c/w$, for $c > 0$. (This c is the same as the c described in the previous section, and a is the same as λ in the previous section.) The Riemann surface $\mathscr{M}_a^2$ and meromorphic functions g and f are the Weierstrass data for a genus-one catenoid. Let $F(z,w) \in \mathrm{SL}(2,\mathbb{C})$ satisfy Bryant's equation

$$F^{-1}dF = \begin{pmatrix} g & -g^2 \\ 1 & -g \end{pmatrix} f\,dz$$

with initial condition $F =$ identity at $z = 0$, $w = 1$. Hence $\Phi = F^{-1}\overline{F^{-1}}^t$ is a surface of constant mean curvature 1 in the Hermitian model for $\mathbb{H}^3$, and this surface is defined on the universal cover of $\mathscr{M}_a^2$. Representing Φ in this way, we have already done the dilation of hyperbolic space that produces a surface of constant mean curvature 1 in $\mathbb{H}^3(-1)$ from a surface of constant mean curvature c in $\mathbb{H}^3(-c^2)$.

We don't yet know that Φ is well-defined on $\mathscr{M}_a^2$ itself (which must be the case if Φ has finite total curvature). For this to happen, F must satisfy the SU(2) condition, which we now state. Suppose that γ is a loop in $\mathscr{M}_a^2$ with base point $p \in \mathscr{M}_a^2$. Suppose that the value of F at p is $F(p)$. Starting with the initial condition $F(p)$ and evaluating F along γ using Bryant's equation above, we return to the base point p with a new value $\breve{F}(p)$ for F at p. If the loop γ is nontrivial, we can expect that $\breve{F}(p) \neq F(p)$. However, since both $\breve{F}(p)$ and $F(p)$ are in $\mathrm{SL}(2,\mathbb{C})$, there exists a matrix $P \in \mathrm{SL}(2,\mathbb{C})$ such that $\breve{F}(p) = P \cdot F(p)$. If $P \in \mathrm{SU}(2)$, it follows that

$$\breve{F}^{-1}\,\overline{\breve{F}^{-1}}^t = F^{-1}\,\overline{F^{-1}}^t.$$

Thus if $P \in \mathrm{SU}(2)$ for any loop γ, then Φ is well-defined on $\mathscr{M}_a^2$ itself. We say that the SU(2) condition is satisfied on γ if $P \in \mathrm{SU}(2)$.

It is enough to check the SU(2) condition on the following three loops, since they generate the fundamental group of $\mathscr{M}_a^2$ (see Figure 6):

The curve $\gamma_1 \subset \mathscr{M}_a^2$ starts at $(0,1) \in \mathscr{M}_a^2$. Its first portion has z coordinate in the first quadrant of the z plane and ends at a point (z,w) where $z \in \mathbb{R}$ and $1 < z < a$. Its second portion starts at (z,w) and ends at $(0,-1)$ and has z coordinate



FIGURE 6. Projection on the z-plane of the curves γ_1, γ_2, and γ_3, which generate the fundamental group of $\mathscr{M}_a^2$.

in the second quadrant. Its third portion starts at $(0,-1)$ and ends at $(-z,\ 1/\bar{w})$ and has z coordinate in the third quadrant. Its fourth and last portion starts at $(-z,\ 1/\bar{w})$ and returns to the base point $(0,1)$ and has z coordinate in the fourth quadrant.

The curve $\gamma_2 \subset \mathscr{M}_a^2$ starts at $(0,1)$. Its first portion has z coordinate in the first quadrant and ends at a point (z,w) where $z \in \mathbb{R}$ and $z > a$. Its second and last portion starts at (z,w) and returns to $(0,1)$ and has z coordinate in the second quadrant.

The curve $\gamma_3 \subset \mathscr{M}_a^2$ starts at $(0,1)$. Its first portion has z coordinate in the third quadrant and ends at a point (z,w) where $z \in \mathbb{R}$ and $z < -a$. Its second and last portion starts at (z,w) and returns to $(0,1)$ and has z coordinate in the fourth quadrant.

Consider the symmetries

$$\begin{aligned}\varphi_1(z,w) &= (\bar{z},\bar{w}),\\ \varphi_2(z,w) &= (-z,\ 1/w),\\ \varphi_3(z,w) &= (-\bar{z},\ 1/\bar{w})\end{aligned}$$

from $\mathscr{M}_a^2$ to $\mathscr{M}_a^2$. If $(z(t),w(t))$, for $t \in [0,1]$, is a curve in $\mathscr{M}_a^2$ that begins at $(0,1)$ when $t=0$ and ends at some point (z,w) when $t=1$, then we can consider how F changes along $(z(t),w(t))$. At the beginning point of $(z(t),w(t))$ let $F(0,1)$ be the identity, and denote the value of F at the ending point of $(z(t),w(t))$ by $F(z,w)$. Then if we consider the curve $\varphi_i(z(t),w(t))$, F is the identity at the beginning of this curve as well, and we denote the value of F at the end of this curve by $F(\varphi_i(z,w))$. The following lemma gives the relationships between $F(z,w)$ and $F(\varphi_i(z,w))$.

Lemma 5.1. *If* $F(z,w) = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$, *then*

$$\begin{aligned}F(\varphi_1(z,w)) &= \begin{pmatrix} \bar{A} & \bar{B} \\ \bar{C} & \bar{D} \end{pmatrix},\\ F(\varphi_2(z,w)) &= \begin{pmatrix} D & C \\ B & A \end{pmatrix},\\ F(\varphi_3(z,w)) &= \begin{pmatrix} \bar{D} & \bar{C} \\ \bar{B} & \bar{A} \end{pmatrix}.\end{aligned}$$

Proof. Suppose $F(0,1)$ is the identity and $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$ is a solution to the equation

$$\begin{pmatrix} dA & dB \\ dC & dD \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} 1 & -g \\ 1/g & -1 \end{pmatrix} c\,dz$$

on $(z(t),w(t))$. Equivalently,

$$\begin{pmatrix} d\bar{A} & d\bar{B} \\ d\bar{C} & d\bar{D} \end{pmatrix} = \begin{pmatrix} \bar{A} & \bar{B} \\ \bar{C} & \bar{D} \end{pmatrix} \begin{pmatrix} 1 & -\bar{g} \\ 1/\bar{g} & -1 \end{pmatrix} c\,d\bar{z}$$

on $(z(t),w(t))$. Since when $(z,w) \to \varphi_1(z,w)$, we have that $z \to \bar{z}$ and $g \to \bar{g}$, we conclude that $\begin{pmatrix} \bar{A} & \bar{B} \\ \bar{C} & \bar{D} \end{pmatrix}$ is a solution on $\varphi_1(z(t),w(t))$. Since the initial condition that F is the identity is left unchanged by conjugation, we conclude the first part of the lemma. The above equation could also be equivalently written as

$$\begin{pmatrix} dD & dC \\ dB & dA \end{pmatrix} = \begin{pmatrix} D & C \\ B & A \end{pmatrix} \begin{pmatrix} 1 & -1/g \\ g & -1 \end{pmatrix} c\,d(-z).$$

Since when $(z,w) \to \varphi_2(z,w)$, we have $z \to -z$ and $g \to 1/g$, we can conclude the second part of the lemma in the same way. Since $\varphi_3 = \varphi_2 \circ \varphi_1$, the first two parts of the lemma imply the final part. □

In the next lemma, we consider the map $\varphi_4(z,w) = (\bar{z},-\bar{w})$. This map is different from $\varphi_1,\varphi_2,\varphi_3$ in that $(0,1)$ is not in the fixed point set of φ_4. Thus when $(z(t),w(t))$ is a curve that begins at $(0,1)$, the image $\varphi_4(z(t),w(t))$ is a curve that begins at $(0,-1)$, not $(0,1)$.

Lemma 5.2. *Suppose that* $(z(t),w(t)) \subset \mathscr{M}_a^2$ *is a curve that starts at* $(0,1)$ *and ends at a point* (z,w) *such that* $z \in \mathbb{R}$ *and* $1 < z < a$. *Evaluating Bryant's equation along* $(z(t),w(t))$ *with initial condition* $F(0,1) =$ *identity*, *we denote the value of* F *at the endpoint* (z,w) *by* $F(z,w) = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. *Then* $\varphi_4(z(t),w(t))$ *starts at* $(0,-1)$ *and ends at the same endpoint* (z,w). *If we evaluate Bryant's equation along* $\varphi_4(z(t),w(t))$ *with initial condition*

$F(0,-1)$ = identity, then the value of F at the endpoint (z,w) of $\varphi_4(z(t),w(t))$ is

$$F(\varphi_4(z,w)) = \begin{pmatrix} \bar{A} & -\bar{B} \\ -\bar{C} & \bar{D} \end{pmatrix}.$$

Proof. Bryant's equation can be equivalently written as

$$\begin{pmatrix} d\bar{A} & -d\bar{B} \\ -d\bar{C} & d\bar{D} \end{pmatrix} = \begin{pmatrix} \bar{A} & -\bar{B} \\ -\bar{C} & \bar{D} \end{pmatrix} \begin{pmatrix} 1 & \bar{g} \\ -1/\bar{g} & -1 \end{pmatrix} c\, d\bar{z}.$$

The result follows just as in the previous proof. □

Let $\alpha_1(t)$, for $t \in [0,1]$, be a curve starting at (z,w)= $(0,1)$ whose projection to the z-plane is an embedded curve in the first quadrant, and whose endpoint has a z coordinate that is real and larger than 1 and less than a. Let $\alpha_2(t)$, for $t \in [0,1]$, be a curve starting at (z,w)= $(0,1)$ whose projection to the z-plane is an embedded curve in the first quadrant, and whose endpoint has a z coordinate that is real and larger than a. With F = identity at $(z,w) = (0,1)$, we solve Bryant's equation along these two paths to find

$$F(\alpha_1(1)) = \begin{pmatrix} A_1 & B_1 \\ C_1 & D_1 \end{pmatrix},$$
$$F(\alpha_2(1)) = \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix}.$$

Then traveling about the loop γ_1, it follows from Lemmas 5.1 and 5.2 that F changes from the identity to the matrix φ given by

$$\begin{pmatrix} A_1 & B_1 \\ C_1 & D_1 \end{pmatrix} \begin{pmatrix} \bar{D}_1 & \bar{B}_1 \\ \bar{C}_1 & \bar{A}_1 \end{pmatrix} \begin{pmatrix} D_1 & -C_1 \\ -B_1 & A_1 \end{pmatrix} \begin{pmatrix} \bar{A}_1 & -\bar{C}_1 \\ -\bar{B}_1 & \bar{D}_1 \end{pmatrix}.$$

Traveling about the loop γ_2, it follows from Lemma 5.1 that F changes from the identity to the matrix

$$\psi := \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix} \begin{pmatrix} \bar{D}_2 & -\bar{B}_2 \\ -\bar{C}_2 & \bar{A}_2 \end{pmatrix}.$$

And traveling about γ_3, F changes from the identity to the matrix

$$\begin{pmatrix} D_2 & C_2 \\ B_2 & A_2 \end{pmatrix} \begin{pmatrix} \bar{A}_2 & -\bar{C}_2 \\ -\bar{B}_2 & \bar{D}_2 \end{pmatrix}.$$

Changing the initial condition from $F(0,1) =$ identity to

$$F(0,1) = \begin{pmatrix} \alpha & \beta \\ \beta & \alpha \end{pmatrix},$$

where $\alpha, \beta \in \mathbb{R}$, $\alpha^2 - \beta^2 = 1$, we see that solving the SU(2) conditions on all three loops γ_1, γ_2, and γ_3 is equivalent to showing that

$$\begin{pmatrix} \alpha & \beta \\ \beta & \alpha \end{pmatrix} \varphi \begin{pmatrix} \alpha & -\beta \\ -\beta & \alpha \end{pmatrix}$$

and

$$\begin{pmatrix} \alpha & \beta \\ \beta & \alpha \end{pmatrix} \psi \begin{pmatrix} \alpha & -\beta \\ -\beta & \alpha \end{pmatrix}$$

are both in SU(2). We can choose α and β so that this holds precisely when

$$f_1 := \frac{-2(\bar{A}_1 D_1 + \bar{D}_1 A_1 + \bar{C}_1 B_1 + \bar{B}_1 C_1)}{\bar{D}_1 C_1 + \bar{C}_1 D_1 + \bar{B}_1 A_1 + \bar{A}_1 B_1}$$
$$= \frac{2(\bar{A}_2 D_2 - \bar{D}_2 A_2 + \bar{C}_2 B_2 - \bar{B}_2 C_2)}{\bar{D}_2 C_2 - \bar{C}_2 D_2 + \bar{B}_2 A_2 - \bar{A}_2 B_2} =: f_2,$$

and the absolute value of this number is greater than 2. If this holds, we choose α and β so that

$$f_1 = \frac{1 + 2\beta^2}{\beta\sqrt{1+\beta^2}} = f_2,$$

and then the SU(2) conditions are satisfied.

In order to prove Theorem 1.2, we need to show there exist values c and a so that $c > 0$, $a > 1$, $|f_1| = |f_2| > 2$, and $f_1 = f_2$. In this next section we check that such values for c and a exist, by doing a mathematically rigorous analysis of the error bounds for our numerical approximations. (See Figure 7.)

6. ERROR ESTIMATES

Here we shall prove that for some given value of $a > 1$ there exists a positive value for c so that $f_1 = f_2 > 2$. We do this by showing that for one particular value for a, there exists a value of $c > 0$, call it c_1, so that $f_1 > f_2$ at c_1, and there exists another value of $c > c_1$, call it c_2, so that $f_1 < f_2$ at c_2. We also show that $f_1, f_2 \in (2, +\infty)$ for all



FIGURE 7. The functions f_1 (thin curve) and f_2 (thick curve) when $a = 1.78$. The horizontal axis represents c, and the vertical axis represents f_1 and f_2. We see that $f_1, f_2 > 2$ for $c \in (0.0495, 0.0505)$, and $f_1 = f_2$ at some value of c, and $a > 1$.

values of $c \in (c_1, c_2)$. Then, by the continuity of f_i and the intermediate value theorem, we conclude that there exists a $c \in [c_1, c_2]$ such that $f_1 = f_2 > 2$.

Furthermore, by continuity, for any other value of a sufficiently close to our chosen value of a, there also exists a positive value for c so that $f_1 = f_2 > 2$, and hence the genus-one catenoid exists for all a sufficiently close to our chosen value of a. Thus, with our method, showing existence for one value of a is sufficient to conclude the existence of a one-parameter family of genus-one catenoid cousins. (However, we cannot draw any conclusions about the possible range of the parameter a for this one-parameter family.)

Thus, to prove Theorem 1.2, it is sufficient to do the following:

- We choose a suitable value for a and call it a_0. Then we choose suitable values for $c_1 > 0$ and $c_2 > c_1$.
- Using the initial condition F=identity, and the value a_0 for a, and using the value c_i for c, we evaluate the Runge–Kutta algorithm approximation for the solution to Bryant's equation along the path $\alpha_j(t)$. With each evaluation of the Runge–Kutta algorithm, we make the evaluation by both rounding each mathematical operation upward and rounding each mathematical operation downward. Thus for each output of the algorithm, we can find a range in which the theoretical value of the output of the algorithm must lie.
- We then use Lemma 6.1, which gives an upper bound on the absolute value of the difference between the theoretical value of the output of the algorithm and the actual value of the solution of Bryant's equation. Using Lemma 6.1, we can find a single bound which is valid for $a = a_0$ and all $c \in [c_1, c_2]$.
- We then have enough information to determine that any possible approximation errors are small enough to ensure that $f_1 > f_2$ at c_1 and that $f_1 < f_2$ at c_2.
- Then, it only remains to show that f_1 and f_2 are both bounded and greater than 2 for all $c \in [c_1, c_2]$. We do this by showing that the derivative with respect to c of the theoretical value of the output of the algorithm is bounded by a certain constant, for all $c \in [c_1, c_2]$. This is the purpose of Lemma 6.2; it allows us to place limits on the rate at which the output of the algorithm can change with respect to c. This enables us to conclude that $2 < f_i < \infty$ for all $c \in [c_1, c_2]$ simply by checking that this is so at a finite number of values of c in $[c_1, c_2]$.

Lemma 6.1. *Let $\alpha(t)$, for $t \in [0, 1]$, be a path in the complex plane. Let*

$$F(\alpha(t)) = \begin{pmatrix} A(t) & B(t) \\ C(t) & D(t) \end{pmatrix}$$

be an $\mathrm{SL}(2, \mathbb{C})$-valued function on $\alpha(t)$ such that $F(\alpha(0))$=identity and $F(\alpha(t))$ satisfies the equation

$$\begin{pmatrix} dA/dt & dB/dt \\ dC/dt & dD/dt \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} ch_1 & ch_3 \\ ch_2 & ch_4 \end{pmatrix},$$

where c is a real positive constant and h_i are functions on the complex plane satisfying the bounds $|h_i| < M$, $|h_i'| < M_1$, $|h_i''| < M_2$, $|h_i'''| < M_3$ on

$\alpha(t)$ *for* $i = 1, 2, 3, 4$. *Assume also that* h_1 *and* h_4 *are constant functions, and choose* $n \in \mathbb{Z}^+$ *so that* $Mc/n < 0.01$. *Applying the standard Runge–Kutta algorithm on the interval* $t \in [0, 1]$, *using* n *intervals of equal length, let the resulting approximate value for* $F(\alpha(1))$ *produced by the Runge–Kutta algorithm be denoted by the matrix*

$$\begin{pmatrix} \tilde{A} & \tilde{B} \\ \tilde{C} & \tilde{D} \end{pmatrix}.$$

Then $|A(1)-\tilde{A}|$, $|B(1)-\tilde{B}|$, $|C(1)-\tilde{C}|$, $|D(1)-\tilde{D}|$ *are all bounded by*

$$\frac{e^{2.1cM} + e^{4.1cM}}{4.2\, cM\, n^{12}}\,\zeta,$$

where $\zeta = \zeta(c, n, M, M_1, M_2, M_3)$ *is a certain polynomial in six variables.*

We include the condition that h_1 and h_4 are constant in Lemma 6.1, because this is sufficient for our application, and this will later allow us to assume a smaller lower bound for n. It also simplifies the proof somewhat. However, it is not necessary to assume h_1 and h_4 are constant in order to produce a lemma of this type.

Proof. The system of equations in the lemma can be separated into two systems of two equations each: one the system with variables A and B, and the other with C and D. We consider now the system involving A and B:

$$\frac{dA}{dt} = ch_1A + ch_2B, \quad \frac{dB}{dt} = ch_3A + ch_4B.$$

Since $|h_i| < M$ for all $t \in [0, 1]$ and all $i = 1, 2, 3, 4$, we conclude that

$$\left|\frac{dA}{dt}\right| \le cM|A| + cM|B|,$$
$$\left|\frac{dB}{dt}\right| \le cM|A| + cM|B|.$$

If we replace the inequalities in these equations by equalities, we would be able to evaluate the system explicitly with $A(0) = 1$ and $B(0) = 0$. It follows that

$$|A(t)| \le 1 + \frac{1}{2}\sum_{j=1}^{\infty}\frac{(2tcM)^j}{j!} = \tfrac{1}{2} + \tfrac{1}{2}e^{2tcM},$$
$$|B(t)| \le \frac{1}{2}\sum_{j=1}^{\infty}\frac{(2tcM)^j}{j!} = -\tfrac{1}{2} + \tfrac{1}{2}e^{2tcM}.$$

Now we run the standard Runge–Kutta algorithm on $t \in [0, 1]$ for a system of two equations with n steps of equal size $\frac{1}{n}$. The initial conditions are $A_0 = 1$ and $B_0 = 0$. The algorithm at step k is this:

$$k_0 = \tfrac{c}{n}\big(h_1\big(\tfrac{k}{n}\big)A_k + h_2\big(\tfrac{k}{n}\big)B_k\big),$$
$$m_0 = \tfrac{c}{n}\big(h_3\big(\tfrac{k}{n}\big)A_k + h_4\big(\tfrac{k}{n}\big)B_k\big),$$
$$k_1 = \tfrac{c}{n}\big(h_1\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(A_k+\tfrac{1}{2}k_0\big) + h_2\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(B_k+\tfrac{1}{2}m_0\big)\big),$$
$$m_1 = \tfrac{c}{n}\big(h_3\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(A_k+\tfrac{1}{2}k_0\big) + h_4\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(B_k+\tfrac{1}{2}m_0\big)\big),$$
$$k_2 = \tfrac{c}{n}\big(h_1\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(A_k+\tfrac{1}{2}k_1\big) + h_2\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(B_k+\tfrac{1}{2}m_1\big)\big),$$
$$m_2 = \tfrac{c}{n}\big(h_3\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(A_k+\tfrac{1}{2}k_1\big) + h_4\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(B_k+\tfrac{1}{2}m_1\big)\big),$$
$$k_3 = \tfrac{c}{n}\big(h_1\big(\tfrac{k}{n}+\tfrac{1}{n}\big)\big(A_k+k_2\big) + h_2\big(\tfrac{k}{n}+\tfrac{1}{n}\big)\big(B_k+m_2\big)\big),$$
$$m_3 = \tfrac{c}{n}\big(h_3\big(\tfrac{k}{n}+\tfrac{1}{n}\big)\big(A_k+k_2\big) + h_4\big(\tfrac{k}{n}+\tfrac{1}{n}\big)\big(B_k+m_2\big)\big),$$
$$A_{k+1} = A_k + \tfrac{1}{6}\big(k_0+2k_1+2k_2+k_3\big),$$
$$B_{k+1} = B_k + \tfrac{1}{6}\big(m_0+2m_1+2m_2+m_3\big).$$

We define the local discretization errors for A and B to be

$$d^A_{k+1} := A\big(\tfrac{k+1}{n}\big) - A\big(\tfrac{k}{n}\big) - \tfrac{1}{6}\big(\hat{k}_0+2\hat{k}_1+2\hat{k}_2+\hat{k}_3\big),$$
$$d^B_{k+1} := B\big(\tfrac{k+1}{n}\big) - B\big(\tfrac{k}{n}\big) - \tfrac{1}{6}\big(\hat{m}_0+2\hat{m}_1+2\hat{m}_2+\hat{m}_3\big),$$

where

$$\hat{k}_0 = \tfrac{c}{n}\big(h_1\big(\tfrac{k}{n}\big)A\big(\tfrac{k}{n}\big) + h_2\big(\tfrac{k}{n}\big)B\big(\tfrac{k}{n}\big)\big),$$
$$\hat{m}_0 = \tfrac{c}{n}\big(h_3\big(\tfrac{k}{n}\big)A\big(\tfrac{k}{n}\big) + h_4\big(\tfrac{k}{n}\big)B\big(\tfrac{k}{n}\big)\big),$$
$$\hat{k}_1 = \tfrac{c}{n}\big(h_1\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(A\big(\tfrac{k}{n}\big) + \tfrac{1}{2}\hat{k}_0\big) + h_2\big(\tfrac{k}{n}+\tfrac{1}{2n}\big)\big(B\big(\tfrac{k}{n}\big) + \tfrac{1}{2}\hat{m}_0\big)\big),$$

and $\hat{m}_1, \hat{k}_2, \hat{m}_2, \hat{k}_3, \hat{m}_3$ are defined analogously to the way m_1, k_2, m_2, k_3, m_3 were defined. We define the maximums of the local discretization errors by

$$\begin{aligned} D^A &:= \max_k |d_k^A|, \\ D^B &:= \max_k |d_k^B|, \\ D &:= \max(D^A, D^B). \end{aligned}$$

We define the global discretization errors by

$$\begin{aligned} g_k^A &:= A\left(\tfrac{k}{n}\right) - A_k, \\ g_k^B &:= B\left(\tfrac{k}{n}\right) - B_k, \\ g_k &:= \max\left(|g_k^A|, |g_k^B|\right). \end{aligned}$$

Since

$$A\Big(\frac{k+1}{n}\Big) = A\Big(\frac{k}{n}\Big) + \tfrac{1}{6}(\hat{k}_0 + 2\hat{k}_1 + 2\hat{k}_2 + \hat{k}_3) + d_{k+1}^A,$$

we have

$$\begin{aligned} g_{k+1}^A = g_k^A &+ \tfrac{1}{6}(\hat{k}_0 + 2\hat{k}_1 + 2\hat{k}_2 + \hat{k}_3) \\ &- \tfrac{1}{6}(k_0 + 2k_1 + 2k_2 + k_3) + d_{k+1}^A, \end{aligned}$$

and therefore we can compute that

$$\begin{aligned} |g_{k+1}^A| \le\ & |g_k^A| \\ &+ \left(\frac{cM}{n} + \frac{c^2M^2}{n^2} + \frac{2c^3M^3}{3n^3} + \frac{c^4M^4}{3n^4}\right)\left|A\Big(\frac{k}{n}\Big) - A_k\right| \\ &+ \left(\frac{cM}{n} + \frac{c^2M^2}{n^2} + \frac{2c^3M^3}{3n^3} + \frac{c^4M^4}{3n^4}\right)\left|B\Big(\frac{k}{n}\Big) - B_k\right| \\ &+ |d_{k+1}^A|. \end{aligned}$$

We assumed that $n > 100cM$, so

$$|g_{k+1}^A| \le \Big(1 + \frac{1.05cM}{n}\Big)|g_k^A| + \frac{1.05cM}{n}|g_k^B| + D^A.$$

Similarly,

$$|g_{k+1}^B| \le \frac{1.05cM}{n}|g_k^A| + \Big(1 + \frac{1.05cM}{n}\Big)|g_k^B| + D^B.$$

Thus,

$$g_{k+1} \le (1 + \frac{2.1cM}{n})g_k + D.$$

By repeated application of this inequality we have

$$g_n \le \Big(1 + \frac{2.1cM}{n}\Big)^n g_0 + \frac{(1 + 2.1cM/n)^n - 1}{2.1cM/n} D.$$

And since $g_0 = 0$, we have

$$g_n \le \frac{e^{2.1cMn/n} - 1}{2.1cM/n} D < \frac{ne^{2.1cM}}{2.1cM} D.$$

Here we have used the fact that e^x is convex on $\mathbb{R}$, so $1 + x \le e^x$ and therefore also $(1+x)^n \le (e^x)^n = e^{xn}$ for any positive x.

Note that $h_i\big(\frac{k}{n} + \frac{1}{2n}\big)$, $h_i\big(\frac{k}{n} + \frac{1}{n}\big)$, and $A\big(\frac{k}{n} + \frac{1}{n}\big)$ have the following Taylor expansions:

$$\begin{aligned} h_i\Big(\frac{k}{n} + \frac{1}{2n}\Big) &= h_i\Big(\frac{k}{n}\Big) \\ &\quad + \frac{1}{2n}h_i'\Big(\frac{k}{n}\Big) + \frac{1}{8n^2}h_i''\Big(\frac{k}{n}\Big) + \frac{1}{48n^3}h_i'''\Big(\frac{k}{n} + \theta\frac{1}{2n}\Big), \\ h_i\Big(\frac{k}{n} + \frac{1}{n}\Big) &= h_i\Big(\frac{k}{n}\Big) \\ &\quad + \frac{1}{n}h_i'\Big(\frac{k}{n}\Big) + \frac{1}{2n^2}h_i''\Big(\frac{k}{n}\Big) + \frac{1}{6n^3}h_i'''\Big(\frac{k}{n} + \theta\frac{1}{n}\Big), \\ A\Big(\frac{k}{n} + \frac{1}{n}\Big) &= A\Big(\frac{k}{n}\Big) + \frac{1}{n}A'\Big(\frac{k}{n}\Big) \\ &\quad + \frac{1}{2n^2}A''\Big(\frac{k}{n}\Big) + \frac{1}{6n^3}A'''\Big(\frac{k}{n}\Big) + \frac{1}{24n^4}A''''\Big(\frac{k}{n} + \theta\frac{1}{n}\Big), \end{aligned}$$

for appropriate values of $\theta \in [0, 1]$. Here the symbol $'$ denotes derivative with respect to t.

Repeatedly using the equalities $A' = ch_1A + ch_2B$ and $B' = ch_3A + ch_4B$, the above Taylor expansion for $A\big(\frac{k}{n} + \frac{1}{n}\big)$ can be rewritten in a longer form so that it does not contain any terms of the form A', B', A'', B'', A''', B''', A'''', or B''''. Using this longer form for $A\big(\frac{k}{n} + \frac{1}{n}\big)$, and using the above Taylor expansions for $h_i\big(\frac{k}{n} + \frac{1}{2n}\big)$ and $h_i\big(\frac{k}{n} + \frac{1}{n}\big)$, we can make a direct (but long) calculation to determine d_{k+1}^A and d_{k+1}^B in terms of A, B, n, c, h_i, and the derivatives (up to third order) of h_i. These formulas are extremely long, so we do not include them here. However, for each of them we can take the sum of the absolute values of all of the terms, and make the following replacements: $|h_i|$ by its upper bound M, $|h_i'|$ by its upper bound

M_1, $|h_i''|$ by its upper bound M_2, $|h_i'''|$ by its upper bound M_3, and $|A|$ and $|B|$ by their upper bound $\frac{1}{2}(1+e^{2cM})$. We then get upper bounds for $|d_{k+1}^A|$ and $|d_{k+1}^B|$. We can then find that a sufficient upper bound for both $|d_{k+1}^A|$ and $|d_{k+1}^B|$ is

$$D \le \frac{1+e^{2cM}}{2n^{13}}\,\zeta,$$

for an appropriate polynomial ζ. Thus

$$g_n < \frac{ne^{2.1cM}}{2.1cM} D \le \frac{ne^{2.1cM}}{2.1cM}\frac{1+e^{2cM}}{2n^{13}}\,\zeta.$$

An identical argument gives the same conclusion for C and D. □

Lemma 6.2. *Suppose that the conditions of Lemma* 6.1 *hold for all c contained in some interval $[c_1, c_2]$. Then $|\partial\tilde{A}/\partial c|$, $|\partial\tilde{B}/\partial c|$, $|\partial\tilde{C}/\partial c|$, and $|\partial\tilde{D}/\partial c|$ are all bounded by $2.48Me^{2.4Mc}$ for all $c \in [c_1, c_2]$.*

Proof. We consider the system of two equations for A and B here. The case for C and D is identical.

Recall that the Runge–Kutta algorithm is

$$A_{j+1} = A_j + \tfrac{1}{6}(k_0 + 2k_1 + 2k_2 + k_3),$$
$$B_{j+1} = B_j + \tfrac{1}{6}(m_0 + 2m_1 + 2m_2 + m_3).$$

We can expand out the terms of these two equations so that everything is written in terms of only $A_j, B_j, A_{j+1}, B_{j+1}$ and h_i, c, n. We then have

$$A_{j+1} = A_j + ZA_j + ZB_j, \quad B_{j+1} = B_j + ZA_j + ZB_j,$$

where Z is a polynomial that consists of the sum of two terms of the form

$$\frac{ch_*(*)}{6n},$$

two terms of the form

$$\frac{ch_*(*)}{3n},$$

six terms of the form

$$\frac{c^2h_*(*)h_*(*)}{6n^2},$$

eight terms of the form

$$\frac{c^3h_*(*)h_*(*)h_*(*)}{12n^3},$$

and eight terms of the form

$$\frac{c^4h_*(*)h_*(*)h_*(*)h_*(*)}{24n^4}.$$

We will use Z to denote any polynomial of this form, regardless of what the subindices are for the functions $h_i(z)$ and regardless of the value of $z(t)$ at which we are evaluating the functions $h_i(z)$. (It is for this reason that we are writing the functions $h_i(z)$ merely as $h_*(*)$.) Although Z is not a well-defined notation, for our purposes it will be sufficient. It follows from the assumptions $|h_i| < M$ and $c > 0$ and $Mc/n < 0.01$ that

$$|Z| < 1.2\,\frac{Mc}{n} \quad \text{and} \quad \left|\frac{\partial Z}{\partial c}\right| < \frac{M}{n}\left(1 + 2.4\,\frac{Mc}{n}\right),$$

regardless of what the indices of h_i are and regardless of at which values of $z(t)$ we evaluate the functions h_i.

Applying the Runge–Kutta algorithm on n steps of equal length, we find that the resulting estimates for A and B at $\alpha(1)$ are of the form

$$A_n = A_0 + p(Z)A_0 + p(Z)B_0,$$
$$B_n = B_0 + p(Z)A_0 + p(Z)B_0,$$

where $p(Z)$ is a polynomial in Z with

$$\frac{2^{j-1}n(n-1)(n-2)\cdots(n-j+1)}{j!}$$

terms of the form Z^j for each $j = 1, \ldots, n$. Thus we have the bounds

$$\left|\frac{\partial A_n}{\partial c}\right|, \left|\frac{\partial B_n}{\partial c}\right| \le 2\sum_{j=1}^{n} \frac{2^{j-1}n(n-1)\cdots(n-j+1)}{j!}\left|\frac{\partial Z^j}{\partial c}\right|$$
$$\le \sum_{j=1}^{n} \frac{2^j n(n-1)\cdots(n-j+1)}{j!}\, j\,|Z|^{j-1}\left|\frac{\partial Z}{\partial c}\right|$$

$$\le \sum_{j=1}^{n} \frac{2^j n(n-1)\cdots(n-j+1)}{j!}\, j \left(1.2\frac{Mc}{n}\right)^{j-1} \times \left(\frac{M}{n}\left(1+2.4\,\frac{Mc}{n}\right)\right)$$

$$\le \sum_{j=1}^{n} \frac{2^j}{(j-1)!}\,(1.2Mc)^{j-1}\left(M\left(1+2.4\,\frac{Mc}{n}\right)\right)$$

$$\le 2M\left(1+2.4\frac{Mc}{n}\right)\sum_{j=1}^{n}\frac{1}{(j-1)!}\,(2.4Mc)^{j-1}$$

$$< 2.48M\sum_{j=0}^{n-1}\frac{1}{j!}\,(2.4Mc)^j$$

$$< 2.48M\sum_{j=0}^{\infty}\frac{1}{j!}\,(2.4Mc)^j = 2.48Me^{2.4Mc}. \qquad \square$$

Proof of Theorem 1.2. Recall that our surface is described by the equation

$$\begin{pmatrix} dA/dz & dB/dz \\ dC/dz & dD/dz \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix}\begin{pmatrix} g & -g^2 \\ 1 & -g \end{pmatrix}\frac{c}{g},$$

where

$$g = \sqrt{\frac{(z+1)(z-a)}{(z-1)(z+a)}}.$$

This system separates into two systems: one involving A and B, and the other involving C and D. We now consider the system involving A and B:

$$\frac{dA}{dz} = cA + \frac{c}{g}B, \quad \frac{dB}{dz} = -cgA - cB.$$

If we wish to evaluate this system along a curve from z_0 to z_1, linearly defined as $(b-t)z_0/(b-a)+(t-a)z_1/(b-a)$ for $t \in [a,b]$, then

$$\frac{dz}{dt} = \frac{z_1 - z_0}{b-a}$$

for all $t \in [a,b]$. Our system can then be written as

$$\frac{dA}{dt} = c\,\frac{z_1-z_0}{b-a}\,A + c\,\frac{z_1-z_0}{b-a}\,\frac{1}{g}\,B,$$

$$\frac{dB}{dt} = -c\,\frac{z_1-z_0}{b-a}\,gA - c\,\frac{z_1-z_0}{b-a}\,B.$$

Set

$$h_1 = \frac{z_1-z_0}{b-a}, \qquad h_2 = \frac{z_1-z_0}{b-a}\,\frac{1}{g},$$

$$h_3 = -\frac{z_1-z_0}{b-a}\,g, \quad h_4 = -\frac{z_1-z_0}{b-a}.$$

Now we choose the paths $\alpha_1(t)$ and $\alpha_2(t)$, where $t \in [0,1]$. The paths will start at the point $\alpha_1(0) = \alpha_2(0) = (0,1)$ in the base Riemann surface $\mathscr{M}_a^2$ and will be defined by their z coordinates. The z coordinates of the paths will be polygonal and t will be defined linearly with respect to z-length on each line segment. The path $\alpha_1(t)$ will project to a line segment from $z=0$ $(t=0)$ to $z = 1+0.4i$ $(t=0.67)$, then a line segment from $z = 1+0.4i$ $(t=0.67)$ to $z = \frac{1}{2}(1+a)$ $(t=1)$. The path $\alpha_2(t)$ will project to a line segment from $z=0$ $(t=0)$ to $z = (a+0.2)+0.7i$ $(t=0.686)$, then a line segment from $z=(a+0.2)+0.7i$ $(t=0.686)$ to $z = a+\frac{1}{2}$ $(t=1)$.

We now solve Bryant's differential equation along $\alpha_j(t)$. At the beginning point $(z,w)=(0,1)$, that is, at $t=0$, the initial condition will be $F =$ identity. Suppose that the true value of F at the endpoints is

$$F(\alpha_j(1)) = \begin{pmatrix} A_j & B_j \\ C_j & D_j \end{pmatrix}, \quad \text{for } j = 1,2,$$

and that the approximate value of F at the endpoints produced by the Runge–Kutta algorithm using n steps of equal length is

$$\begin{pmatrix} \tilde{A}_j & \tilde{B}_j \\ \tilde{C}_j & \tilde{D}_j \end{pmatrix}, \quad \text{for } j = 1,2.$$

Of course, the exact values of $\tilde{A}_j, \tilde{B}_j, \tilde{C}_j, \tilde{D}_j$ cannot be computed, but by considering the possible round-off error for each mathematical operation in the algorithm, and keeping track of the possible cumulative round-off error, we can find intervals in which they must lie. That is, we can find real numbers $\tilde{A}_j^{ur}$, $\tilde{A}_j^{ui}$, $\tilde{A}_j^{lr}$, $\tilde{A}_j^{li}$, $\tilde{B}_j^{ur}$, $\tilde{B}_j^{ui}$, $\tilde{B}_j^{lr}$, $\tilde{B}_j^{li}$, $\tilde{C}_j^{ur}$, $\tilde{C}_j^{ui}$, $\tilde{C}_j^{lr}$, $\tilde{C}_j^{li}$, $\tilde{D}_j^{ur}$, $\tilde{D}_j^{ui}$, $\tilde{D}_j^{lr}$, $\tilde{D}_j^{li}$ (where ur stands

for the upper bound of the real part, and so on) such that

$$\tilde{T}_j^{lr} \le \operatorname{Re}\tilde{T}_j \le \tilde{T}_j^{ur},$$
$$\tilde{T}_j^{li} \le \operatorname{Im}\tilde{T}_j \le \tilde{T}_j^{ui},$$

for T taking the values A, B, C, D and $j = 1, 2$. These bounds can be computed using code written in a programming language such as C++ or Fortran. They also could be computed using a shorter code written in a scientific programming language such as CXSC or PROFIL.

Choosing a to be 1.78, we find that we have the following bounds for both paths: $M = 4.6$, $M_1 = 48$, $M_2 = 850$, $M_3 = 25000$. (These bounds are defined in Lemma 6.1.) Then, if we choose any $c \in [0.0495, 0.0505]$, from Lemma 6.1 we see that if $n > 500$, the errors incurred by the Runge–Kutta algorithm on A_j, B_j, C_j, and D_j are less than ε=0.00001. That is, $|\tilde{A}_j - A_j| < \varepsilon$, $|\tilde{B}_j - B_j| < \varepsilon$, $|\tilde{C}_j - C_j| < \varepsilon$, and $|\tilde{D}_j - D_j| < \varepsilon$, for all $c \in [0.0495, 0.0505]$. It follows that

$$\tilde{T}_j^{lr} - \varepsilon \le \operatorname{Re}(T_j) \le \tilde{T}_j^{ur} + \varepsilon,$$
$$\tilde{T}_j^{li} - \varepsilon \le \operatorname{Im}(T_j) \le \tilde{T}_j^{ui} + \varepsilon,$$

for $T = A, B, C, D$ and $j = 1, 2$.

The value of a is fixed, but c is arbitrary in the range $[0.0495, 0.0505]$, so the numbers $A_j, \ldots, D_j$, $\tilde{A}_j, \ldots, \tilde{D}_j$, $\tilde{A}_j^{ur}, \ldots, \tilde{D}_j^{li}$ are all functions of c. To show this dependence, we write $A_j(c), \ldots, D_j(c)$, $\tilde{A}_j(c), \ldots, \tilde{D}_j(c)$, $\tilde{A}_j^{ur}(c), \ldots, \tilde{D}_j^{li}(c)$.

By Lemma 6.2, the numbers

$$\left|\frac{\partial \tilde{A}_j(c)}{\partial c}\right|, \quad \left|\frac{\partial \tilde{B}_j(c)}{\partial c}\right|, \quad \left|\frac{\partial \tilde{C}_j(c)}{\partial c}\right|, \quad \left|\frac{\partial \tilde{D}_j(c)}{\partial c}\right|$$

are all bounded by $2.48Me^{2.4Mc} < 20$ for all $c \in [0.0495, 0.0505]$. It follows that, if we choose any $c \in [0.04999, 0.05001]$, then $\tilde{A}_j(c)$, $\tilde{B}_j(c)$, $\tilde{C}_j(c)$, and $\tilde{D}_j(c)$ can vary from their values at $c = 0.05$ by at most $\hat{\varepsilon}$ =0.0002. That is, $|\tilde{A}_j(c) - \tilde{A}_j(0.05)| < \hat{\varepsilon}$, $|\tilde{B}_j(c) - \tilde{B}_j(0.05)| < \hat{\varepsilon}$, $|\tilde{C}_j(c) - \tilde{C}_j(0.05)| < \hat{\varepsilon}$, and $|\tilde{D}_j(c) - \tilde{D}_j(0.05)| < \hat{\varepsilon}$, for all $c \in [0.04999, 0.05001]$. We conclude that

$$\tilde{T}_j^{lr}(0.05) - \varepsilon - \hat{\varepsilon} \le \operatorname{Re} T_j(c) \le \tilde{T}_j^{ur}(0.05) + \varepsilon + \hat{\varepsilon},$$
$$\tilde{T}_j^{li}(0.05) - \varepsilon - \hat{\varepsilon} \le \operatorname{Im} T_j(c) \le \tilde{T}_j^{ui}(0.05) + \varepsilon + \hat{\varepsilon},$$

for $T = A, B, C, D$ and $j = 1, 2$ and all c in the interval $[0.04999, 0.05001]$. This is sufficient to conclude that both $f_1, f_2 \in (2, +\infty)$ for all c in the same interval. Checking in this way on many small intervals (a finite number of intervals), we can conclude that $2 < f_1, f_2 < \infty$ for all $c \in [0.0495, 0.0505]$.

Then, as we saw before, solving the period problem means solving $f_1 = f_2 > 2$. Running the Runge–Kutta algorithm with $c = 0.0495$, we conclude that

$$\tilde{T}_j^{lr}(0.0495) - \varepsilon \le \operatorname{Re} T_j(0.0495) \le \tilde{T}_j^{ur}(0.0495) + \varepsilon,$$
$$\tilde{T}_j^{li}(0.0495) - \varepsilon \le \operatorname{Im} T_j(0.0495) \le \tilde{T}_j^{ui}(0.0495) + \varepsilon,$$

for $T = A, B, C, D$ and $j = 1, 2$. These estimates are sufficient to show that $f_1 > f_2$ at $c = 0.0495$. Similarly we can show that $f_1 < f_2$ at $c = 0.0505$. We conclude that there exists a value of $c \in [0.0495, 0.0505]$ so that $f_1 = f_2 > 2$.

We have thus shown of existence of at least one genus-one catenoid cousin. Then, since the problem is continuous in a, we know that for all a sufficiently close to 1.78 there exists a positive value for c so that $f_1 = f_2 > 2$. This proves existence of a one-parameter family of genus-one catenoid cousins. □

REFERENCES

[Berglund and Rossman 1995] J. Berglund and W. Rossman, "Minimal surfaces with catenoid ends", *Pacific J. Math.* **171**:2 (1995), 353–371.

[Bryant 1987] R. L. Bryant, "Surfaces of mean curvature one in hyperbolic space", pp. 321–347 in *Théorie des variétés minimales et applications* (Palaiseau, 1983–1984), Astérisque **154-155**, Soc. math. France, Paris, 1987.

[do Carmo and Lawson 1983] M. P. do Carmo and J. Lawson, H. Blaine, "On Alexandrov–Bernstein

theorems in hyperbolic space", *Duke Math. J.* **50**:4 (1983), 995–1003.

[do Carmo et al. 1986] M. P. do Carmo, J. Gomes, and G. Thorbergsson, "The influence of the boundary behaviour on hypersurfaces with constant mean curvature in H^{n+1}", *Comment. Math. Helv.* **61**:3 (1986), 429–441.

[Hass et al. 1995] J. Hass, M. Hutchings, and R. Schlafly, "The double bubble conjecture", *Electron. Res. Announc. Amer. Math. Soc.* **1**:3 (1995), 98–102.

[Kapouleas 1990] N. Kapouleas, "Complete constant mean curvature surfaces in Euclidean three-space", *Ann. of Math.* (2) **131**:2 (1990), 239–330.

[Karcher 1989] H. Karcher, "The triply periodic minimal surfaces of Alan Schoen and their constant mean curvature companions", *Manuscripta Math.* **64**:3 (1989), 291–357.

[Karcher et al. 1988] H. Karcher, U. Pinkall, and I. Sterling, "New minimal surfaces in S^3", *J. Differential Geom.* **28**:2 (1988), 169–185.

[Korevaar et al. 1989] N. J. Korevaar, R. Kusner, and B. Solomon, "The structure of complete embedded surfaces with constant mean curvature", *J. Differential Geom.* **30**:2 (1989), 465–503.

[Korevaar et al. 1992] N. J. Korevaar, R. Kusner, I. Meeks, William H., and B. Solomon, "Constant mean curvature surfaces in hyperbolic space", *Amer. J. Math.* **114**:1 (1992), 1–43.

[Levitt and Rosenberg 1985] G. Levitt and H. Rosenberg, "Symmetry of constant mean curvature hypersurfaces in hyperbolic space", *Duke Math. J.* **52**:1 (1985), 53–59.

[Osserman 1969] R. Osserman, *A survey of minimal surfaces*, Van Nostrand Reinhold Co., New York, 1969. Reprinted by Dover Publications, New York, 1969.

[Polthier 1991] K. Polthier, "New periodic minimal surfaces in H^3", pp. 201–210 in *Workshop on Theoretical and Numerical Aspects of Geometric Variational Problems* (Canberra, 1990), edited by G. Dziuk et al., Proc. Centre Math. Appl. Austral. Nat. Univ. **26**, Austral. Nat. Univ., Canberra, 1991.

[Rossman et al. 1997] W. Rossman, M. Umehara, and K. Yamada, "Irreducible constant mean curvature 1 surfaces in hyperbolic space with positive genus", *Tohoku Math. J.* **49** (1997), 449–484.

[Schoen 1983] R. M. Schoen, "Uniqueness, symmetry, and embeddedness of minimal surfaces", *J. Differential Geom.* **18**:4 (1983), 791–809.

[Umehara and Yamada 1992] M. Umehara and K. Yamada, "A parametrization of the Weierstrass formulae and perturbation of complete minimal surfaces in $\mathbb{R}^3$ into the hyperbolic 3-space", *J. Reine Angew. Math.* **432** (1992), 93–116.

[Umehara and Yamada 1993] M. Umehara and K. Yamada, "Complete surfaces of constant mean curvature 1 in the hyperbolic 3-space", *Ann. of Math.* (2) **137**:3 (1993), 611–638.

[Umehara and Yamada 1996] M. Umehara and K. Yamada, "Surfaces of constant mean curvature c in $\mathbb{H}^3(-c^2)$ with prescribed hyperbolic Gauss map", *Math. Ann.* **304**:2 (1996), 203–224.

[Umehara and Yamada 1997] M. Umehara and K. Yamada, "A duality on CMC-1 surfaces in hyperbolic space, and a hyperbolic analogue of the Osserman inequality", *Tsukuba J. Math.* **21**:1 (1997), 229–237.

Wayne Rossman, Graduate School of Mathematics, School of Engineering, Kyushu University, Fukuoka 812-81, Japan (wayne@math.kyushu-u.ac.jp)

Katsunori Sato, Department of Mathematics and Information Science, Tokyo Institute of Technology, Oh-okayama, Meguro-ku, Tokyo 152, Japan (ksato@is.titech.ac.jp)

Received April 17, 1996; accepted in revised from May 12, 1997