#A04 INTEGERS 9 (2009), 47-52

ON RAPID GENERATION OF $SL_2(\mathbb{F}_Q)$

Jeremy Chapman
Department of Mathematics, University of Missouri, Columbia, Missouri 65211
jeremy@math.missouri.edu

Alex Iosevich
Department of Mathematics, University of Missouri, Columbia, Missouri 65211
iosevich@math.missouri.edu

Received: 9/30/08, Accepted: 11/19/08

Abstract

We prove that if $A \subset \mathbb{F}_q \backslash \{0\}$ with $|A| > Cq^{\frac{5}{6}}$, then $|R(A) \cdot R(A)| \geq C'q^3$, where

$$R(A) = \left\{ \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in SL_2(\mathbb{F}_q) : a_{11}, a_{12}, a_{21} \in A \right\}.$$

The proof relies on a result, previously established by D. Hart and the second author, which implies that if $|A|$ is much larger than $q^{\frac{3}{4}}$ then

$$|\{(a_{11}, a_{12}, a_{21}, a_{22}) \in A \times A \times A \times A : a_{11}a_{22} - a_{12}a_{21} = 1\}| = |A|^4 q^{-1}(1 + o(1)).$$

1. Introduction

Let $SL_2(\mathbb{F}_q)$ denote the set of two by two matrices with determinant one over the finite field with q elements.

Definition 1. *Given $A \subset \mathbb{F}_q$, let*

$$R(A) = \left\{ \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in SL_2(\mathbb{F}_q) : a_{11}, a_{12}, a_{21} \in A \right\}.$$

Observe that the size of $R(A)$ is exactly $|A|^3$. The purpose of this paper is to determine how large A needs to be to ensure that the product set

$$R(A) \cdot R(A) = \{M \cdot M' : M, M' \in R(A)\}$$

contains a positive proportion of all the elements of $SL_2(\mathbb{F}_q)$, q prime. This question is partly motivated by the following result due to Harald Helfgott ([5]). See his paper for further background on this problem and related references. See also [2] where Helfgott's result is proved for general fields.

Theorem 2. (Helfgott) *Let p be a prime. Let E be a subset of $SL_2(\mathbb{Z}/p\mathbb{Z})$ not contained in any proper subgroup.*

- *Assume that* $|E| < p^{3-\delta}$ *for some fixed* $\delta > 0$. *Then* $|E \cdot E \cdot E| > c|E|^{1+\epsilon}$, *where* $c > 0$ *and* $\epsilon > 0$ *depend only on* δ.
- *Assume that* $|E| > p^{\delta}$ *for some fixed* $\delta > 0$. *Then there is an integer* $k > 0$, *depending only on* δ, *such that every element of* $SL_2(\mathbb{Z}/p\mathbb{Z})$ *can be expressed as a product of at most* k *elements of* $E \cup E^{-1}$.

Our main result is the following.

Theorem 3. *Let* $A \subset \mathbb{F}_q \backslash \{0\}$ *with* $|A| \geq Cq^{\frac{5}{6}}$. *Then there exists* $C' > 0$ *such that*

$$|R(A) \cdot R(A)| \geq C'|SL_2(\mathbb{F}_q)| \geq C''q^3. \tag{1}$$

Remark 4. *Observe that if* $q = p^2$, *then* $\mathbb{F}_q$ *contains* $\mathbb{F}_p$ *as a sub-field. Since* $R(\mathbb{F}_p)$ *is a sub-group of* $SL_2(\mathbb{F}_q)$ *we see that the threshold assumption on the size of* A *in Theorem 3 cannot be improved beyond* $|A| \geq q^{\frac{1}{2}}$.

We shall make use of the following result due to D. Hart and A. Iosevich ([4]).

Theorem 5. *Let* $E \subset \mathbb{F}_q^d$, $d \geq 2$, *and define*

$$\nu(t) = |\{(x, y) \in E \times E : x \cdot y \equiv x_1y_1 + \cdots + x_dy_d = t\}|.$$

Then $\nu(t) = |E|^2 q^{-1} + \mathcal{D}(t)$, *where for every* $t > 0$, $|\mathcal{D}(t)| < |E|q^{\frac{d-1}{2}}$. *In particular, if* $|E| > q^{\frac{d+1}{2}}$, *then* $\nu(t) > 0$ *and as* E *grows beyond this threshold,*

$$\nu(t) = |E|^2 q^{-1}(1 + o(1)).$$

Remark 6. *The proof of Theorem 5 goes through unchanged if* $x \cdot y$ *is replaced by any non-degenerate bi-linear form* $B(x, y)$. *In particular, we can replace* $x_1y_1 + x_2y_2$ *by* $x_1y_1 - x_2y_2$ *in the case* $d = 2$ *and this is what we actually use in this paper. More precisely, we shall use the fact that if* $E = A \times A$ *and the size of* A *is much greater than* $q^{\frac{3}{4}}$, *then*

$$|\{(a, b, c, d) \in A \times A \times A \times A : ad - bc = 1\}| = |A|^4 q^{-1}(1 + o(1)). \tag{2}$$

1.1. Structure of the Proof of Estimate (1)

The basic idea behind the argument below is the following. Let $T \in SL_2(\mathbb{F}_q)$ and define

$$\nu(T) = |\{(S, S') \in R(A) \times R(A) : S \cdot S' = T\}.$$

We prove below that $\sqrt{\text{var}(\nu)} \leq C|A|^3 q^{-\frac{1}{2}}$, where variance is defined in the usual way as $\mathbb{E}\left((\nu - \mathbb{E}(\nu))^2\right)$, with the expectation defined, also in the usual way, as

$$\mathbb{E}(\nu) = |SL_2(\mathbb{F}_q)|^{-1} \sum_{T \in SL_2(\mathbb{F}_q)} \nu(T) = |A|^6 |SL_2(\mathbb{F}_q)|^{-1} = |A|^6 q^{-3}(1 + o(1)).$$

One can then check by a direct computation that $\sqrt{\mathrm{var}(\nu)}$ is much smaller than $\mathbb{E}(\nu)$ if $|A| \geq Cq^{\frac{5}{6}}$, with C sufficiently large, and we conclude that in this regime, $\nu(T)$ is concentrated around its expected value $\mathbb{E}(\nu) = |A|^6 q^{-3}(1+o(1))$.

1.2. Fourier Analysis Used in This Paper

We shall make use of the following basic formulas of Fourier analysis on $\mathbb{F}_q^d$. Let $f : \mathbb{F}_q^d \to \mathbb{C}$ and let χ denote a non-trivial additive character on $\mathbb{F}_q$. Define

$$\widehat{f}(m) = q^{-d} \sum_{x \in \mathbb{F}_q^d} \chi(-x \cdot m) f(x).$$

It is not difficult to check that

$$f(x) = \sum_{m \in \mathbb{F}_q^d} \chi(x \cdot m) \widehat{f}(m) \qquad \text{(Inversion)}$$

and

$$\sum_{m \in \mathbb{F}_q^d} \left|\widehat{f}(m)\right|^2 = q^{-d} \sum_{x \in \mathbb{F}_q^d} |f(x)|^2. \qquad \text{(Plancherel)}$$

2. Proof of Theorem 3 (Estimate 1)

We are looking to solve the equation

$$\begin{pmatrix} a_{11} & a_{12} \\ a_{21} & \frac{1+a_{12}a_{21}}{a_{11}} \end{pmatrix} \cdot \begin{pmatrix} b_{11} & b_{21} \\ b_{12} & \frac{1+b_{12}b_{21}}{b_{11}} \end{pmatrix} = \begin{pmatrix} t & \alpha \\ \beta & \frac{1+\alpha\beta}{t} \end{pmatrix},$$

which leads to the equations

$$a_{11}b_{11} + a_{12}b_{12} = t, \quad \frac{b_{21}}{b_{11}}t + \frac{a_{12}}{b_{11}} = \alpha, \quad \text{and} \quad \frac{a_{21}}{a_{11}}t + \frac{b_{12}}{a_{11}} = \beta. \tag{3}$$

Let D_t denote the characteristic function of the set

$$\{(a_{11}, b_{11}, a_{12}, b_{12}) \in A \times A \times A \times A : a_{11}b_{11} + a_{12}b_{12} = t\}$$

and let $E = A \times A$. Then the number of six-tuplets satisfying the equations (3) above equals

$$\begin{aligned} \nu(t,\alpha,\beta) &= \frac{1}{q^2} \sum_{u,v} \sum_{\substack{a_{11},b_{11},a_{12} \\ b_{12},a_{21},b_{21}}} \Big(D_t(a_{11},b_{11},a_{12},b_{12}) E(a_{21},b_{21}) \\ &\qquad \chi(u(b_{21}t + a_{12} - \alpha b_{11})) \chi(v(a_{21}t + b_{12} - \beta a_{11}) \Big) \\ &= q^{-2}|D_t||E| + q^4 \sum_{\mathbb{F}_q^2 \setminus \{(0,0)\}} \widehat{D}_t(\beta v, \alpha u, -u, -v) \widehat{E}(tv, tu) \\ &= \nu_0(t,\alpha,\beta) + \nu_{main}(t,\alpha,\beta). \end{aligned}$$

By (2), $\nu_0(t,\alpha,\beta) = q^{-3}|A|^6(1+o(1))$, which implies that

$$\sum_{t,\alpha,\beta} \nu_0^2(t,\alpha,\beta) = q^{-3}|A|^{12}(1+o(1)).$$

We now estimate $\sum_{t,\alpha,\beta} \nu_{main}^2(t,\alpha,\beta)$. By Cauchy–Schwarz and Plancherel,

$$\begin{aligned}\nu_{main}^2(t,\alpha,\beta) &\leq q^8 \textstyle\sum_{u,v} \left|\widehat{D}_t(\beta v, \alpha u, -u, -v)\right|^2 \cdot \sum_{u,v} \left|\widehat{E}(tv,tu)\right|^2 \\ &\leq |E|q^6 \textstyle\sum_{u,v} \left|\widehat{D}_t(\beta v, \alpha u, -u, -v)\right|^2.\end{aligned}$$

Now,

$$|E|q^6 \sum_{\alpha,\beta} \sum_{u,v} \left|\widehat{D}_t(\beta v, \alpha u, -u, -v)\right|^2 = |E|q^6 q^{-4}|A|^4 q^{-1}(1+o(1))$$

as long as $|E|$ is much larger than $q^{\frac{3}{2}}$. It follows that $\sum_{t\neq 0,\alpha,\beta} \nu_{main}^2(t,\alpha,\beta) \leq |A|^6 q^2$. Hence,

$$\sum_{t,\alpha,\beta} \nu^2(t,\alpha,\beta) \leq C(|A|^{12}q^{-3} + |A|^6 q^2). \tag{4}$$

In view of (4), we have

$$\begin{aligned}\left(|A|^6 - \sum_{\alpha,\beta} \nu(0,\alpha,\beta)\right)^2 &= \left(\sum_{t\neq 0,\alpha,\beta} \nu(t,\alpha,\beta)\right)^2 \\ &\leq C|\text{support}(\nu)| \cdot (|A|^{12}q^{-3} + |A|^6 q^2).\end{aligned}$$

If we can show that

$$\sum_{\alpha,\beta} \nu(0,\alpha,\beta) \leq \frac{1}{2}|A|^6, \tag{5}$$

then it would follow that

$$|\text{support}(\nu)| \gtrsim C \min\left\{q^3, \frac{|A|^6}{q^2}\right\}.$$

This expression is not less than $C|SL_2(\mathbb{F}_q)| = q^3(1+o(1))$ if $|A| \geq Cq^{\frac{5}{6}}$, as desired.

We are left to establish (5). Observe that if $t = 0$, then $\beta = -\alpha^{-1}$. Plugging this into (3) we see that this forces $a_{11} = -\alpha b_{12}$ and $a_{12} = \alpha b_{11}$, which implies that $\nu(0,\alpha,\beta) = \nu(0,\alpha,-\alpha^{-1}) \leq q^4$. This, in turn, implies that $\sum_{\alpha,\beta} \nu(0,\alpha,\beta) = \sum_{\alpha} \nu(0,\alpha,-\alpha^{-1}) \leq q^5$. Now, since $q^5 \leq \frac{1}{2}|A|^6$ if $|A| \geq Cq^{\frac{5}{6}}$, the proof is complete.

3. Proof of Theorem 5

To prove Theorem 5, we start out by observing that $\nu(t)=\sum_{x,y\in E} q^{-1}\sum_{s\in\mathbb{F}_q}\chi(s(x\cdot y-t))$, where χ is a non-trivial additive character on $\mathbb{F}_q$. It follows that $\nu(t)=|E|^2q^{-1}+\mathcal{D}$, where

$$\mathcal{D}=\sum_{x,y\in E} q^{-1}\sum_{s\neq 0}\chi(s(x\cdot y-t)).$$

Viewing $\mathcal{D}$ as a sum in x, applying the Cauchy-Schwarz inequality, and dominating the sum over $x\in E$ by the sum over $x\in\mathbb{F}_q^d$, we see that

$$\mathcal{D}^2\leq |E|\sum_{x\in\mathbb{F}_q^d} q^{-2}\sum_{s,s'\neq 0}\sum_{y,y'\in E}\chi(sx\cdot y-s'x\cdot y')\chi(t(s'-s)). \tag{6}$$

Orthogonality in the x variable yields that the right-hand side of (3.1) equals

$$|E|q^{d-2}\sum_{\substack{sy=s'y'\\ s,s'\neq 0}}\chi(t(s'-s))E(y)E(y').$$

If $s\neq s'$ we may set $a=s/s', b=s'$ and obtain

$$|E|q^{d-2}\sum_{\substack{y\neq y'\\ ay=y'\\ a\neq 1,b}}\chi(tb(1-a))E(y)E(y')=-|E|q^{d-2}\sum_{y\neq y',a\neq 1}E(y)E(ay),$$

and the absolute value of this quantity is at most

$$|E|q^{d-2}\sum_{y\in E}|E\cap l_y|\leq |E|^2q^{d-1},$$

since $|E\cap l_y|\leq q$ by the virtue of the fact that each line contains exactly q points.

If $s=s'$, then we get $|E|q^{d-2}\sum_{s,y}E(y)=|E|^2q^{d-1}$. It follows that $\nu(t)=|E|^2q^{-1}+\mathcal{D}(t)$, where $\mathcal{D}^2(t)\leq -Q(t)+|E|^2q^{d-1}$, with $Q(t)\geq 0$. This gives us $\mathcal{D}^2(t)\leq |E|^2q^{d-1}$, so that

$$|\mathcal{D}(t)|\leq |E|q^{\frac{d-1}{2}}. \tag{7}$$

We conclude that $\nu(t)=|E|^2q^{-1}+\mathcal{D}(t)$ with $|\mathcal{D}(t)|$ bounded as in (7). This quantity is strictly positive if $|E|>q^{\frac{d+1}{2}}$ with a sufficiently large constant $C>0$. This completes the proof of Theorem 5.

4. Remarks and Questions

- It has been recently pointed out to us by O. Dinai that the conclusion of our main result can be obtained using the methods in [1].
- The Fourier analysis used in the proof of both the first and second assertions of Theorem 3 is almost entirely formal as no hard estimates are used, even on the level of Gauss sums. This suggests that the result should be generalizable to a much wider setting.
- A natural analog of the second part of Theorem 3 is proved in [4] in all dimensions. Thus in principle there is a launching mechanism to attack the second part, though it is certainly more difficult technically.
- One of the consequences of the main result of this paper is to give a quantitative version of Helfgott's result (Theorem 2) for a class of relatively large subsets of $SL_2(\mathbb{F}_q)$. In analogy with the results in [3] it should be possible to address the question of obtaining explicit exponents for relatively small sets as well.

References

[1] L. Babai, N. Nikolov and L. Pyber *Product Growth and Mixing in Finite Groups*, Proceedings of the 19th annual ACM-SIAM symposium on discrete algorithms, 2008.

[2] O. Dinai, Ph.D. Dissertation (2008).

[3] M. Garaev, *An explicit sum-product estimate in* $\mathbb{F}_p$, Int. Math. Res. Notes **2007** (2007), article ID rnm035

[4] D. Hart and A. Iosevich, *Sums and products in finite fields: an integral geometric viewpoint*, Contemporary Math., to appear.

[5] H. A. Helfgott, *Growth and generation in* $SL_2(\mathbb{Z}/p\mathbb{Z})$, Ann. of Math. **167** (2008), 601-623.