Journal of Integer Sequences, Vol. 20 (2017),
Article 17.9.7

Divisors on Overlapped Intervals and Multiplicative Functions

José Manuel Rodríguez Caballero
Département de Mathématiques
UQAM
Case Postale 8888, Succ. Centre-ville
Montréal, Québec H3C 3P8
Canada
rodriguez_caballero.jose_manuel@uqam.ca

Abstract

Reutenauer and Kassel introduced a family $P_n(q)$ of polynomials defined in terms of divisors of n on overlapped intervals. The evaluation of $P_n(q)$ at roots of unity of order 2, 3, 4, 6 form well-known integer sequences related to the number of integer solutions of the equations $x^2 + y^2 = n$, $x^2 + 2y^2 = n$, and $x^2 + xy + y^2 = n$. Also, $P_n(1)$ is the sum of divisors of n. In this paper we define a new family $L_n(q)$ of polynomials defined in terms of divisors of n on overlapped intervals, slightly modifying the definition of $P_n(q)$. The values of $L_n(q)$ at $q = 1$ and $q = -1$ are related to the sum of divisors of n and to the number of integer solutions of the equations $x^2 + xy + y^2 = n$ and $x^2 + 3y^2 = n$.

1 Introduction

For a given integer $n \geq 1$, consider the two-sided sequence

$$p_{n,k} = \ln\left(k + \sqrt{k^2 + 2\,n}\right),$$

where $k \in \mathbb{Z}$ and define the intervals

$$\mathcal{P}_{n,k} = \left(p_{n,k} - \ln 2, p_{n,k}\right].$$

Kassel and Reutenauer [2] introduced the polynomials[1]

$$\frac{P_n(q)}{q^{n-1}} = \sum_{d|n} \sum_{k \in \mathbb{Z}} \mathbf{1}_{\mathcal{P}_{n,k}} (\ln d)\ q^k,$$

where $\mathbf{1}_A(x)$ is the characteristic function of the set A, i.e., $\mathbf{1}_A(x) = 1$ if $x \in A$, otherwise $\mathbf{1}_A(x) = 0$. Each polynomial $P_n(q)$ is monic of degree $2n-2$, its coefficients are non-negative integers and it is self-reciprocal [3]. The evaluations of $P_n(q)$ at some complex roots of 1 have number-theoretical interpretations [3], e.g.,

$$\begin{aligned}
\sigma(n) &= P_n(1), \\
\frac{r_{1,0,1}(n)}{4} &= P_n(-1), \\
\frac{r_{1,0,2}(n)}{2} &= \left|P_n\left(\sqrt{-1}\right)\right|, \\
\frac{r_{1,1,1}(n)}{6} &= \mathrm{Re}\ P_n\left(\frac{-1+\sqrt{-3}}{2}\right),
\end{aligned}$$

where $\sigma(n)$, $\frac{r_{1,0,1}(n)}{4}$, $\frac{r_{1,0,2}(n)}{2}$ and $\frac{r_{1,1,1}(n)}{6}$ are multiplicative functions [5] given by

$$\begin{aligned}
\sigma(n) &= \sum_{d|n} d, \\
r_{a,b,c}(n) &= \#\left\{(x,y) \in \mathbb{Z}^2 : \quad a\,x^2 + b\,x\,y + c\,y^2 = n\right\}.
\end{aligned}$$

Furthermore, for $q = \frac{1+\sqrt{-3}}{2}$, the same sequence $n \mapsto P_n(q)$ is related to $r_{1,0,1}(n)$ in three ways [4], depending on the congruence class of n in $\mathbb{Z}/3\mathbb{Z}$,

$$\left|P_n\left(\frac{1+\sqrt{-3}}{2}\right)\right| = \begin{cases} r_{1,0,1}(n), & \text{if } n \equiv 0 \pmod 3; \\ \frac{1}{4}\, r_{1,0,1}(n), & \text{if } n \equiv 1 \pmod 3; \\ \frac{1}{2}\, r_{1,0,1}(n), & \text{if } n \equiv 2 \pmod 3. \end{cases}$$

For any integer $n \geq 1$, consider the two-sided sequence

$$\ell_{n,k} = \ln\left(\tfrac{3}{2}\,k + \sqrt{\left(\tfrac{3}{2}\,k\right)^2 + 3\,n}\right)$$

and the intervals

$$\mathcal{L}_{n,k} = \left(\ell_{n,k} - \ln 3, \ell_{n,k}\right],$$

[1] The original definition of $P_n(q)$, which we refer to as *Kassel-Reutenauer polynomials* [2] is rather different, but equivalent, to the one presented here. We preferred to take the logarithm of the divisors in place of the divisors themselves in order to work with intervals $\mathcal{P}_{n,k}$ of constant length.



Figure 1: Representation of $L_6(q)$.

where k runs over the integers. Define a variation of the polynomials $P_n(q)$ as follows:

$$\frac{L_n(q)}{q^{n-1}} = \sum_{d|n} \sum_{k\in\mathbb{Z}} \mathbf{1}_{\mathcal{L}_{n,k}} (\ln d)\ q^k.$$

For example, in order to compute $L_6(q)$ from the definition, we need to consider the intervals $(\ell_{6,k} - \ln 3, \ell_{6,k}]$ on the real line and to count the number of values of $\ln d$ inside each interval, where d runs over the divisors of n. These data are shown in Figure 1, where the numbers $\ell_{6,k}$ are plotted on the line below (the corresponding values of k are labelled) whereas the numbers $\ln d$ are plotted on the line above (the corresponding values of d are labelled). Counting the number of intersections between the horizontal and the vertical lines, we obtain that the coefficients of $\frac{L_6(q)}{q^{6-1}}$ are as follows:

$$\frac{L_6(q)}{q^{6-1}} = q^5 + q^4 + q^3 + 2\,q^2 + 2\,q + 2\,q^0 + 2\,q^{-1} + 2\,q^{-2} + q^{-3} + q^{-4} + q^{-5}.$$

Like $P_n(q)$, the polynomial $L_n(q)$ is monic of degree $2\,n - 2$, self-reciprocal and its coefficients are non-negative integers. The aim of this paper is to express the multiplicative functions [5, p. 421] $\frac{r_{1,1,1}(n)}{6}$ and $\frac{r_{1,0,3}(n)}{2}$ in terms of the evaluations of $L_n(q)$ at roots of the unity. More precisely, we will prove the following result.

Theorem 1. *For each $n \geq 1$,*

$$\text{A002324}(n) \ := \ \frac{r_{1,1,1}(n)}{6} = 4\,\sigma(n) - 3\,L_n\,(1)\,, \tag{1}$$

$$\text{A096936}(n) \ := \ \frac{r_{1,0,3}(n)}{2} = L_n\,(-1)\,. \tag{2}$$

2 Auxiliary results for the first identity of Theorem 1

For any $n \geq 1$, we will use the notation

$$d_{a,m}(n) := \#\left\{d|n : \quad d \equiv a \pmod{m}\right\}.$$

We will use the following well-known result [1].

Lemma 2. *For all integers $n \geq 1$,*

$$\frac{r_{1,1,1}(n)}{6} = d_{1,3}(n) - d_{2,3}(n).$$

Lemma 3. *For any integer $n \geq 1$,*

$$3\left\lceil 3^{-1}\,n\right\rceil - n \;=\; \begin{cases} 0, & \text{if } n \equiv 0 \pmod 3; \\ 2, & \text{if } n \equiv 1 \pmod 3; \\ 1, & \text{if } n \equiv 2 \pmod 3. \end{cases}$$

$$n - 3\left\lfloor 3^{-1}\,n\right\rfloor \;=\; \begin{cases} 0, & \text{if } n \equiv 0 \pmod 3; \\ 1, & \text{if } n \equiv 1 \pmod 3; \\ 2, & \text{if } n \equiv 2 \pmod 3. \end{cases}$$

Proof. It is enough to evaluate $3\left\lceil 3^{-1}\,n\right\rceil - n$ and $3\left\lceil 3^{-1}\,n\right\rceil - n$ at $n = 3k + r$, for $k \in \mathbb{Z}$ and $r \in \{0, 1, 2\}$. ∎

Lemma 4. *For any pair of integers $n \geq 1$ and k, the inequalities*

$$\ell_{n,k} - \ln 3 < \ln d \leq \ell_{n,k}$$

hold if and only if the inequalities

$$3^{-1}\,d - \frac{n}{d} \leq k < d - 3^{-1}\,\frac{n}{d}$$

hold.

Proof. The inequalities

$$\ell_{n,k} - \ln 3 < \ln d \leq \ell_{n,k}$$

are equivalent to

$$\ln d \leq \ell_{n,k} < \ln d + \ln 3.$$

Applying the strictly increasing function $x \mapsto \frac{e^x}{3} - n\,e^{-x}$ to the last inequalities we obtain the following equivalent inequalities

$$3^{-1}\,d - \frac{n}{d} \leq k < d - 3^{-1}\,\frac{n}{d}.$$

Indeed, $\frac{e^{\ln d}}{3} - n\,e^{-\ln d} = 3^{-1}\,d - \frac{n}{d}$, $\frac{e^{\ln d + \ln 3}}{3} - n\,e^{-(\ln d + \ln 3)} = d - 3^{-1}\frac{n}{d}$ and

$$
\begin{aligned}
\frac{e^{\ell_{n,k}}}{3} - n\, e^{-\ell_{n,k}} &= \frac{\frac{3}{2}\,k + \sqrt{\left(\frac{3}{2}\,k\right)^2 + 3\,n}}{3} - \frac{n}{\frac{3}{2}\,k + \sqrt{\left(\frac{3}{2}\,k\right)^2 + 3\,n}} \\
&= \frac{\frac{3}{2}\,k + \sqrt{\left(\frac{3}{2}\,k\right)^2 + 3\,n}}{3} + \frac{\frac{3}{2}\,k - \sqrt{\left(\frac{3}{2}\,k\right)^2 + 3\,n}}{3} \\
&= k.
\end{aligned}
$$

So the lemma is proved. □

Lemma 5. *Let $n \geq 1$ be an integer. For all $d|n$,*

$$
\sum_{k\in\mathbb{Z}} \mathbf{1}_{\mathcal{L}_{n,k}}(\ln d) = \left\lceil d - 3^{-1}\,\frac{n}{d}\right\rceil - \left\lceil 3^{-1}\,d - \frac{n}{d}\right\rceil.
$$

Proof. For all integers $n \geq 1$ and k, we have that

$$
\begin{aligned}
& \sum_{k\in\mathbb{Z}} \mathbf{1}_{\mathcal{L}_{n,k}}(\ln d) \\
= \; & \#\left\{k \in \mathbb{Z} : \quad \ell_{n,k} - \ln 3 < \ln d \leq \ell_{n,k}\right\} \\
= \; & \#\left\{k \in \mathbb{Z} : \quad 3^{-1}\,d - \frac{n}{d} \leq k < d - 3^{-1}\,\frac{n}{d}\right\} \quad \text{(Lemma 4)} \\
= \; & \#\left\{k \in \mathbb{Z} : \quad \left\lceil 3^{-1}\,d - \frac{n}{d}\right\rceil \leq k < \left\lceil d - 3^{-1}\,\frac{n}{d}\right\rceil\right\} \\
= \; & \left\lceil d - 3^{-1}\,\frac{n}{d}\right\rceil - \left\lceil 3^{-1}\,d - \frac{n}{d}\right\rceil.
\end{aligned}
$$

So the lemma is proved. □

3 Auxiliary results for the second identity of Theorem 1

We will use the following well-known result [5, p. 421].

Lemma 6. *The function $\frac{r_{1,0,3}(n)}{2}$ is multiplicative.*

We will use the following well-known result [1].

Lemma 7. *For all integers $n \geq 1$,*

$$
\frac{r_{1,0,3}(n)}{2} = d_{1,3}(n) - d_{2,3}(n) + 2\left(d_{4,12}(n) - d_{8,12}(n)\right).
$$

Recall that the nonprincipal Dirichlet character mod 3 is the 3-periodic arithmetic function $\chi_3(n)$ given by $\chi_3(0) = 0$, $\chi_3(1) = 1$ and $\chi_3(2) = -1$.

Lemma 8. *For all $n \geq 1$,*

$$\frac{(-1)^{\lfloor 3^{-1}\, n \rfloor} - (-1)^{\lceil 3^{-1}\, n \rceil}}{2} = (-1)^{n-1}\, \chi_3(n).$$

Proof. It is enough to substitute $n = 3k + r$, with $k \in \mathbb{Z}$ and $r \in \{0, 1, 2\}$, in both sides in order to check that they are equal. $\square$

Lemma 9. *For all $n \geq 1$,*

$$\sum_{d|n} (-1)^{\frac{n}{d}-1}\, (-1)^{d-1}\, \chi_3(d) = (-1)^{n-1}\, \frac{r_{1,0,3}(n)}{2}.$$

Proof. By Lemma 6, the function $\frac{r_{1,0,3}(n)}{2}$ is multiplicative. Also, it is easy to check that the functions $(-1)^{n-1}$ and $\chi_3(n)$ are multiplicative. So the functions $f(n) = (-1)^{n-1}\, \frac{r_{1,0,3}(n)}{2}$ and $(-1)^{n-1}\, \chi_3(n)$ are multiplicative, because the multiplicative property is preserved by ordinary product. The function $g(n) = \sum_{d|n} (-1)^{\frac{n}{d}-1}\, (-1)^{d-1}\, \chi_3(d)$ is multiplicative, because Dirichlet convolution preserves the multiplicative property. So it is enough to prove that $f\left(p^k\right) = g\left(p^k\right)$ for each prime power p^k.

Considering the case $p = 2$. The following elementary equivalences hold for any integer $m \geq 0$,

$$\begin{aligned}
2^m \equiv 1 \pmod{3} &\iff m \equiv 0 \pmod{2}, \\
2^m \equiv 2 \pmod{3} &\iff m \equiv 1 \pmod{2}, \\
2^m \equiv 4 \pmod{12} &\iff m \equiv 0 \pmod{2} \text{ and } m \neq 0, \\
2^m \equiv 8 \pmod{12} &\iff m \equiv 1 \pmod{2} \text{ and } m \neq 1.
\end{aligned}$$

So, for each integer $k \geq 1$,

$$\begin{aligned}
d_{1,3}\left(2^k\right) &= \#[0, k] \cap 2\mathbb{Z} = \left\lfloor \frac{k}{2} \right\rfloor + 1, \\
d_{2,3}\left(2^k\right) &= \#[1, k] \cap (2\mathbb{Z} + 1) = \left\lceil \frac{k}{2} \right\rceil, \\
d_{4,12}\left(2^k\right) &= \#[2, k] \cap 2\mathbb{Z} = \left\lfloor \frac{k}{2} \right\rfloor, \\
d_{8,12}\left(2^k\right) &= \#[3, k] \cap (2\mathbb{Z} + 1) = \left\lceil \frac{k}{2} \right\rceil - 1.
\end{aligned}$$

For any $k \geq 1$, it follows that

$$
\begin{aligned}
g\left(2^k\right) &= \sum_{j=0}^{k} (-1)^{2^{k-j}-1} (-1)^{2^j-1} \chi_3\left(2^j\right) \\
&= \sum_{j=0}^{k} (-1)^{2^{k-j}-1} (-1)^{2^j-1} (-1)^j \\
&= -1-(-1)^k + \sum_{j=1}^{k-1} (-1)^j \\
&= -1-(-1)^k + \frac{-1-(-1)^k}{2} \\
&= -3\,\frac{1+(-1)^k}{2} \\
&= -3\left(1+\left\lfloor \frac{k}{2} \right\rfloor - \left\lceil \frac{k}{2} \right\rceil\right) \\
&= -\left(\left(\left\lfloor \frac{k}{2} \right\rfloor + 1\right) - \left\lceil \frac{k}{2} \right\rceil + 2\left(\left\lfloor \frac{k}{2} \right\rfloor - \left(\left\lceil \frac{k}{2} \right\rceil - 1\right)\right)\right) \\
&= (-1)^{2^k-1}\left(d_{1,3}\left(2^k\right) - d_{2,3}\left(2^k\right) + 2\left(d_{4,12}\left(2^k\right) - d_{8,12}\left(2^k\right)\right)\right) \\
&= f\left(2^k\right) \quad \text{(Lemma 7).}
\end{aligned}
$$

Let p and $k \geq 1$ be an odd prime and an integer respectively. Noticing that $(-1)^{p^j-1} = 1$ for all $0 \leq j \leq k$. Also, $d_{4,12}\left(p^k\right) = d_{8,12}\left(p^k\right) = 0$, because p^k has not even divisor. So, for any $k \geq 1$,

$$
\begin{aligned}
g\left(p^k\right) &= \sum_{j=0}^{k} (-1)^{p^{k-j}-1} (-1)^{p^j-1} \chi_3\left(p^j\right) \\
&= \sum_{j=0}^{k} \chi_3\left(p^j\right) \\
&= d_{1,3}\left(p^k\right) - d_{2,3}\left(p^k\right) \\
&= (-1)^{p^k-1}\left(d_{1,3}\left(p^k\right) - d_{2,3}\left(p^k\right) + 2\left(d_{4,12}\left(p^k\right) - d_{8,12}\left(p^k\right)\right)\right) \\
&= f\left(p^k\right) \quad \text{(Lemma 7).}
\end{aligned}
$$

Therefore, $f(n) = g(n)$ for all $n \geq 1$. $\square$

Lemma 10. *For each $d|n$,*

$$\sum_{k\in\mathbb{Z}} \mathbf{1}_{\mathcal{L}_{n,k}}(\ln d)\,(-1)^k = \frac{1}{2}\left((-1)^{\left\lceil 3^{-1}\,d-\frac{n}{d}\right\rceil} - (-1)^{\left\lceil d-3^{-1}\,\frac{n}{d}\right\rceil}\right).$$

Proof. For any integer $n \geq 1$ and any $d|n$,

$$\begin{aligned}\sum_{k\in\mathbb{Z}} \mathbf{1}_{\mathcal{L}_{n,k}}(\ln d)\,(-1)^k &= \sum_{3^{-1}\,d-\frac{n}{d}\leq k< d-3^{-1}\,\frac{n}{d}} (-1)^k \quad \text{(Lemma 4)}\\ &= \sum_{\left\lceil 3^{-1}\,d-\frac{n}{d}\right\rceil\leq k<\left\lceil d-3^{-1}\,\frac{n}{d}\right\rceil} (-1)^k.\end{aligned}$$

Substituting $a = \lceil 3^{-1}\,d - \frac{n}{d}\rceil$, $b = \lceil d - 3^{-1}\,\frac{n}{d}\rceil$ and $q = -1$ in the geometric sum

$$\sum_{a\leq k<b} q^k = \frac{q^a - q^b}{1-q}$$

we obtain

$$\begin{aligned}\sum_{k\in\mathbb{Z}} \mathbf{1}_{\mathcal{L}_{n,k}}(\ln d)\,(-1)^k &= \frac{(-1)^{\left\lceil 3^{-1}\,d-\frac{n}{d}\right\rceil} - (-1)^{\left\lceil d-3^{-1}\,\frac{n}{d}\right\rceil}}{1-(-1)}\\ &= \frac{1}{2}\left((-1)^{\left\lceil 3^{-1}\,d-\frac{n}{d}\right\rceil} - (-1)^{\left\lceil d-3^{-1}\,\frac{n}{d}\right\rceil}\right).\end{aligned}$$

So the lemma is proved. □

4 Proof of the main result

We proceed now with the proof of the main result of this paper.

Proof of Theorem 1. Identity (1) follows from the following transformations,

$$\begin{aligned}L_n(1) &= \sum_{d|n}\sum_{k\in\mathbb{Z}} \mathbf{1}_{\mathcal{L}_{n,k}}(\ln d)\\ &= \sum_{d|n}\left(\left\lceil d-3^{-1}\,\frac{n}{d}\right\rceil - \left\lceil 3^{-1}\,d-\frac{n}{d}\right\rceil\right) \quad \text{(Lemma 5)}\\ &= \sum_{d|n}\left(d+\frac{n}{d}\right) + \sum_{d|n}\left\lceil -3^{-1}\,\frac{n}{d}\right\rceil - \sum_{d|n}\left\lceil 3^{-1}d\right\rceil\\ &= \sum_{d|n}\left(d+\frac{n}{d}\right) - \sum_{d|n}\left\lfloor 3^{-1}\,\frac{n}{d}\right\rfloor - \sum_{d|n}\left\lceil 3^{-1}d\right\rceil\end{aligned}$$

$$
\begin{aligned}
&= \frac{2}{3}\sum_{d|n}\left(d+\frac{n}{d}\right)+\frac{1}{3}\sum_{d|n}\left(\frac{n}{d}-3\,\left\lfloor 3^{-1}\,\frac{n}{d}\right\rfloor\right)-\frac{1}{3}\sum_{d|n}\left(3\,\left\lceil 3^{-1}d\right\rceil-d\right)\\
&= \frac{4\sigma(n)}{3}+\frac{d_{1,3}(n)+2d_{2,3}(n)}{3}-\frac{2d_{1,3}(n)+d_{2,3}(n)}{3} \quad \text{(Lemma 3)}\\
&= \frac{4\sigma(n)}{3}-\frac{d_{1,3}(n)-d_{2,3}(n)}{3}\\
&= \frac{4}{3}\,\sigma(n)-\frac{1}{3}\,\frac{r_{1,1,1}(n)}{6} \quad \text{(Lemma 2).}
\end{aligned}
$$

Identity (2) follows from the following transformations,

$$
\begin{aligned}
\frac{L_n\,(-1)}{(-1)^{n-1}} &= \sum_{d|n}\sum_{k\in\mathbb{Z}}\mathbf{1}_{\mathcal{L}_{n,k}}\,(\ln d)\,\,(-1)^k\\
&= \sum_{d|n}\frac{1}{2}\,\left((-1)^{\left\lceil 3^{-1}\,d-\frac{n}{d}\right\rceil}-(-1)^{\left\lceil \frac{n}{d}-3^{-1}\,d\right\rceil}\right) \quad \text{(Lemma 10)}\\
&= \sum_{d|n}\frac{1}{2}\,\left((-1)^{\left\lceil 3^{-1}\,d\right\rceil-\frac{n}{d}}-(-1)^{\frac{n}{d}-\left\lfloor 3^{-1}\,d\right\rfloor}\right)\\
&= \sum_{d|n}(-1)^{\frac{n}{d}-1}\,\frac{(-1)^{\left\lfloor 3^{-1}\,d\right\rfloor}-(-1)^{\left\lceil 3^{-1}\,d\right\rceil}}{2}\\
&= \sum_{d|n}(-1)^{\frac{n}{d}-1}\,(-1)^{d-1}\,\chi_3(d) \quad \text{(Lemma 8)}\\
&= (-1)^{n-1}\,\frac{r_{1,0,3}(n)}{2} \quad \text{(Lemma 9).}
\end{aligned}
$$

So the theorem is proved. □

5 Final remarks

1. Let k be a field and $\mathcal{R}$ be a k-algebra. The *codimension* of an ideal I of $\mathcal{R}$ is the dimension of the quotient $\mathcal{R}/I$ as a vector space over k.

   We let $\mathbb{Z}\oplus\mathbb{Z}$ denote the free abelian group of rank 2. Let $k=\mathbb{F}_q$ be the finite field with q elements and $\mathcal{R}=\mathbb{F}_q\,[\mathbb{Z}\oplus\mathbb{Z}]$ be its group algebra. Kassel and Reutenauer [2] proved that, for any prime power q, the number of ideals of codimension $n\geq 1$ of $\mathbb{F}_q\,[\mathbb{Z}\oplus\mathbb{Z}]$ is $(q-1)^2\,\,P_n(q)$. So it is natural to look for connections between the values of $L_n(q)$, when q is a prime power, and the algebraic structures related to $\mathbb{F}_q$.

2. The polynomials $P_n(q)$ are generated by the product [3]

$$\prod_{m\geq 1} \frac{(1-t^m)^2}{(1-q\,t^m)\,(1-q^{-1}\,t^m)} = 1 + \left(q + q^{-1} - 2\right) \sum_{n=1}^{\infty} \frac{P_n(q)}{q^{n-1}}\, t^n.$$

It would be interesting to find a similar generating function for $L_n(q)$.

6 Acknowledgments

The author deeply thanks Prof. S. Brlek and Prof. C. Reutenauer for reading carefully the paper and for providing useful comments. The author gratefully acknowledges the referee.

References

[1] M. D. Hirschhorn, Three classical results on representations of a number, in D. Foata and G.-N. Han, eds., *The Andrews Festschrift*, Springer, 2001, pp. 159–165.

[2] C. Kassel and C. Reutenauer, Counting the ideals of given codimension of the algebra of Laurent polynomials in two variables, *Michigan Math. J.*, to appear.

[3] C. Kassel and C. Reutenauer, Complete determination of the zeta function of the Hilbert scheme of n points on a two-dimensional torus, *Ramanujan J.*, to appear.

[4] C. Kassel and C. Reutenauer, The Fourier expansion of $\eta(z)\,\eta(2z)\,\eta(3z)\,/\,\eta(6z)$, *Archiv Math.* **108** (2017), 453–463.

[5] L. V. Lorenz, *Œuvres scientifiques de L. Lorenz*, Vol. 2, Lehmann & Stage, 1899.

[6] N. J. A. Sloane et al., The On-Line Encyclopedia of Integer Sequences, available at `http://oeis.org`.

2010 *Mathematics Subject Classification*: Primary 11A25; Secondary 11A07, 11A05.
Keywords: multiplicative function, representation by quadratic form, polynomial, divisor.

(Concerned with sequences A002324 and A096936.)

Received September 16 2017; revised versions received October 13 2017; October 21 2017. Published in *Journal of Integer Sequences*, October 29 2017.

Return to Journal of Integer Sequences home page.