Journal of Integer Sequences, Vol. 22 (2019),
Article 19.1.4

The γ-Vectors of Pascal-like Triangles Defined by Riordan Arrays

Paul Barry
School of Science
Waterford Institute of Technology
Ireland
pbarry@wit.ie

Abstract

We define and characterize the γ-matrix associated with Pascal-like matrices that are defined by ordinary and exponential Riordan arrays. We also define and characterize the γ-matrix of the reversions of these triangles, in the case of ordinary Riordan arrays. We are led to the γ-matrices of a one-parameter family of generalized Narayana triangles. Thus these matrices generalize the matrix of γ-vectors of the associahedron. The principal tools used are the bivariate generating functions of the triangles and Jacobi continued fractions.

1 Introduction

A polynomial $P_n(x) = \sum_{k=0}^n a_{n,k} x^k$ of degree n is said to be *reciprocal* if

$$P_n(x) = x^n P_n(1/x).$$

Thus we have

$$[x^k]P_n(x) = a_{n,k} = [x^k]x^n P_n(1/x).$$

Now

$$\begin{aligned}
[x^k]x^n P_n(1/x) &= [x^{k-n}] \sum_{i=0} a_{n,i} \frac{1}{x^i} \\
&= [x^{k-n}] \sum_{i=0} a_{n,i} x^{-i} \\
&= a_{n,n-k}.
\end{aligned}$$

Thus $P_n(x) = \sum_{k=0}^n a_{n,k}x^k$ defines a family of reciprocal polynomials if and only if $a_{n,k} = a_{n,n-k}$. We shall call a lower-triangular matrix $(a_{n,k})$ *Pascal-like* if

1. $a_{n,k} = a_{n,n-k}$
2. $a_{n,0} = a_{n,n} = 1$.

Such a matrix will then be the coefficient array of a family of monic reciprocal polynomials.

We have the following well-known result [7]

Proposition 1. *Let $P_n(x)$ be a reciprocal polynomial of degree n. Then there exists a unique polynomial γ_n of degree $\lfloor \frac{n}{2} \rfloor$ with the property*

$$P_n(x) = (1+x)^n \gamma_n \left(\frac{x}{(1+x)^2} \right).$$

If $P_n(x)$ has integer coefficients then so does $\gamma_n(x)$.

By this means, we can associate with every Pascal-like matrix $(a_{n,k})$ a matrix $(\gamma_{n,k})$ so that for all n, we have

$$P_n(x) = \sum_{k=0}^n a_{n,k}x^k = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_{n,k} x^k (1+x)^{n-2k}.$$

We shall call this matrix the γ-matrix associated with the coefficient array $(a_{n,k})$ of the family of polynomials $P_n(x)$.

We can characterize the matrix $(a_{n,k})$ in terms of the γ-matrix $(\gamma_{n,k})$ as follows. Before we do this, we shall change our notation somewhat. In algebraic topology, it is customary to use the notation $h(x)$ for palindromic (reciprocal) polynomials [9, 15]. Thus we shall set $h_n(x) = \sum_{k=0}^n h_{n,k}x^k$, where $(h_{n,k})$ now denotes a Pascal-like matrix. We shall denote by $h(x,y)$ the bivariate generating function of this matrix.

Proposition 2. *For a Pascal-like matrix $(h_{n,k})$ we have*

$$h_{n,k} = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-2i}{k-i} \gamma_{n,i}.$$

Proof. We have

$$\begin{aligned}
h_{n,k} &= [x^k]\sum_{i=0}^{n} h_{n,i}x^i \\
&= [x^k]\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_{n,i}x^i(1+x)^{n-2i} \\
&= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_{n,i}[x^k]x^i(1+x)^{n-2i} \\
&= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_{n,i}[x^{k-i}]\sum_{j=0}^{n-2i}\binom{n-2i}{j}x^j \\
&= \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_{n,i}\binom{n-2i}{k-i}.
\end{aligned}$$

□

Example 3. The identity

$$\binom{n}{k} = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \binom{n-2i}{k-i}\delta_{i,0}$$

shows that the matrix that begins

$$\begin{pmatrix}
1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0 \\
1 & 0 & 0 & 0 & 0 & 0 & 0
\end{pmatrix}$$

is the γ-matrix for the binomial matrix $\mathbf{B} = (\binom{n}{k})$ A007318. Here, we have used the *Annnnnn* number of the On-Line Encyclopedia of Integer Sequences [13, 14] for the binomial matrix (Pascal's triangle).

When $(\gamma_{n,k})$ is the γ-matrix for $(h_{n,k})$, we shall say the $(\gamma_{n,k})$ *generates*, or *is the generator of*, the matrix $(h_{n,k})$.

Example 4. The matrix that begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

with $\gamma_{n,0} = 1$, $\gamma_{n,\lfloor \frac{n}{2} \rfloor} = 1$, and 0 otherwise, generates the matrix $(h_{n,k})$ that begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 \\ 1 & 3 & 3 & 1 & 0 & 0 \\ 1 & 4 & 7 & 4 & 1 & 0 \\ 1 & 5 & 10 & 10 & 5 & 1 \end{pmatrix}.$$

2 Pascal-like matrices defined by Riordan arrays

We now wish to characterize the γ-matrices that are generators for the family of Pascal-like matrices that are determined by the one-parameter family of Riordan arrays

$$\left(\frac{1}{1-x}, \frac{x(1+rx)}{1-x}\right).$$

We shall also determine the (generalized) γ-matrices associated with the reversion of these triangles. We recall that an ordinary Riordan array $(g(x), f(x))$ is defined [1, 10, 11] by two power series

$$g(x) = 1 + g_1x + g_2x^2 + \cdots,$$
$$f(x) = x + f_2x^2 + f_3x^3 + \cdots,,$$

where the (n, k)-th element of the resulting lower-triangular matrix is given by

$$a_{n,k} = [x^n]g(x)f(x)^k.$$

Such matrices are invertible. When they have integer entries, the inverse again is an integer matrix (note that we have $a_{n,n} = 1$ in our case because $g_0 = 1$ and $f_1 = 1$). The bivariate generating function of the Riordan array (g, f) is given by

$$\frac{g(x)}{1 - yf(x)}.$$

Matrices defined in a similar manner but with $f(x)$ replaced by $\phi(x) = x^2 + \phi_3x^3 + \ldots$ are called "stretched" Riordan arrays [5]. They are not invertible but they do possess left inverses.

Example 5. The stretched Riordan array $\left(\frac{1}{1-x}, x^2\right)$ begins

$$\left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 \end{array}\right).$$

It is the γ-matrix for the Pascal-like triangle that begins

$$\left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 4 & 4 & 1 & 0 & 0 & 0 \\ 1 & 5 & 9 & 5 & 1 & 0 & 0 \\ 1 & 6 & 14 & 14 & 6 & 1 & 0 \\ 1 & 7 & 20 & 29 & 20 & 7 & 1 \end{array}\right).$$

Example 6. The matrix $\binom{n-k}{k}$ is the stretched Riordan array $\left(\frac{1}{1-x}, \frac{x^2}{1-x}\right)$ that begins

$$\left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 4 & 3 & 0 & 0 & 0 & 0 \\ 1 & 5 & 6 & 1 & 0 & 0 & 0 \end{array}\right).$$

It generates the Pascal-like matrix that begins

$$\left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 5 & 5 & 1 & 0 & 0 & 0 \\ 1 & 7 & 13 & 7 & 1 & 0 & 0 \\ 1 & 9 & 25 & 25 & 9 & 1 & 0 \\ 1 & 11 & 41 & 63 & 41 & 11 & 1 \end{array}\right).$$

We shall see that this is the Riordan array $\left(\frac{1}{1-x}, \frac{x(1+x)}{1-x}\right)$, which is A008288, the triangle of Delannoy numbers.

The bivariate generating function of the stretched Riordan array $(g(x), \phi(x))$ is given by

$$\frac{g(x)}{1-y\phi(x)}.$$

We have the following proposition [4].

Proposition 7. *The Riordan array* $\left(\frac{1}{1-x}, \frac{x(1+rx)}{1-x}\right)$ *is Pascal-like (for any* $r \in \mathbb{Z}$*).*

This is clear since in this case we have

$$h_{n,k} = \sum_{j=0}^{k} \binom{k}{j}\binom{n-j}{n-k-j} r^j = \sum_{j=0}^{k} \binom{k}{j}\binom{n-k}{n-k-j}(r+1)^j.$$

We can now characterize the γ-matrices that generate these Pascal-like matrices.

Proposition 8. *The* γ*-matrices that generate the Pascal-like matrices* $\left(\frac{1}{1-x}, \frac{x(1+rx)}{1-x}\right)$ *defined by ordinary Riordan arrays are given by the stretched Riordan arrays*

$$\left(\frac{1}{1-x}, \frac{rx^2}{1-x}\right),$$

with (n,k)*-th term*

$$\gamma_{n,k} = \binom{n-k}{k} r^k.$$

Proof. The generating function of the Pascal-like matrix $\left(\frac{1}{1-x}, \frac{x(1+rx)}{1-x}\right)$ is given by

$$h(x,y) = \frac{1}{1-x}\frac{1}{1-y\frac{x(1+rx)}{1-x}} = \frac{1}{1-(1+y)x-rx^2y}.$$

Similarly, the generating function of the matrix $(\binom{n-k}{k} r^k)$ is given by

$$\gamma(x,y) = \frac{1}{1-x}\frac{1}{1-y\frac{rx^2}{1-x}} = \frac{1}{1-x-rx^2y}.$$

We now have

$$h(x,y) = \gamma\left((1+y)x, \frac{y}{(1+y)^2}\right).$$

∎

We recall that for a generating function $f(x)$, its INVERT(α) transform is the generating function

$$\frac{f(x)}{1+\alpha x f(x)}.$$

Note that

$$\frac{\frac{v}{1+\alpha x v}}{1-\alpha x\frac{v}{1+\alpha x v}}=v,$$

and thus the inverse of the INVERT(α) transform is the INVERT($-\alpha$) transform.

Corollary 9. *The generating function $h(x,y)$ of the Pascal-like matrix $\left(\frac{1}{1-x},\frac{x(1+rx)}{1-x}\right)$ is the INVERT(y) transform of the generating function $\gamma(x,y)$ of the corresponding γ-matrix.*

Proof. A direct calculation shows that for $\gamma(x,y)=\frac{1}{1-x-rx^2y}$ we have

$$\frac{\gamma(x,y)}{1-yx\gamma(x,y)}=\frac{1}{1-(y+1)x-rx^2y}=h(x,y).$$

$\square$

Equivalently, we can say that the generating function of the γ-matrix is the INVERT($-y$) transform of the generating function of the corresponding Pascal-like matrix.

We make the following observation, which will be relevant when we discuss a family of generalized Narayana triangles. The γ-matrix corresponding to the signed Pascal-like matrix

$$\left(\frac{1}{1+x},\frac{-x(1+rx)}{1+x}\right)$$

has generating function

$$\frac{1}{1+x+rx^2y}.$$

This is the matrix with general term $(-1)^{n-k}r^k\binom{n-k}{k}$. By a signed Pascal-like matrix in this case we mean that $a_{n,k}=a_{n,n-k}$ but we now have $a_{n,0}=a_{n,n}=(-1)^n$.

We close this section by recalling the formula

$$\gamma_n=(1+x)^n\gamma_n\left(\frac{x}{(1+x)^2}\right).$$

We now note that the inverse of the Riordan array

$$\left(1,\frac{x}{(1+x)^2}\right)$$

is given by

$$\left(1,xc(x)^2\right),$$

where

$$c(x) = \frac{1-\sqrt{1-4x}}{2x}$$

is the generating function of the Catalan numbers $C_n = \frac{1}{n+1}\binom{2n}{n}$ A000108. In fact, we have the following result [9].

Proposition 10. ***(Zeilberger's Lemma).***

$$\gamma_{n,k} = [x^k]\frac{h_n(xc(x)^2)}{c(x)^n}.$$

We can use this result to find an explicit formula for $\gamma_{n,k}$ in terms of $h_{n,k}$. We let $\alpha_{n,k}$ be the general (n,k)-th element of the Riordan array $(1, xc(x)^2)$ [8]. We have

$$\alpha_{n,k} = \begin{cases} 1, & \text{if } n=0 \text{ and } k=0; \\ \binom{2n-1}{n-k}\frac{2k}{n+k}, & \text{otherwise;} \end{cases}$$

or, equivalently,

$$\alpha_{n,k} = \binom{2n-1}{n-k}\frac{2k+0^{n+k}}{n+k+0^{n+k}} = \binom{2n-2}{n-k} - \binom{2n-2}{n-k-2}.$$

We let $\beta_{n,k}$ be the general (n,k)-th term of the Riordan array $\left(1, \frac{x}{c(x)}\right)$. We have $\beta_{n,n}=1$, and

$$\beta_{n,k} = \sum_{j=0}^{n-k} \frac{(-1)^j}{n-k}\binom{k+j-1}{j}\binom{2(n-k)}{n-k-j},$$

otherwise. This is essentially A271875. Then we have the following result.

Corollary 11. *We have*

$$\gamma_{n,k} = \sum_{i=0}^{k}\left(\sum_{j=0}^{n} h_{n,j}\alpha_{i,j}\right)\beta_{n+k-i,n}.$$

Proof. We have

$$\begin{aligned} [x^k][x^k]\frac{h_n(xc(x)^2)}{c(x)^n} &= \sum_{i=0}^{n}[x^i]\sum_{j=0}^{n} h_{n,j}(xc(x^2))^j[x^{k-i}]\frac{1}{c(x)^n} \\ &= \sum_{i=0}^{k}\left(\sum_{j=0}^{n} h_{n,j}[x^i](xc(x)^2)^j\right)[x^{k-1+n}]\frac{x^n}{c(x)^n} \\ &= \sum_{i=0}^{k}\left(\sum_{j=0}^{n} h_{n,j}\alpha_{i,j}\right)\beta_{n+k-i,n}. \end{aligned}$$

$\square$

This gives us the following formula:

$$\gamma_{n,k}=\sum_{i=0}^{k}\sum_{j=0}^{n}h_{n,j}\binom{2i-1}{i-j}\frac{2j+0^{i+j}}{i+j+0^{i+j}}\cdot\begin{cases}1, & \text{if } i=k;\\ \sum_{m=0}^{k-i}\frac{m(-1)^m}{k-i}\binom{n-1+m}{m}\binom{2(k-i)}{k-i-m}, & \text{otherwise;}\end{cases}$$

which we can also write as

$$\gamma_{n,k}=\sum_{i=0}^{k}\sum_{j=0}^{n}h_{n,j}\binom{2i-1}{i-j}\frac{2j+0^{i+j}}{i+j+0^{i+j}}\,\mathrm{If}\left[i=k,1,\sum_{m=0}^{k-i}\frac{m(-1)^m}{k-i}\binom{n-1+m}{m}\binom{2(k-i)}{k-i-m}\right].$$

Example 12. If we take $(h_{n,k})$ to be the triangle of Eulerian numbers A008292 that begins

$$\begin{pmatrix}1&0&0&0&0&0&0\\1&1&0&0&0&0&0\\1&4&1&0&0&0&0\\1&11&11&1&0&0&0\\1&26&66&26&1&0&0\\1&57&302&302&57&1&0\\1&120&1191&2416&1191&120&1\end{pmatrix}$$

we find that the γ-matrix $(\gamma_{n,k})$ is the triangle A101280 that begins

$$\begin{pmatrix}1&0&0&0&0&0&0\\1&0&0&0&0&0&0\\1&2&0&0&0&0&0\\1&8&0&0&0&0&0\\1&22&16&0&0&0&0\\1&52&136&0&0&0&0\\1&114&720&272&0&0&0\end{pmatrix}.$$

This is the triangle of γ-vectors for the permutahedra (of type A). It also gives the number of permutations of n objects with k descents such that every descent is a peak [12].

Example 13. We consider the Pascal-like matrix $(h_{n,k})=\left(\frac{1}{1-x},x\right)$ that begins

$$\begin{pmatrix}1&0&0&0&0&0&0\\1&1&0&0&0&0&0\\1&1&1&0&0&0&0\\1&1&1&1&0&0&0\\1&1&1&1&1&0&0\\1&1&1&1&1&1&0\\1&1&1&1&1&1&1\end{pmatrix}.$$

We note that the row elements are constant. We have that

$$\gamma_{n,k} = \sum_{i=0}^{k}\sum_{j=0}^{n}\binom{2i-1}{i-j}\frac{2j+0^{i+j}}{i+j+0^{i+j}}\,\text{If}\left[i=k,1,\sum_{m=0}^{k-i}\frac{m(-1)^m}{k-i}\binom{n-1+m}{m}\binom{2(k-i)}{k-i-m}\right].$$

We find that the γ-matrix in this case begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & -2 & 0 & 0 & 0 & 0 & 0 \\ 1 & -3 & 1 & 0 & 0 & 0 & 0 \\ 1 & -4 & 3 & 0 & 0 & 0 & 0 \\ 1 & -5 & 6 & -1 & 0 & 0 & 0 \end{pmatrix}.$$

This is the matrix $\left(\binom{n-k}{k}(-1)^k\right)$. Thus

$$\sum_{i=0}^{\lfloor \frac{n}{2} \rfloor}\binom{n-2i}{k-i}\binom{n-i}{i}(-1)^i = \text{If}[k \le n, 1, 0].$$

3 Stretched Riordan arrays as γ-matrices

Every stretched Riordan array of the form

$$\left(\frac{1}{1-x}, x^2 g(x)\right),$$

where

$$g(x) = 1 + g_1 x + g_2 x^2 + \cdots$$

can be used to generate a Pascal-like matrix. Thus with each power series $g(x)$ above we can associate a Pascal-like matrix whose γ-matrix is given by this stretched Riordan array.

In this section, we shall concentrate on the case when $g(x) = \frac{1+rx}{1-x}$.

Example 14. For $r = 1$, we obtain the γ-matrix that begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 0 & 0 & 0 & 0 & 0 \\ 1 & 5 & 1 & 0 & 0 & 0 & 0 \\ 1 & 7 & 5 & 0 & 0 & 0 & 0 \\ 1 & 9 & 13 & 1 & 0 & 0 & 0 \end{pmatrix}.$$

The corresponding Pascal-like matrix then begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 6 & 6 & 1 & 0 & 0 & 0 \\ 1 & 9 & 17 & 9 & 1 & 0 & 0 \\ 1 & 12 & 36 & 36 & 12 & 1 & 0 \\ 1 & 15 & 64 & 101 & 64 & 15 & 1 \end{pmatrix}.$$

The row sums of this matrix, which begin

$$1, 2, 5, 14, 37, 98, 261, \ldots$$

give A077938, with generating function

$$\frac{1}{1-2x-x^2-2x^3}.$$

The diagonal sums, which begin

$$1, 1, 2, 4, 8, 16, 31, \ldots$$

are the Pentanacci numbers A001591 with generating function

$$\frac{1}{1-x-x^2-x^3-x^4-x^5}.$$

We have the following proposition.

Proposition 15. *The Pascal-like triangle that begins*

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & r+5 & r+5 & 1 & 0 & 0 & 0 \\ 1 & 2r+7 & 4r+13 & 2r+7 & 1 & 0 & 0 \\ 1 & 3r+9 & 11r+25 & 11r+25 & 3r+9 & 1 & 0 \\ 1 & 4r+11 & r^2+22r+41 & 2r^2+36r+63 & r^2+22r+41 & 4r+11 & 1 \end{pmatrix}$$

with γ-matrix given by the stretched Riordan array $\left(\frac{1}{1-x}, \frac{x^2(1+rx)}{1-x}\right)$, has row sums with generating function

$$\frac{1}{1-2x-x^2-2rx^3},$$

and diagonal sums given by the generalized Pentanacci numbers with generating function

$$\frac{1}{1-x-x^2-x^3-rx^4-rx^5}.$$

4 Reverting triangles

Let $h(x,y)$ be the generating function of the lower-triangular matrix $h_{n,k}$, with $h_{0,0}=1$. By the reversion of this triangle, we shall mean the triangle whose generating function $h^*(x,y)$ is given by

$$h^*(x,y)=\frac{1}{x}\text{Rev}_x(xh(x,y)).$$

Procedurally, this means that we solve the equation

$$uh(u,y)=x$$

and then we divide the solution $u(x,y)$ that satisfies $u(0,y)=0$ by x.

Proposition 16. *The generating function of the reversion of the Pascal-like matrix defined by the Riordan array* $\left(\frac{1}{1-x},\frac{x(1+rx)}{1-x}\right)$ *is given by*

$$h^*(x,y)=\frac{1}{1+x(y+1)}c\left(\frac{-rx^2y}{(1+x(y+1))^2}\right),$$

where

$$c(x)=\frac{1-\sqrt{1-4x}}{2x}$$

is the generating function of the Catalan numbers $C_n=\frac{1}{n+1}\binom{2n}{n}$*. (A000108).*

Proof. Solving the equation

$$\frac{u}{1-u(y+1)-ru^2y}=x$$

gives us

$$h^*(x,y)=\frac{-1-x(y+1)+\sqrt{1+2x(y+1)+x^2(1+2y(2r+1)+y^2)}}{2rx^2y}.$$

Thus

$$h^*(x,y)=\frac{1}{1+x(y+1)}c\left(\frac{-rx^2y}{(1+x(y+1))^2}\right).$$

□

We note that we can now calculate an expression for the terms of the reverted triangle, since, using the language of Riordan arrays, we have

$$h^*(x,y)=\left(\frac{1}{1+y(x+1)},\frac{-rx^2y}{(1+x(y+1))^2}\right)\cdot c(x).$$

Proposition 17. *We have*

$$[x^n][y^i]h^*(x,y) = h^*_{n,i} = \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^n(-r)^k \binom{n}{2k} C_k \binom{n-2k}{i-k}.$$

The γ-matrix of the reverted triangle $(h^*_{n,k})$ is given by

$$\gamma^*_{n,k} = (-1)^n(-r)^k \binom{n}{2k} C_k.$$

The γ-matrix $(\gamma^*_{n,k})$ of the reverted triangle $(h^*_{n,k})$ is the reversion of the triangle $\gamma_{n,k}$.

Proof. The expression for $h^*_{n,k}$ results from a direct calculation. Reverting the expression $\gamma(x,y) = \frac{1}{1-x-rx^2y}$ in the sense above gives us

$$\gamma^*(x,y) = \frac{1}{1+x} c\left(\frac{-rx^2y}{(1+x)^2}\right),$$

from which we deduce the other statements. $\square$

Example 18. For $r = -1, 0, 1$, the triangles $(h_{n,k})$ begin, respectively,

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ 1 & 3 & 3 & 1 & 0 & 0 & 0 \\ 1 & 4 & 6 & 4 & 1 & 0 & 0 \\ 1 & 5 & 10 & 10 & 5 & 1 & 0 \\ 1 & 6 & 15 & 20 & 15 & 6 & 1 \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 5 & 5 & 1 & 0 & 0 & 0 \\ 1 & 7 & 13 & 7 & 1 & 0 & 0 \\ 1 & 9 & 25 & 25 & 9 & 1 & 0 \\ 1 & 11 & 41 & 63 & 41 & 11 & 1 \end{pmatrix}.$$

The corresponding reverted triangles $(h^*_{n,k})$, are, respectively,

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ -1 & -6 & -6 & -1 & 0 & 0 & 0 \\ 1 & 10 & 20 & 10 & 1 & 0 & 0 \\ -1 & -15 & -50 & -50 & -15 & -1 & 0 \\ 1 & 21 & 105 & 175 & 105 & 21 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ -1 & -3 & -3 & -1 & 0 & 0 & 0 \\ 1 & 4 & 6 & 4 & 1 & 0 & 0 \\ -1 & -5 & -10 & -10 & -5 & -1 & 0 \\ 1 & 6 & 15 & 20 & 15 & 6 & 1 \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & -1 & 0 & 0 & 0 \\ 1 & -2 & -4 & -2 & 1 & 0 & 0 \\ -1 & 5 & 10 & 10 & 5 & -1 & 0 \\ 1 & -9 & -15 & -15 & -15 & -9 & 1 \end{pmatrix}.$$

Note that for $r=-1$, the reverted triangle is $(-1)^n$ times the Narayana triangle A001263.

The corresponding γ-matrices $(\gamma_{n,k})$ are given by, respectively,

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & -2 & 0 & 0 & 0 & 0 & 0 \\ 1 & -3 & 1 & 0 & 0 & 0 & 0 \\ 1 & -4 & 3 & 0 & 0 & 0 & 0 \\ 1 & -5 & 6 & -1 & 0 & 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 4 & 3 & 0 & 0 & 0 & 0 \\ 1 & 5 & 6 & 1 & 0 & 0 & 0 \end{pmatrix}.$$

The corresponding reverted γ-matrices $(\gamma^*_{n,k})$ are then, respectively,

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ -1 & -3 & 0 & 0 & 0 & 0 & 0 \\ 1 & 6 & 2 & 0 & 0 & 0 & 0 \\ -1 & -10 & -10 & 0 & 0 & 0 & 0 \\ 1 & 15 & 30 & 5 & 0 & 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 \\ -1 & 3 & 0 & 0 & 0 & 0 & 0 \\ 1 & -6 & 2 & 0 & 0 & 0 & 0 \\ -1 & 10 & -10 & 0 & 0 & 0 & 0 \\ 1 & -15 & 30 & -5 & 0 & 0 & 0 \end{pmatrix}.$$

It is interesting to represent the generating functions of the $(\gamma^*_{n,k})$ and the $(h^*_{n,k})$ triangles as Jacobi continued fractions. We have

Proposition 19. *The generating function $h^*(x,y)$ can be expressed as the Jacobi continued fraction*

$$\mathcal{J}(-(y+1), -(y+1), -(y+1), \ldots; -ry, -ry, -ry, \ldots).$$

The generating function $\gamma^*(x,y)$ can be expressed as the Jacobi continued fraction

$$\mathcal{J}(-1, -1, -1, \ldots; -ry, -ry, -ry, \ldots).$$

Proof. We solve the continued fraction equation

$$u = \frac{1}{1+(y+1)x+rx^2u}$$

to retrieve the generating function $h^*(x,y)$. Similarly, we solve the continued fraction equation

$$u = \frac{1}{1+x+rx^2u}$$

to retrieve the generating function $\gamma^*(x,y)$. □

Note that we have used the notation $\mathcal{J}(a,b,c,\ldots;r,s,t,\ldots)$ to denote the Jacobi continued fraction [2, 16]

$$\cfrac{1}{1-ax-\cfrac{rx^2}{1-bx-\cfrac{sx^2}{1-cx-\cfrac{tx^2}{1-\cdots}}}}.$$

We can now express the relationship between the generating functions $h^*(x,y)$ and $\gamma^*(x,y)$ in terms of repeated binomial transforms.

Corollary 20. *The generating function $h^*(x,y)$ is the $(-y)$-th binomial transform of the γ generating function $\gamma^*(x,y)$:*

$$h^*(x,y) = \frac{1}{1+xy}\gamma^*\left(\frac{x}{1+xy}, y\right).$$

Equivalently, the γ generating function $\gamma^*(x,y)$ is the y-th binomial transform of the generating function $h^*(x,y)$:

$$\gamma^*(x,y) = \frac{1}{1-xy}h^*\left(\frac{x}{1-xy}, y\right).$$

This reflects the general assertion that the reversion of an INVERT transform is a binomial transform.

5 The γ-vectors of generalized Narayana numbers

The Riordan array $\left(\frac{1}{1+x}, \frac{-x(1+rx)}{1+x}\right)$, with bivariate generating function

$$\frac{1}{1+x(y+1)+rx^2y},$$

has a γ-matrix with generating function

$$\frac{1}{1+x+rx^2y}.$$

We shall call elements of the reversions of the Riordan array $\left(\frac{1}{1+x}, \frac{-x(1+rx)}{1+x}\right)$ r-Narayana numbers. The Narayana numbers $N_{n,k} = \frac{1}{k+1}\binom{n+1}{k}\binom{n}{k}$ are then the 1-Narayana numbers. The bivariate generating function for the r-Narayana numbers is given by

$$\frac{1}{1-x(y+1)} c\left(\frac{rx^2y}{(1-x(y+1))^2}\right).$$

The bivariate generating function for the γ-matrix of the r-Narayana numbers is then obtained by reverting the generating function $\frac{1}{1+x+rx^2y}$. We thus obtain the following result.

Proposition 21. *The γ-matrix for the r-Narayana numbers has generating function*

$$\frac{1}{1-x} c\left(\frac{rx^2y}{(1-x)^2}\right).$$

This is the matrix that begins

$$\left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & r & 0 & 0 & 0 & 0 & 0 \\ 1 & 3r & 0 & 0 & 0 & 0 & 0 \\ 1 & 6r & 2r^2 & 0 & 0 & 0 & 0 \\ 1 & 10r & 10r^2 & 0 & 0 & 0 & 0 \\ 1 & 15r & 30r^2 & 5r^3 & 0 & 0 & 0 \end{array}\right),$$

with general term

$$\binom{n}{2k} r^k C_k.$$

For $r = -1, 0, 1$, the matrices $\left(\frac{1}{1+x}, \frac{-x(1+rx)}{1+x}\right)$ begin, respectively,

$$\left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ -1 & -5 & -5 & -1 & 0 & 0 & 0 \\ 1 & 7 & 13 & 7 & 1 & 0 & 0 \\ -1 & -9 & -25 & -25 & -9 & -1 & 0 \\ 1 & 11 & 41 & 63 & 41 & 11 & 1 \end{array}\right), \left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ -1 & -3 & -3 & -1 & 0 & 0 & 0 \\ 1 & 4 & 6 & 4 & 1 & 0 & 0 \\ -1 & -5 & -10 & -10 & -5 & -1 & 0 \\ 1 & 6 & 15 & 20 & 15 & 6 & 1 \end{array}\right),$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ -1 & -1 & -1 & -1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ -1 & -1 & -1 & -1 & -1 & -1 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{pmatrix}.$$

The corresponding matrices of r-Narayana numbers are, respectively,

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 1 & -2 & -4 & -2 & 1 & 0 & 0 \\ 1 & -5 & -10 & -10 & -5 & 1 & 0 \\ 1 & -9 & -15 & -15 & -15 & -9 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 1 & 0 & 0 & 0 & 0 \\ 1 & 3 & 3 & 1 & 0 & 0 & 0 \\ 1 & 4 & 6 & 4 & 1 & 0 & 0 \\ 1 & 5 & 10 & 10 & 5 & 1 & 0 \\ 1 & 6 & 15 & 20 & 15 & 6 & 1 \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 6 & 6 & 1 & 0 & 0 & 0 \\ 1 & 10 & 20 & 10 & 1 & 0 & 0 \\ 1 & 15 & 50 & 50 & 15 & 1 & 0 \\ 1 & 21 & 105 & 175 & 105 & 21 & 1 \end{pmatrix}.$$

This last matrix, as expected, is the Narayana triangle A001263. The corresponding γ-matrices for these r-Narayana triangles are, respectively,

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & -1 & 0 & 0 & 0 & 0 & 0 \\ 1 & -3 & 0 & 0 & 0 & 0 & 0 \\ 1 & -6 & 2 & 0 & 0 & 0 & 0 \\ 1 & -10 & 10 & 0 & 0 & 0 & 0 \\ 1 & -15 & 30 & -5 & 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 0 & 0 & 0 & 0 & 0 \\ 1 & 6 & 2 & 0 & 0 & 0 & 0 \\ 1 & 10 & 10 & 0 & 0 & 0 & 0 \\ 1 & 15 & 30 & 5 & 0 & 0 & 0 \end{pmatrix}.$$

This last matrix is A055151. The rows of this triangle are the γ-vectors of the n-dimensional (type A) associahedra [9]. We have seen that its elements are given by

$$\gamma_{n,k} = \sum_{i=0}^{k} \sum_{j=0}^{n} N_{n,j} \binom{2i-1}{i-j} \frac{2j+0^{i+j}}{i+j+0^{i+j}} \text{ If } \left(k=i, 1, \sum_{m=0}^{k-i} \frac{m(-1)^m}{k-i} \binom{n-1+m}{m} \binom{2(k-i)}{k-i-m} \right),$$

where $N_{n,k}$ denotes the (n,k)-th Narayana number A001263.

The relationship between the γ-matrix and the r-Narayana numbers can be further clarified as follows.

Proposition 22. *The generating function of the r-Narayana numbers can be expressed as the Jacobi continued fraction*

$$\mathcal{J}((y+1),(y+1),(y+1),\ldots;ry,ry,ry,\ldots).$$

The generating function of the corresponding γ-matrix can be expressed as the Jacobi continued fraction

$$\mathcal{J}(1,1,1,\ldots;ry,ry,ry,\ldots).$$

Corollary 23. *The generating function of the r-Narayana numbers is the y-th binomial transform of the generating function of the corresponding γ-matrix.*

$$h^*(x,y) = \frac{1}{1-xy}\gamma^*\left(\frac{x}{1-xy}, y\right).$$

Equivalently, the γ generating function $\gamma^*(x,y)$ is the $(-y)$-th binomial transform of the generating function $h^*(x,y)$:

$$\gamma^*(x,y) = \frac{1}{1+xy}h^*\left(\frac{x}{1+xy}, y\right).$$

6 Pascal-like triangles defined by exponential Riordan arrays

We recall that an exponential Riordan array $[g(x), f(x)]$ [1, 6] is defined by two exponential generating functions

$$g(x) = 1 + g_1\frac{x}{1!} + g_2\frac{x}{2!} + \cdots,$$

and

$$f(x) = \frac{x}{1!} + f_2\frac{x^2}{2!} + \cdots,$$

with its (n,k)-th term $a_{n,k}$ given by

$$a_{n,k} = \frac{n!}{k!}[x^n]g(x)f(x)^k.$$

In the context of Pascal-like matrices, we have that the exponential Riordan array

$$[e^x, x(1+rx/2)],$$

with general term

$$h_{n,k} = \frac{n!}{k!} \sum_{j=0}^{k} \frac{r^j}{(n-k-j)!2^j},$$

is a Pascal-like matrix [3]. This matrix begins

$$\left(\begin{array}{ccccccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & r+2 & 1 & 0 & 0 & 0 & 0 \\ 1 & 3r+3 & 3r+3 & 1 & 0 & 0 & 0 \\ 1 & 6r+4 & 3r^2+12r+6 & 6r+4 & 1 & 0 & 0 \\ 1 & 10r+5 & 15r^2+30r+10 & 15r^2+30r+10 & 10r+5 & 1 & 0 \\ 1 & 15r+6 & 45r^2+60r+15 & 15r^3+90r^2+90r+20 & 45r^2+60r+15 & 15r+6 & 1 \end{array}\right).$$

We have the following result.

Proposition 24. *The γ-matrix of the Pascal-like exponential Riordan array $[e^x, x(1+rx/2)]$ is the matrix with general term*

$$\binom{n}{2k} r^k (2k-1)!!$$

In fact, the generating function of the exponential Riordan array $[e^x, x(1+rx/2)]$ is given by

$$\mathcal{J}(y+1, y+1, y+1, \ldots; ry, 2ry, 3ry, \ldots)$$

while that of its γ-matrix is given by

$$\mathcal{J}(1, 1, 1, \ldots; ry, 2ry, 3ry, \ldots).$$

Proposition 25. *The generating function of the γ-matrix of the Pascal-like exponential Riordan array $[e^x, x(1+rx/2)]$ has generating function*

$$e^{x(1+rxy/2)}.$$

Proof. By the theory of exponential Riordan arrays, the generating function of the Riordan array $[e^x, x(1+rx/2)]$ is given by

$$e^x e^{xy(1+rx/2)}.$$

Taking the $(-y)$-th binomial transform of this, we obtain

$$e^{x(1+rxy/2)}.$$

∎

Example 26. For $r = 1$, we get the γ-matrix that begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 0 & 0 & 0 & 0 & 0 \\ 1 & 6 & 3 & 0 & 0 & 0 & 0 \\ 1 & 10 & 15 & 0 & 0 & 0 & 0 \\ 1 & 15 & 45 & 15 & 0 & 0 & 0 \end{pmatrix}.$$

This is A100861, the triangle of Bessel numbers that count the number of k-matchings of the complete graph $K(n)$. The corresponding Pascal-like matrix begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 3 & 1 & 0 & 0 & 0 & 0 \\ 1 & 6 & 6 & 1 & 0 & 0 & 0 \\ 1 & 10 & 21 & 10 & 1 & 0 & 0 \\ 1 & 15 & 55 & 55 & 15 & 1 & 0 \\ 1 & 21 & 120 & 215 & 120 & 21 & 1 \end{pmatrix}.$$

This is A100862, which counts the number of k-matchings of the corona $K'(n)$ of the complete graph $K(n)$ and the complete graph $K(1)$.

Example 27. For $r = 2$, we obtain the γ-matrix that begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 2 & 0 & 0 & 0 & 0 & 0 \\ 1 & 6 & 0 & 0 & 0 & 0 & 0 \\ 1 & 12 & 12 & 0 & 0 & 0 & 0 \\ 1 & 20 & 60 & 0 & 0 & 0 & 0 \\ 1 & 30 & 180 & 120 & 0 & 0 & 0 \end{pmatrix}.$$

This is A059344, where row n consists of the nonzero coefficients of the expansion of $2^n x^n$ in terms of Hermite polynomials with decreasing subscripts. The corresponding Pascal-like matrix begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 1 & 4 & 1 & 0 & 0 & 0 & 0 \\ 1 & 9 & 9 & 1 & 0 & 0 & 0 \\ 1 & 16 & 42 & 16 & 1 & 0 & 0 \\ 1 & 25 & 130 & 130 & 25 & 1 & 0 \\ 1 & 36 & 315 & 680 & 315 & 36 & 1 \end{pmatrix}.$$

The row sums of this matrix are given by A000898, the number of symmetric involutions of $[2n]$ (Deutsch).

7 Conclusion

It is the case that the set of Pascal-like matrices defined by Riordan arrays is a restricted one. Nevertheless, we hope that this note indicates that they have interesting properties, including in particular their generating γ-matrices. In the case of Pascal-like matrices defined by ordinary Riordan arrays, we have seen that be reverting them, we find additional (signed) Pascal-like triangles, including triangles of Narayana type. The γ-matrices of these new triangles are again the reversions of the original triangles' γ-matrices.

We have also shown that stretched Riordan arrays play a useful role, and in particular can lead to further (non-Riordan) Pascal-like matrices. We have also found it useful to use Riordan array techniques to find an explicit closed form formula for the elements $\gamma_{n,k}$ of the γ-matrix in terms of $h_{n,k}$.

References

[1] P. Barry, *Riordan Arrays: A Primer*, Logic Press, 2017.

[2] P. Barry, Continued fractions and transformations of integer sequences, *J. Integer Sequences*, **12** (2009), Article 09.7.6.

[3] P. Barry, On a family of generalized Pascal triangles defined by exponential Riordan array, *J. Integer Sequences*, **10** (2007), Article 07.3.5.

[4] P. Barry, On integer-sequence-based constructions of generalized Pascal triangles, *J. Integer Sequences*, **9** (2006), Article 06.2.4.

[5] C. Corsani, D. Merlini, and R. Sprugnoli, Left-inversion of combinatorial sums, *Discrete Math.*, **180** (1998), 107–122.

[6] E. Deutsch and L. Shapiro, Exponential Riordan arrays, Lecture Notes, Nankai University, 2004, available electronically at `http://www.combinatorics.net/ppt2004/Louis%20W.%20Shapiro/shapiro.htm`.

[7] S. R. Gal, Real root conjecture fails for five and higher dimensional spheres, *Discrete Comput. Geom.*, **34** (2005), 269–284.

[8] Tian-Xiao He and L. W. Shapiro, Fuss-Catalan matrices, their weighted sums, and stabilizer subgroups of the Riordan group, *Linear Algebra Appl.*, **552** (2017), 25–42.

[9] K. Petersen, *Eulerian Numbers*, Birkhäuser, 2015.

[10] L. Shapiro, A survey of the Riordan group, available electronically at `http://www.combinatorics.cn/activities/Riordan%20Group.pdf`. Center for Combinatorics, Nankai University, 2018.

[11] L. W. Shapiro, S. Getu, W.-J. Woan, and L. C. Woodson, The Riordan group, *Discr. Appl. Math.* **34** (1991), 229–239.

[12] L. W. Shapiro, W.-J. Woan, and S. Getu, Runs, slides and moments, *SIAM J. Alg. Discrete Methods*, **4** (1983), 459–466.

[13] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*. Published electronically at `https://oeis.org`, 2018.

[14] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, *Notices Amer. Math. Soc.*, **50** (2003), 912–915.

[15] R. P. Stanley, f-vectors and h-vectors of simplicial posets, *J. Pure Appl. Algebra*, **71** (1991), 319–331.

[16] H. S. Wall, *Analytic Theory of Continued Fractions*, AMS Chelsea Publishing, 2001.

2010 *Mathematics Subject Classification*: Primary 11B83; Secondary 33C45, 42C-5, 15B36, 15B05, 14N10, 11C20.
Keywords: Gamma vector, Pascal-like triangle, Riordan array, Narayana number, Eulerian number, associahedron, permutahedron.

(Concerned with sequences A000108, A000898, A001263, A001591, A007318, A008288, A008292, A055151, A059344, A077938, A100861, A100862, A101280, and A271875.)

Received April 13 2018; revised version received December 19 2018. Published in *Journal of Integer Sequences*, December 19 2018.

Return to Journal of Integer Sequences home page.