124 (1999) MATHEMATICA BOHEMICA No. 2–3, 149–166

POSITIVE SOLUTIONS OF CRITICAL QUASILINEAR ELLIPTIC EQUATIONS IN $\mathbb{R}^N$

Paul A. Binding, Calgary, Pavel Drábek, Plzeň, Yin Xi Huang, Memphis

(Received September 25, 1998)

Dedicated to Professor Alois Kufner on the occasion of his 65th birthday

Abstract. We consider the existence of positive solutions of

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u + \alpha h(x)|u|^{q-2}u + f(x)|u|^{p^*-2}u \tag{1}$$

in $\mathbb{R}^N$, where $\lambda, \alpha \in \mathbb{R}$, $1 < p < N$, $p^* = Np/(N-p)$, the critical Sobolev exponent, and $1 < q < p^*$, $q \neq p$. Let $\lambda_1^+ > 0$ be the principal eigenvalue of

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u \quad \text{in } \mathbb{R}^N, \qquad \int_{\mathbb{R}^N} g(x)|u|^p > 0, \tag{2}$$

with $u_1^+ > 0$ the associated eigenfunction. We prove that, if $\int_{\mathbb{R}^N} f|u_1^+|^{p^*} < 0$, $\int_{\mathbb{R}^N} h|u_1^+|^q > 0$ if $1 < q < p$ and $\int_{\mathbb{R}^N} h|u_1^+|^q < 0$ if $p < q < p^*$, then there exist $\lambda^* > \lambda_1^+$ and $\alpha^* > 0$, such that for $\lambda \in [\lambda_1^+, \lambda^*)$ and $\alpha \in [0, \alpha^*)$, (1) has at least one positive solution.

Keywords: the p-Laplacian, positive solutions, critical exponent

MSC 2000: 35J70, 35P30

1. Introduction

We study the existence of positive solutions to the following problem in $\mathbb{R}^N$

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u + \alpha h(x)|u|^{q-2}u + f(x)|u|^{p^*-2}u, \tag{1.1}$$

Research of the authors was supported by NSERC of Canada, Grant # 201/97/0395 of the Grant Agency of the Czech Republic, a University of Memphis Faculty Research Grant, and NATO Collaborative Research Grant OUTR.CRG 961190

where $\lambda, \alpha \in \mathbb{R}$, $1 < p < N$, $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2}\nabla u)$ is the p-Laplacian, $p^* = Np/(N-p)$, $1 < q < p^*$, $q \neq p$, f, g and h satisfy $g^+ \not\equiv 0$, $f^\pm \not\equiv 0$, $h^+ \not\equiv 0$, and other conditions. The problem is closely related to the following eigenvalue problem,

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u \quad \text{in } \mathbb{R}^N, \qquad \int_{\mathbb{R}^N} g(x)|u|^p > 0. \tag{1.2}_\lambda$$

It is known that $(1.2)_\lambda$ has an eigenvalue $\lambda_1^+ > 0$ associated with a positive eigenfunction u_1^+ (see [6]).

Equations involving critical Sobolev exponents have been studied extensively, and there exists a large body of literature. We refer to [5] and the references therein. Specifically, Swanson and Yu [12] studied $(1.1)_\lambda$ for the case $\lambda \in (0, \lambda_1^+)$ and $p < q < p^*$. It is shown in [12] that if $g \geqslant 0$, $g \in L^{N/p}(\mathbb{R}^N)$, $f \geqslant 0$, and $h \geqslant h_0 > 0$ in $\mathbb{R}^N$, then $(1.1)_\lambda$ has a positive solution if $\lambda \in (0, \lambda_1^+)$. Noussair, Swanson and Yang [11] investigated the problem

$$-\Delta_m u = p(x)u^\tau + q(x)u^\gamma$$

on an open connected smooth domain, where $2 \leqslant m < N$, $m - 1 < \gamma < \tau$, and $\tau + 1 = Nm/(N-m)$. The existence of at least one positive solution was obtained for both p and q nonnegative and satisfying other local conditions. More recently Noussair and Swanson [10] considered

$$-\Delta u = p|u|^{\tau-2}u + q|u|^{\gamma-2}u \quad \text{in } \mathbb{R}^N, \tag{1.3}$$

where $2 < \gamma < \tau = 2N/(N-2)$, and showed, under suitable assumptions, including nonnegativity of p and q, that (1.3) has two positive decaying solutions. The existence of two positive solutions of $(1.1)_\lambda$ was studied for the case $p < q < p^*$ and $f \equiv 0$ in [4], and for the case $h(x) \equiv 0$ in [5]. Various forms of the equation

$$-\Delta_p u + a(x)|u|^{p-2}u = \beta h(x)|u|^{q-2}u + k(x)|u|^{p^*-2}u \quad \text{in } \mathbb{R}^N \tag{1.4}$$

are treated by Alves, Gonçalves and Miyagaki in [1], [2] and [7], where a, h and k are nonnegative, $1 < q \neq p$, $q < p^*$, and $\beta \geqslant 0$. The existence of nonnegative solutions was obtained via Mountain Pass arguments. Specifically, [1] deals with the case $1 < q < p$, $a \equiv 0$, $k \equiv 1$; [2] the case $1 < q < p$ and $\beta = 1$; and [7] the case $a \equiv 0$, $\beta = 1$, $k \equiv 1$, and $1 < q < p^*$, $q \neq p$, $p \geqslant 2$.

In this paper we are mainly concerned with the situation where $\lambda \geqslant \lambda_1^+$. We note that, for $\lambda \in (0, \lambda_1^+)$, the functional $\int_{\mathbb{R}^N}(|\nabla u|^p - \lambda g|u|^p)$ is always positive for $u \neq 0$, so one can use a Mountain Pass type argument to show that $(1.1)_\lambda$ has a positive solution. Assuming $h > 0$ in some open set in $\mathbb{R}^N$, one can even prove the

existence of two positive solutions by first finding a local nonzero minimizer of the associated functional and then using the Mountain Pass Theorem to find a saddle point. This is the approach used in [1], [2] and [7]. For $\lambda \geqslant \lambda_1^+$, the situation is different. The problem is that in this case, the functional $\int_{\mathbb{R}^N}(|\nabla u|^p - \lambda g|u|^p)$ is no longer positive definite. Even a local minimizer is difficult to find. Specifically, for $\lambda > \lambda_1^+$, $\int_{\mathbb{R}^N}(|\nabla u|^p - \lambda g|u|^p)$ will always approach $-\infty$ as $\|u\| \to \infty$ in the direction of u_1^+, while it can achieve positive values in other directions. For $\lambda = \lambda_1^+$, $\int_{\mathbb{R}^N}(|\nabla u|^p - \lambda g|u|^p)$ will always be zero in the direction of u_1^+. This destroys the Mountain Pass structure. Here we use a procedure devised by Tarantello [13] and further utilized in [4] and [5]. The conditions

$$\int_{\mathbb{R}^N} f(u_1^+)^{p^*} < 0, \quad \int_{\mathbb{R}^N} h(u_1^+)^q > 0, \tag{1.5}$$

$$\int_{\mathbb{R}^N} f(u_1^+)^{p^*} < 0, \quad \int_{\mathbb{R}^N} h(u_1^+)^q < 0, \tag{1.6}$$

are essential in our presentation. Under further related local conditions on g, h and f, we can prove the existence of positive solutions of $(1.1)_\lambda$.

Main Result. *Assume* (1.5) *if* $1 < q < p$ *and* (1.6) *if* $p < q < p^*$. *Then there exist* $\lambda^* > \lambda_1^+$ *and* $\alpha^* > 0$, *such that for any* $\lambda \in [\lambda_1^+, \lambda^*)$ *and* $\alpha \in [0, \alpha^*)$, $(1.1)_\lambda$ *has a positive solution (see Theorems* 3.8 *and* 4.7 *for precise assumptions on* f, g *and* h*).*

In our setting, g and h are allowed more flexibility than in [1], [2] and [7], e.g., they may or may not change sign. But (1.5) forces f to change sign, and (1.6) forces both f and h to change sign. We note that here we need an additional condition that α is small enough. While this is the case for $1 < q < p$ in [1], [2] and [7], no such smallness restriction is postulated to h in [7] and [12], for the case $p < q < p^*$.

This paper is organized as follows: In Section 2 we study the geometric structure of certain solution manifolds of the associated functional. Section 3 provides the proof of the existence result for the case $1 < q < p$. The case $p < q < p^*$ is discussed in Section 4.

2. Geometry of the solution manifolds for $1 < q < p$

We collect our basic assumptions and recall some known results. We assume throughout this paper that $1 < p < N$, $p^* = Np/(N-p)$, $1 < q < p^*$ and $q \neq p$. We also assume

(g0) $g(x) = g^+(x) - g^-(x)$, $g^+, g^- \geqslant 0$, and $g^+ \in L^\infty_{\mathrm{loc}}(\mathbb{R}^N) \cap L^{N/p}(\mathbb{R}^N)$, $g^- \in L^\infty_{\mathrm{loc}}(\mathbb{R}^N)$,

(h0) $h \in L^\infty_{\rm loc}(\mathbb{R}^N) \cap L^Q(\mathbb{R}^N)$, where $Q = Np[Np - q(N-p)]^{-1}$.

Let

$$\omega(x) = \frac{1}{(1+|x|)^p},\ x \in \mathbb{R}^N,$$
$$w(x) = \max\{g^-(x), \omega(x)\} > 0,\ x \in \mathbb{R}^N.$$

Let V be the completion of $C_0^\infty(\mathbb{R}^N)$ with respect to the norm $\|\cdot\|$ defined by

$$\|u\| = \left(\int |\nabla u|^p + \int w(x)|u|^p \right)^{1/p}.$$

Here and henceforth the integrals are taken on $\mathbb{R}^N$ unless otherwise stated. Then V is a uniformly convex Banach space. In this paper $\|\cdot\|_p$ will denote the usual L^p norm, and $D^{1,p}(\mathbb{R}^N)$ the completion of $C_0^\infty(\mathbb{R}^N)$ with respect to the norm

$$\|u\|_D = \left(\int |\nabla u|^p \right)^{1/p}.$$

Note that since $V \subset D^{1,p}(\mathbb{R}^N)$, a weakly convergent sequence in V is also weakly convergent in $D^{1,p}(\mathbb{R}^N)$. By Hardy's inequality, $D^{1,p}(\mathbb{R}^N)$ is embedded continuously in $L^p(\mathbb{R}^N, \omega(x))$, so a strongly convergent sequence in $D^{1,p}(\mathbb{R}^N)$ is also strongly convergent in $L^p(\mathbb{R}^N, \omega(x))$.

Throughout this paper the function f is always assumed to satisfy

(f0) $f^\pm \not\equiv 0$ and $f(x) \in L^\infty(\mathbb{R}^N) \cap C(\mathbb{R}^N)$.

We have (from Lemma 2.3 of [6]):

Proposition 2.1. *Assume the above conditions are satisfied. Then there exists a unique, simple isolated eigenvalue $\lambda_1^+ > 0$, such that the eigenvalue problem $(1.2)_\lambda$ has a positive eigenfunction $u_1^+ \in V$ associated with λ_1^+.*

Next we introduce the following functional

$$I_\lambda(u) = \frac{1}{p}\int (|\nabla u|^p - \lambda g|u|^p) - \frac{\alpha}{q}\int h|u|^q - \frac{1}{p^*}\int f|u|^{p^*}. \tag{2.1}$$

It is clear that the functional I_λ is well defined on V. Obviously a critical point of I_λ in V is a (weak) solution of $(1.1)_\lambda$. We can always assume that critical points of I_λ are nonnegative functions since I_λ is an even functional. For simplicity, we will assume in the sequel that $\alpha > 0$, for the case $\alpha = 0$ has been covered in [5].

Define

$$J_\lambda(u) = \int (|\nabla u|^p - \lambda g|u|^p),$$
$$\Lambda_\lambda = \{u \in V\colon\ \Psi_\lambda(u) := \langle I'_\lambda(u), u\rangle = 0\}$$
$$= \left\{u \in V\colon\ J_\lambda(u) = \alpha \int h|u|^q + \int f|u|^{p^*}\right\},$$

and

$$\Lambda_\lambda^- = \{u \in \Lambda_\lambda\colon\ \langle \Psi'_\lambda(u), u\rangle < 0\}. \tag{2.2}$$

We list the following equivalent expressions of this set.

$$\begin{aligned}\Lambda_\lambda^- &= \left\{u \in \Lambda_\lambda\colon\ (p-q)J_\lambda(u) < (p^*-q)\int f|u|^{p^*}\right\}\\ &= \left\{u \in \Lambda_\lambda\colon\ (p^*-p)J_\lambda(u) > \alpha(p^*-q)\int h|u|^q\right\} \qquad (2.3)\\ &= \left\{u \in \Lambda_\lambda\colon\ \alpha(p-q)\int h|u|^q < (p^*-p)\int f|u|^{p^*}\right\}.\end{aligned}$$

We note that it is not entirely clear whether Λ_λ^- is nonempty for general g, h and f. To show that $\Lambda_\lambda^- \neq \emptyset$, we introduce other conditions on g, f and h.

(f1) $f(0) = \|f\|_\infty$ and for some $r > 0$, $f(x) > 0$ for $x \in B(0, 2r)$,
(h1) $h(x) \geqslant h_0 > 0$ in $B(0, 2r)$,
(g1) $g(x) \geqslant g_0 > 0$ in $B(0, 2r)$.

Lemma 2.2. *Suppose* (f0), (f1), (g0), (g1), (h0) *and* (h1) *hold. Then for* $\lambda > 0$ *in any bounded interval, there exists* $\alpha_1 > 0$ *such that* $\Lambda_\lambda^- \neq \emptyset$ *provided* $\alpha \in (0, \alpha_1)$.

P r o o f. Define, for $\varepsilon > 0$,

$$u_\varepsilon(x) = \frac{\psi(x)}{(\varepsilon + |x|^{p/(p-1)})^{(N-p)/p}}, \quad v_\varepsilon(x) = \frac{u_\varepsilon(x)}{\|u_\varepsilon(x)\|_{p^*}},$$

where $\psi \in C_0^\infty(B(0, 2r))$ is such that $0 \leqslant \psi(x) \leqslant 1$ and $\psi(x) \equiv 1$ on $B(0, r)$.

Consider for $t > 0$,

$$\Psi_\lambda(tv_\varepsilon) = t^p J_\lambda(v_\varepsilon) - \alpha t^q \int h|v_\varepsilon|^q - t^{p^*}\int f|v_\varepsilon|^{p^*}.$$

Let $s_\alpha(t) = at^p - \alpha bt^q - ct^{p^*}$, with $a = J_\lambda(v_\varepsilon)$, $b = \int h|v_\varepsilon|^q$ and $c = \int f|v_\varepsilon|^{p^*}$. It is clear that $b > 0$, $c > 0$. By continuous dependence of the principal eigenvalue

on the domain, $a > 0$ for $\varepsilon > 0$ small enough. Fix this ε so $a,\ b,\ c$ are fixed and let α vary. One easily sees that $s_\alpha(t) \to -\infty$ as $t \to \infty$. Moreover, as $b \to 0$, $s_\alpha(t) \to s_0(t) := at^p - ct^{p^*}$ in C^1 with respect to t. Let t_0 be such that $s_0(t_0) = 0$ and $s_0(t) > 0$ for $t < t_0$. Then $s_0'(t_0) < 0$. By C^1 convergence of s_α to s_0, we easily conclude that there exist $\alpha_1 > 0$ and $\tau > 0$, such that if $0 < \alpha < \alpha_1$, $s_\alpha(t_\alpha) = 0$ and $s_\alpha'(t_\alpha) < 0$ for some $t_\alpha \in (t_0 - \tau, t_0 + \tau)$, that is, $t_\alpha v_\varepsilon \in \Lambda_\lambda^-$. This completes the proof. □

Next we study the geometry of the set Λ_λ^- for $\lambda > 0$. We will seek a critical point of I_λ on Λ_λ^-. Observe that for any $u \in \Lambda_\lambda$,

$$\begin{aligned} I_\lambda(u) &= \frac{1}{N}\int f|u|^{p^*} + \alpha\Big(\frac{1}{p} - \frac{1}{q}\Big)\int h|u|^q \\ &= \frac{1}{N}J_\lambda(u) - \alpha\Big(\frac{1}{q} - \frac{1}{p^*}\Big)\int h|u|^q. \end{aligned} \tag{2.4}$$

We also assume that $\|u_1^+\| = 1$.

The next lemma requires the following conditions.

(f2) $\int f(u_1^+)^{p^*} < 0$.
(h2) $\int h(u_1^+)^q > 0$.

Lemma 2.3. *Assume $p > q$, (f0), (f1), (f2), (g0), (g1), (h0), (h1) and (h2) hold. Then there exist $\lambda^* > \lambda_1^+$ and $\alpha_2 > 0$ with $\alpha_2 \leqslant \alpha_1$, such that for any $\overline{\lambda} \in (0, \lambda_1^+)$, there exists $\sigma > 0$, such that for any $\lambda \in [\overline{\lambda}, \lambda^*)$ and $\alpha \in (0, \alpha_2)$, we have $J_\lambda(u) \geqslant \sigma\|u\|^p$ for any $u \in \Lambda_\lambda^-$.*

P r o o f. We argue by contradiction. Suppose there exist λ_n, α_n and $u_n \in \Lambda_{\lambda_n}^-$ such that

$$\alpha_n \to 0, \quad \lambda_n \to \widehat{\lambda} \in [\overline{\lambda}, \lambda_1^+], \quad J_{\lambda_n}(u_n) < \frac{1}{n}\|u_n\|^p. \tag{2.5}$$

We explicitly note that here $\Lambda_{\lambda_n}^-$ also depends on α_n. Let $v_n = u_n/\|u_n\|$. Without loss of generality we may assume $v_n \to v_0$ weakly in V. Then we have $\int g^+|v_n|^p \to \int g^+|v_0|^p$ by compactness. We then derive by weak lower semicontinuity of the norm that

$$\begin{aligned} 0 &\leqslant \int |\nabla v_0|^p - \widehat{\lambda}\int g^+|v_0|^p + \widehat{\lambda}\int g^-|v_0|^p \\ &\leqslant \liminf_{n\to\infty}\bigg(\int |\nabla v_n|^p - \lambda_n\int g^+|v_n|^p + \lambda_n\int g^-|v_n|^p\bigg) \\ &= \liminf_{n\to\infty} J_{\lambda_n}(v_n) \leqslant \lim_{n\to\infty}\frac{1}{n} = 0. \end{aligned} \tag{2.6}$$

There are two possibilities: (1) $v_0 = 0$, and (2) $v_0 = ku_1^+$ for some $k \neq 0$, and $\widehat{\lambda} = \lambda_1^+$. If $v_0 = 0$, it follows from (2.6) that $\int |\nabla v_n|^p \to 0$ and $\int g^-|v_n|^p \to 0$. Thus $v_n \to 0$ in V, contradicting $\|v_n\| = 1$. If $v_0 = ku_1^+$ for some $k \neq 0$, and $\widehat{\lambda} = \lambda_1^+$, then we have, by the weak convergence of v_n to ku_1^+ and (2.6),

$$\begin{aligned}\lambda_1^+ \int g^+|ku_1^+|^p &= \int (|\nabla ku_1^+|^p + \lambda_1^+ g^-|ku_1^+|^p) \\ &\leqslant \liminf_{n\to\infty} \int |\nabla v_n|^p + \liminf_{n\to\infty} \lambda_n \int g^-|v_n|^p \\ &\leqslant \liminf_{n\to\infty} \int (|\nabla v_n|^p + \lambda_n g^-|v_n|^p) \\ &= \lim_{n\to\infty} \lambda_n \int g^+|v_n|^p = \lambda_1^+ \int g^+|ku_1^+|^p.\end{aligned}$$

It then follows that

$$\liminf_{n\to\infty} \int |\nabla v_n|^p = \int |\nabla ku_1^+|^p, \quad \liminf_{n\to\infty} \int g^-|v_n|^p = \int g^-|ku_1^+|^p.$$

We deduce that (passing to a subsequence if necessary) $v_n \to ku_1^+$ strongly in V. We then derive from (2.3) that

$$\|u_n\|^{p-p^*} J_{\lambda_n}(v_n) < \frac{p^*-q}{p-q} \int f|v_n|^{p^*} \to \frac{p^*-q}{p-q} \int f|ku_1^+|^{p^*} < 0. \tag{2.7}$$

This contradicts (2.6) if $\|u_n\| \not\to 0$ or $J_{\lambda_n}(u_n) \geqslant 0$. Suppose $\|u_n\| \to 0$ and $J_{\lambda_n}(u_n) < 0$. It follows from (2.3) that $\int h|u_n|^q < 0$. That is, $\int h|v_n|^q \leqslant 0$, which contradicts (h2). This proves the lemma. □

R e m a r k 2.4. For $\lambda \in (0, \lambda_1^+)$, conditions (f2) and (h2) are not needed because $J_\lambda(u) \geqslant 0$ for all u. Assumptions (f2) and (h2) are introduced to compensate for the possibility that $J_\lambda(u)$ is negative.

Lemma 2.5. *Assume $p > q$,* (f0), (f1), (f2), (g0), (g1), (h0), (h1) *and* (h2) *hold. For any $\bar{\lambda} \in (0, \lambda_1^+)$, there exist $\varrho > 0$ and $\alpha^* > 0$ with $\alpha^* \leqslant \alpha_2$, such that for any $\lambda \in [\bar{\lambda}, \lambda^*)$, $\alpha \in (0, \alpha^*)$ and $u \in \Lambda_\lambda^-$, we have $-\langle \Psi'_\lambda(u), u\rangle \geqslant \varrho$.*

P r o o f. We first claim that there exists $\zeta > 0$, independent of λ, such that $\|u\| > \zeta$ for all $u \in \Lambda_\lambda^-$. If this were not true, then for some $u_n \in \Lambda_{\lambda_n}^-$, $\lambda_n \in [\bar{\lambda}, \lambda^*)$, $u_n \to 0$. Dividing (2.3) by $\|u_n\|^p$ we obtain, using Lemma 2.3,

$$0 < \sigma \leqslant J_{\lambda_n}(v_n) < \frac{p^*-q}{p-q} \int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-p} \to 0, \tag{2.8}$$

a contradiction, where $v_n = u_n/\|u_n\|$.

Now, by Young's inequality and Lemma 2.3, for any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that

$$\begin{aligned} -\langle \Psi'_\lambda(u), u\rangle &= (p^* - p)J_\lambda(u) - \alpha(p^* - q)\int h|u|^q \\ &\geqslant (p^* - p)\sigma\|u\|^p - \alpha(p^* - q)\|h\|_Q \cdot \|u\|^q \\ &\geqslant ((p^* - p)\sigma - \varepsilon)\zeta^p - \alpha C_\varepsilon \|h\|_Q^{p/(p-q)}. \end{aligned}$$

The proof is complete. □

Corollary 2.6. *Under the conditions of Lemma* 2.5, *for any* $\bar\lambda \in (0, \lambda_1^+)$, *there exists* $\alpha^* > 0$ *such that* Λ_λ^- *is a closed set for* $\lambda \in [\bar\lambda, \lambda^*)$ *provided* $\alpha \in (0, \alpha^*)$.

3. Proof of existence of solutions for $1 < q < p$

Lemma 3.1. *Assume* (f0), (f1), (f2), (g0), (g1), (h0), (h1) *and* (h2) *hold. Then* I_λ *is bounded below on* Λ_λ^- *for* $\lambda \in (0, \lambda^*)$ *and* $\alpha \in (0, \alpha^*)$, *where* α^* *and* λ^* *are given in Lemma* 2.5.

P r o o f. Suppose for some $u_n \in \Lambda_\lambda^-$, $I_\lambda(u_n) \to -\infty$. Then $\|u_n\| \to \infty$. Since $u_n \in \Lambda_\lambda^-$, $\int f|u_n|^{p^*} > 0$ by (2.3) and Lemma 2.3. Dividing $I_\lambda(u_n)$ by $\|u_n\|^p$ we obtain from (2.4) that

$$\frac{I_\lambda(u_n)}{\|u_n\|^p} = \frac{1}{N}\int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-p} - \alpha\Big(\frac{1}{q} - \frac{1}{p}\Big)\int h|v_n|^q \cdot \|u_n\|^{q-p} \to \ell \leqslant 0,$$

with $v_n = u_n/\|u_n\|$. It then follows that $\int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-p} \to N\ell \leqslant 0$. On the other hand, dividing

$$J_\lambda(u_n) = \int f|u_n|^{p^*} + \alpha\int h|u_n|^q$$

by $\|u_n\|^p$ we obtain, using Lemma 2.3,

$$0 < \sigma \leqslant J_\lambda(v_n) = \int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-p} + \alpha\int h|v_n|^q \cdot \|u_n\|^{q-p} \to N\ell \leqslant 0,$$

a contradiction. So I_λ is bounded below on Λ_λ^-. □

Thus we can define $c_0 = \inf\limits_{\Lambda_\lambda^-} I_\lambda(u)$.

Lemma 3.2. *Assume* (f0), (f1), (f2), (g0), (g1), (h0), (h1) *and* (h2) *hold. Then for any* $\lambda \in (0, \lambda^*)$, $\alpha \in (0, \alpha^*)$, *there exists a minimizing sequence* $\{u_n\} \subset \Lambda_\lambda^-$ *of* I_λ *on* Λ_λ^- *which converges weakly to a solution* u *of* $(1.1)_\lambda$.

P r o o f. We first show that any minimizing sequence of I_λ on Λ_λ^- is bounded. Suppose $\{u_n\}$ is an unbounded minimizing sequence of I_λ on Λ_λ^-. Dividing $I_\lambda(u_n)$ by $\|u_n\|^q$, we conclude that, since $I_\lambda(u_n)$ is bounded, $\int f|u_n|^{p^*} \cdot \|u_n\|^{-q}$ is bounded by (2.4). Thus $J_\lambda(u_n) \cdot \|u_n\|^{-q}$ is also bounded by (2.3). Let $v_n = u_n/\|u_n\|$. Then $\sigma \leqslant J_\lambda(v_n) \to 0$, a contradiction. Thus any minimizing sequence $\{u_n\}$ in Λ_λ^- is bounded.

Since Λ_λ^- is a closed set by Corollary 2.6, it follows from Theorem 4.1 and Remark 4.1 of [9] that we can replace $\{u_n\}$ by another minimizing sequence $\{z_n\} \subset \Lambda_\lambda^-$ such that $\|u_n - z_n\| < 1/n$, and for any $y \in \Lambda_\lambda^-$,

$$I_\lambda(y) > I_\lambda(z_n) - \tfrac{1}{n}\|y - z_n\|. \tag{3.1}$$

We want to show that $I'_\lambda(z_n) \to 0$. Choose w_n of unit norm so that

$$\langle I'_\lambda(z_n), w_n\rangle \geqslant \|I'_\lambda(z_n)\| - o(1)$$

as $n \to \infty$. It will suffice to show that

$$\langle I'_\lambda(z_n), w_n\rangle \to 0. \tag{3.2}$$

For each n, let $g_n(t,s) = \Psi_\lambda(tz_n - sw_n)$. Then $g_n(1,0) = 0$ and

$$\tfrac{\partial g_n}{\partial t} = \langle \Psi'_\lambda(z_n), z_n\rangle \neq 0 \text{ at } t = 1, s = 0.$$

It follows from the C^1 Implicit Function Theorem that for each n, for small enough s, there exists $t_n \in C^1$ so that $\Psi_\lambda(t_n(s)z_n - sw_n) = 0$, i.e. $t_n(s)z_n - sw_n \in \Lambda_\lambda$ and

$$\langle \Psi'_\lambda(z_n), z_n\rangle t'_n(0) - \langle \Psi'_\lambda(z_n), w_n\rangle = 0. \tag{3.3}$$

Since z_n is a bounded sequence, so is $\|\Psi'_\lambda(z_n)\|$, and we then conclude from (3.3) and Lemma 2.5 that

$$t'_n(0) \text{ is uniformly bounded in } n. \tag{3.4}$$

We fix n, and consider $v_n(s) = t_n(s)z_n - sw_n - z_n$. Since $\|w_n\| = 1$, we have

$$\|v_n(s)\| \leqslant |s|(1 + (|t'_n(0) + o(1)|)\|z_n\|) \tag{3.5}$$

as $s \to 0$. Moreover $z_n \in \Lambda_\lambda$ gives $\langle I'_\lambda(z_n), z_n\rangle = 0$, so

$$I_\lambda(z_n) - I_\lambda(t_n(s)z_n - sw_n) = \langle I'_\lambda(z_n), -v_n(s)\rangle + o(v_n(s)) = \langle I'_\lambda(z_n), sw_n\rangle + o(s) \tag{3.6}$$

follows from (3.5). By continuity of $\langle \Psi'_\lambda(u), u\rangle$, we have

$$\langle \Psi'_\lambda(t_n(s)z_n - sw_n), t_n(s)z_n - sw_n\rangle - \langle \Psi'_\lambda(z_n), z_n\rangle \to 0$$

as $s \to 0$. We then conclude from this and Lemma 3.4 that

$$\langle \Psi'_\lambda(t_n(s)z_n - sw_n), t_n(s)z_n - sw_n\rangle < 0$$

for s small enough, so $t_n(s)z_n - sw_n \in \Lambda_\lambda^-$.

Dividing (3.6) by s and using (3.1) with $y = t_n(s)z_n - sw_n$ and (3.5), we obtain

$$|\langle I'_\lambda(z_n), w_n\rangle| \leqslant n^{-1}(1 + (|t'_n(0)|)\|z_n\|) + o(1).$$

Letting $n \to \infty$ we conclude that $\langle I'_\lambda(z_n), w_n\rangle$ tends to zero by boundedness of z_n and (3.4). This establishes (3.2).

Assume now that $z_n \to u$ weakly in V. We have, then, as in the proof of Lemma 3.1 of [5], since $I'_\lambda(z_n) \to 0$, that u is a weak solution of $(1.1)_\lambda$, i.e.,

$$-\Delta_p u = \lambda g|u|^{p-2}u + \alpha h|u|^{q-2}u + f|u|^{p^*-2}u$$

in V. This proves the lemma. □

Thus we have obtained a weak solution of $(1.1)_\lambda$. To show that this solution is nontrivial, we need some preparation. Let S be the best Sobolev constant, i.e.,

$$S = \inf\left\{\frac{\|\nabla u\|_p^p}{\|u\|_{p^*}^p} : u \in W_0^{1,p}(\mathbb{R}^N) \setminus \{0\}\right\},$$

and $S_0 = S^{N/p}\|f\|_\infty^{(p-N)/p}/N$. Recall the concentration-compactness principle of P. L. Lions ([8]).

Proposition 3.3. *Let $\{u_n\}$ converge weakly to u in $D^{1,p}(\mathbb{R}^N)$ such that $|u_n|^{p^*}$ and $|\nabla u_n|^p$ converge weakly to nonnegative measures ν and μ on $\mathbb{R}^N$ respectively. Then, for some at most countable set J, we have*

(i) $\nu = |u|^{p^*} + \sum_{j\in J} \nu_j \delta_{x_j}$;

(ii) $\mu \geqslant |\nabla u|^p + \sum_{j\in J} \mu_j \delta_{x_j}$;

(iii) $S\nu_j^{p/p^*} \leqslant \mu_j$,

where $x_j \in \mathbb{R}^N$, δ_{x_j} is the Dirac measure at x_j, and ν_j and μ_j are nonnegative constants.

Lemma 3.4. *Assume* (f0), (f1), (f2), (g0), (g1), (h0), (h1) *and* (h2) *hold. For* $\lambda \in [0, \lambda^*)$ *and* $\alpha \in (0, \alpha^*)$, *any minimizing sequence* $\{u_n\}$ *of* I_λ *on* Λ_λ^- *satisfying* $I_\lambda(u_n) < S_0$ *either converges strongly to a solution* $u \in \Lambda_\lambda^-$, *hence* $u \neq 0$, *or converges weakly to a nontrivial solution* $u \in \Lambda_\lambda$.

P r o o f. Let $\{u_n\}$ be such a minimizing sequence. We can assume without loss of generality that $\{u_n\}$ is bounded (cf. Lemma 3.2).

Assume that $u_n \to u$ weakly in V. We conclude as in the proof of Proposition 2.3 of [5] that

$$-\Delta_p u = \lambda g|u|^{p-2}u + f|u|^{p^*-2}u + \alpha h|u|^{q-2}u$$

in V, that is, $I'_\lambda(u) = 0$ and hence $u \in \Lambda_\lambda$.

Suppose that $u_n \not\to u$ strongly in V and $u = 0$. Then for some j, ν_j given by Proposition 3.3 is not zero. We obtain, using the fact that $\int h|u_n|^q \to 0$ (cf. Proposition 2.3 of [5]),

$$\begin{aligned} S_0 > I_\lambda(u_n) &= \frac{1}{N}\int f|u_n|^{p^*} + \alpha\Big(\frac{1}{p} - \frac{1}{q}\Big)\int h|u_n|^q \\ &\geqslant \frac{1}{N}\sum_{j\in J} f(x_j)\nu_j \geqslant \frac{1}{N}\sum_{j\in J}\frac{S^{N/p}}{f(x_j)^{(N-p)/p}} \geqslant S_0, \end{aligned}$$

a contradiction. Here we used the facts that $f(x_j)\nu_j = \mu_j$ and $\nu_j \geqslant (S/f(x_j))^{N/P}$, which follow from the proof of Proposition 2.4 of [5]. This proves the lemma. □

We need more conditions on f. Assume

(f3) for $x \in B(0, 2r)$,

$$f(x) = f(0) + o(|x|^k), \quad k = \frac{N}{q} \text{ if } q \geqslant \frac{N(p-1)}{N-p}, \quad k = \frac{N-p}{p-1} \text{ if } q < \frac{N(p-1)}{N-p},$$

or

(f3)′ for $x \in B(0, 2r)$,

$$f(x) = f(0) + o(|x|^\delta), \quad \delta = \frac{N-p}{p-1}.$$

Lemma 3.5. *Assume* (f0), (f1), (f2), (f3) or (f3)′, (g0), (g1), (h0) *and* (h1) *hold. Then for* $\lambda > 0$, *and* $\varepsilon > 0$ *small enough, we have*

$$\sup_{t\geqslant 0} I_\lambda(tv_\varepsilon) < \frac{1}{N}S^{N/p}\|f\|_\infty^{(p-N)/p} = S_0, \tag{3.7}$$

where v_ε *is given in the proof of Lemma* 2.2.

P r o o f. Our proof is similar to that of Lemma 4.1 of [12]. Recall for $\varepsilon > 0$,

$$u_\varepsilon(x) = \frac{\psi(x)}{(\varepsilon + |x|^{p/(p-1)})^{(N-p)/p}}, \quad v_\varepsilon(x) = \frac{u_\varepsilon(x)}{\|u_\varepsilon(x)\|_{p^*}},$$

where $\psi \in C_0^\infty(B(0,2r))$ is such that $0 \leqslant \psi(x) \leqslant 1$ and $\psi(x) \equiv 1$ on $B(0,r)$.

Calculations show that (cf. the proof of Lemma 5.6 in [5])

$$\int |v_\varepsilon|^t = \begin{cases} K\varepsilon^{(N(p-t)+tp)(p-1)/p^2}, & \text{if } t > \frac{p^*}{p'}, \\ K\varepsilon^{N(p-1)/p^2}|\ln\varepsilon|, & \text{if } t = \frac{p^*}{p'}, \\ K\varepsilon^{t(N-p)/p^2}, & \text{if } t < \frac{p^*}{p'}, \end{cases} \tag{3.8}$$

and

$$\int |\nabla v_\varepsilon|^t = \begin{cases} K'\varepsilon^{N(p-t)(p-1)/p^2}, & \text{if } t > \frac{N(p-1)}{N-1}, \\ K'\varepsilon^{t(N-p)/p^2}|\ln\varepsilon|, & \text{if } t = \frac{N(p-1)}{N-1}, \\ K'\varepsilon^{t(N-p)/p^2}, & \text{if } t < \frac{N(p-1)}{N-1}. \end{cases} \tag{3.9}$$

In particular, we have

$$\int |v_\varepsilon| = \begin{cases} K\varepsilon^{(N-p)/p^2}, & \text{if } p > \frac{2N}{N+1}, \\ K\varepsilon^{(N-p)/p^2}|\ln\varepsilon|, & \text{if } p = \frac{2N}{N+1}, \\ K\varepsilon^{(N(p-1)+p)(p-1)/p^2}, & \text{if } p < \frac{2N}{N+1}, \end{cases} \tag{3.10}$$

$$\int |v_\varepsilon|^p = \begin{cases} K\varepsilon^{p-1} & p^2 < N, \\ K\varepsilon^{p-1}|\ln\varepsilon| & p^2 = N, \\ K\varepsilon^{(N-p)/p}, & p^2 > N. \end{cases} \tag{3.11}$$

$$\int v_\varepsilon^{p^*} = 1,$$

$$\int |\nabla v_\varepsilon| = \begin{cases} K'\varepsilon^{(N-p)/p^2}, & \text{if } p > \frac{2N-1}{N}, \\ K'\varepsilon^{(N-p)/p^2}|\ln\varepsilon|, & \text{if } p = \frac{2N-1}{N}, \\ K'\varepsilon^{N(p-1)^2/p^2}, & \text{if } p < \frac{2N-1}{N}, \end{cases} \tag{3.12}$$

$$\int |\nabla v_\varepsilon|^{p-1} = K'\varepsilon^{(N-p)(p-1)/p^2}, \tag{3.13}$$

and

$$\int |\nabla v_\varepsilon|^p = \frac{\int |\nabla u_\varepsilon|^p}{\|u_\varepsilon\|_{p^*}^p} = \frac{K_1}{K_2} + O(\varepsilon^{(N-p)/p}), \tag{3.14}$$

where $K_1/K_2 = S$.

Note that for $\varepsilon > 0$ small enough, $J_\lambda(v_\varepsilon) > 0$, so $I_\lambda(tv_\varepsilon)$ attains its maximum at some $t_\varepsilon \in (0,\infty)$ with $s'(t_\varepsilon) = 0$, where $s(t) = I_\lambda(tv_\varepsilon)$. That is,

$$0 = s'(t_\varepsilon) = t_\varepsilon^{p-1}\Big(\int (|\nabla v_\varepsilon|^p - \lambda g|v_\varepsilon|^p) - \alpha t_\varepsilon^{q-p}\int h|v_\varepsilon|^q - t_\varepsilon^{p^*-p}\int f|v_\varepsilon|^{p^*}\Big).$$

Thus, by (g1) and (h1),

$$t_\varepsilon^{p^*-p} \leqslant \frac{\int |\nabla v_\varepsilon|^p}{f(x^*)\int |v_\varepsilon|^{p^*}},$$

where $f(x^*) = \inf\limits_{x\in B(0,r)} f(x) > 0$. It then follows that t_ε is bounded from above. We may also assume that t_ε is bounded from below, otherwise $I_\lambda(t_\varepsilon v_\varepsilon) \to 0$ as $\varepsilon \to 0$. Now,

$$I_\lambda(t_\varepsilon v_\varepsilon) = \sup_{t\geqslant 0} I_\lambda(tv_\varepsilon) = E(\varepsilon) - F(\varepsilon) + V(\varepsilon), \tag{3.15}$$

where

$$\begin{aligned}
E(\varepsilon) &= \frac{t_\varepsilon^p}{p}\int |\nabla v_\varepsilon|^p - \frac{f(0)t_\varepsilon^{p^*}}{p^*}\int v_\varepsilon^{p^*},\\
F(\varepsilon) &= \frac{\lambda t_\varepsilon^p}{p}\int g v_\varepsilon^p + \alpha\frac{t_\varepsilon^q}{q}\int h v_\varepsilon^q,\\
V(\varepsilon) &= \frac{t_\varepsilon^{p^*}}{p^*}\int (f(0) - f(x))v_\varepsilon^{p^*}.
\end{aligned}$$

The maximum of $ap^{-1}t^p - b(p^*)^{-1}t^{p^*}$ is achieved at $t = (a/b)^{(N-p)/p^2}$ for positive a, b, so

$$E(\varepsilon) \leqslant \Big(\frac{1}{p} - \frac{1}{p^*}\Big)[f(0)]^{(p-N)/p}\Big[\int |\nabla v_\varepsilon|^p\Big]^{N/p}\Big[\int v_\varepsilon^{p^*}\Big]^{-N/p^*} = \frac{1}{N}S^{N/p}\|f\|_\infty^{(p-N)/p}.$$

We also have, for k and δ given in (f3) and (f3)′ respectively,

$$V_\varepsilon = O(\varepsilon^{k(p-1)/p}),\ O(\varepsilon^{\delta(p-1)/p}).$$

Assuming (f3) holds, we estimate, using the fact that t_ε is bounded from below,

$$F(\varepsilon) \geqslant \alpha h_0 \int v_\varepsilon^q = \begin{cases} K\varepsilon^{[N(p-q)+qp](p-1)/p^2}, & \text{if } q > \frac{N(p-1)}{N-p},\\ K\varepsilon^{N(p-1)/p^2}|\ln\varepsilon|, & \text{if } q = \frac{N(p-1)}{N-p},\\ K\varepsilon^{q(N-p)/p^2}, & \text{if } q < \frac{N(p-1)}{N-p}. \end{cases}$$

From (f3) we derive that for $\varepsilon > 0$ small enough, $F(\varepsilon)$ dominates $V(\varepsilon)$. Thus we conclude from the above that, for $\varepsilon > 0$ small enough and $K > 0$,

$$I_\lambda(t_\varepsilon v_\varepsilon) \leqslant \begin{cases} S_0 - K\varepsilon^{[N(p-q)+qp](p-1)/p^2}, & \text{if } q > \frac{N(p-1)}{N-p}, \\ S_0 - K\varepsilon^{N(p-1)/p^2}|\ln\varepsilon|, & \text{if } q = \frac{N(p-1)}{N-p}, \\ S_0 - K\varepsilon^{q(N-p)/p^2}, & \text{if } q < \frac{N(p-1)}{N-p}. \end{cases} \tag{3.16}$$

On the other hand, assume (f3)′ holds. We have

$$F(\varepsilon) \geqslant g_0 \int v_\varepsilon^p = \begin{cases} K\varepsilon^{p-1}, & \text{if } p^2 < N, \\ K\varepsilon^{p-1}|\ln\varepsilon|, & \text{if } p^2 = N, \\ K\varepsilon^{(N-p)/p}, & \text{if } p^2 > N. \end{cases}$$

Since $p - 1 \leqslant (N-p)/p$ for $p^2 \leqslant N$ and $\delta(p-1)/p > (N-p)/p$ for $p^2 > N$ by (f3)′, $F(\varepsilon)$ dominates $V(\varepsilon)$. Again we have

$$I_\lambda(t_\varepsilon v_\varepsilon) \leqslant \begin{cases} S_0 - K\varepsilon^{p-1}, & \text{if } p^2 < N, \\ S_0 - K\varepsilon^{p-1}|\ln\varepsilon|, & \text{if } p^2 = N, \\ S_0 - K\varepsilon^{(N-p)/p}, & \text{if } p^2 > N. \end{cases} \tag{3.16}′$$

The lemma then follows. □

Lemma 3.6. *Assume* (f0), (f1), (f2), (f3) or (f3)′, (g0), (g1), (h0), (h1), (h2), *and* λ *and* α *as in Lemma* 2.5. *Then* $c_0 = \inf_{\Lambda_\lambda^-} I_\lambda(u) < S_0$.

This lemma follows from Lemma 3.5 and the fact that $t_\varepsilon v_\varepsilon \in \Lambda_\lambda^-$ for some $t_\varepsilon > 0$ (cf. the proof of Lemma 2.2). Thus we have proved, via Lemmas 3.4 and 3.6, the existence of a nonnegative solution. The next result shows that the solution is actually positive.

Proposition 3.7. *Let* u *be a nonnegative solution of* $(1.1)_\lambda$ *with* $q \leqslant p^*$. *Then* $u > 0$ *in* $\mathbb{R}^N$.

The proof is essentially as that of Lemma 4.3 of [12] and is omitted.

Now we can state our main result.

Theorem 3.8. *Assume that* (f0), (f1), (f2), (f3) *or* (f3)′, (g0), (g1), (h0), (h1), *and* (h2) *hold. Then there exist* $\lambda^* > \lambda_1^+$ *and* $\alpha^* > 0$, *so that the problem*

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u + \alpha h(x)|u|^{q-2}u + f(x)|u|^{p^*-2}u$$

has at least one positive solution in V *for any* $\lambda \in [\lambda_1^+, \lambda^*)$ *and* $\alpha \in (0, \alpha^*)$.

4. The case $p < q < p^*$ and some remarks

For this case, the set Λ_λ^- is defined as in (2.3). We first have the following result.

Lemma 4.1. *Assume* $\operatorname{supp} f^+ \cap \operatorname{supp} h^+$ *contains an open set. Then* $\Lambda_\lambda^- \neq \emptyset$ *for* $\lambda > 0$.

P r o o f. Suppose $\operatorname{supp} f^+ \cap \operatorname{supp} h^+$ contains an open set B and let $\varphi > 0$ be such that $\operatorname{supp}\varphi \subset B$ with $J_\lambda(\varphi) > 0$. Such φ exists as explained in the proof of Lemma 2.2. For $t > 0$, we have

$$\Psi(t\varphi) = t^p J_\lambda(\varphi) - \alpha t^q \int h\varphi^q - t^{p^*} \int f\varphi^{p^*}.$$

Let again $s(t) = at^p - \alpha b t^q - ct^{p^*}$, with $a = J_\lambda(\varphi) > 0$, $b = \int h\varphi^q > 0$ and $c = \int f\varphi^{p^*} > 0$. Obviously $s(t) > 0$ for $t > 0$ small and $s(t) \to -\infty$ as $t \to \infty$. Suppose $s(t_0) = 0$. Then

$$\begin{aligned} s'(t_0) &= t_0^{q-1}(pat_0^{p-q} - \alpha qb - p^* ct_0^{p^*-q}) \\ &= t_0^{q-1}[\alpha(p-q)b - (p^*-p)ct_0^{p^*-q}] < 0 \end{aligned}$$

since $p < q$. That is, $t_0\varphi \in \Lambda_\lambda^-$. This concludes the proof. □

R e m a r k 4.2. Note that (f1) and (h1) imply that $\operatorname{supp} f^+ \cap \operatorname{supp} h^+$ contains an open set. So we will assume for simplicity in the sequel that (f1) and (h1) hold. We also note that Lemma 4.1 holds if $h \equiv 0$.

Instead of (h2), we need
(h2)′ $\int h|u_1^+|^q < 0$.

Lemma 4.3. *Assume* (f0), (f1), (f2), (g0), (g1), (h0), (h1) *and* (h2)′ *hold. Then there exist* $\lambda_1^* > \lambda_1^+$ *and* $\alpha_1 > 0$ *such that for any* $\overline{\lambda} \in (0, \lambda_1^+)$, *there exists* $\sigma > 0$, *such that for any* $\lambda \in [\overline{\lambda}, \lambda_1^*)$ *and* $\alpha \in (0, \alpha_1)$, $J_\lambda(u) \geqslant \sigma\|u\|^p$ *for any* $u \in \Lambda_\lambda^-$.

P r o o f. If the conclusion were false, there would exist λ_n, α_n and $u_n \in \Lambda_{\lambda_n}^-$ such that

$$\alpha_n \to 0, \quad \lambda_n \to \widehat{\lambda} \in [\overline{\lambda}, \lambda_1^+], \quad J_{\lambda_n}(u_n) < \frac{1}{n}\|u_n\|^p.$$

As in the proof of Lemma 2.3, we conclude that $v_n = u_n/\|u_n\| \to ku_1^+$ for some $k \neq 0$. Then instead of (2.7) we have

$$\|u_n\|^{p-p^*} J_{\lambda_n}(v_n) > \frac{p^*-q}{p-q} \int f|v_n|^{p^*} \to \frac{p^*-q}{p-q} \int f|ku_1^+|^{p^*} > 0, \tag{4.1}$$

since $p < q$ and $\int f|ku_1^+|^{p^*} < 0$ by (f2). If $\|u_n\| \not\to 0$, then (4.1) contradicts the fact that $J_{\lambda_n}(v_n) \to 0$. If $\|u_n\| \to 0$, (4.1) implies that $J_{\lambda_n}(v_n) > 0$. We then have

$$\alpha_n \int h|v_n|^q + \int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-q} = J_{\lambda_n}(v_n) \cdot \|u_n\|^{p-q} > 0. \tag{4.2}$$

Note that $\int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-q} \to 0$ since $\|u_n\| \to 0$. Inequality (4.2) then implies $\int h|v_n|^q > 0$, contradicting (h2)$'$. Thus the lemma is proved. □

R e m a r k 4.4. We point out that Lemma 4.3 holds if $h \equiv 0$.

The reason is that instead of (4.2), we now have

$$\int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-p} = J_{\lambda_n}(v_n) > 0. \tag{4.2$'$}$$

This leads to a contradiction again.

Lemma 4.5. *Assume that* (f0), (f1), (f2), (g0), (g1), (h0), (h1) *and* (h2)$'$ *hold. Then for* $\lambda \in (0, \lambda^*)$ *and* $\alpha \in (0, \alpha_1)$,
(i) $I_\lambda(u) > 0$ *for any* $u \in \Lambda_\lambda^-$,
(ii) *any minimizing sequence of* I_λ *on* Λ_λ^- *is uniformly bounded.*

P r o o f. We observe that, for $u \in \Lambda_\lambda^-$, from (2.4) and Lemma 4.3,

$$\begin{aligned} I_\lambda(u) &= \frac{1}{N} J_\lambda(u) - \alpha \frac{p^* - q}{p^* q} \int h|u|^q \\ &> \Big[\frac{1}{N} - \frac{p^* - p}{qp^*}\Big] J_\lambda(u) = \frac{q-p}{Nq} J_\lambda(u) \geqslant \frac{q-p}{Nq} \sigma \|u\|^p. \end{aligned} \tag{4.3}$$

Since σ only depends on α_1 and λ^*, the conclusions then follow directly. This completes the proof. □

Now, by Lemma 4.5, there exists $R > 0$, so that for any $\alpha \in (0, \alpha_1)$ and $\lambda \in (0, \lambda^*)$, for any minimizing sequence $\{u_n\} \subset \Lambda_\lambda^-$ of I_λ (here I_λ also depends on α), we can assume that, by taking a subsequence if necessary, $\|u_n\| \leqslant R$. Define $U_R = \{u \in V\colon \|u\| \leqslant R\}$.

Lemma 4.6. *Assume* (f0), (f1), (f2), (g0), (g1), (h0), (h1) *and* (h2)$'$ *hold. For any* $\overline{\lambda} \in (0, \lambda_1^+)$, *there exist* $\varrho > 0$ *and* α^* *with* $\alpha^* \leqslant \alpha_1$, *such that for any* $\lambda \in [\overline{\lambda}, \lambda^*)$, $\alpha \in (0, \alpha^*)$ *and* $u \in \Lambda_\lambda^- \cap U_{2R}$, $-\langle \Psi'_\lambda(u), u\rangle \geqslant \varrho$.

P r o o f. We first show that for some $\eta > 0$, depending only on α_1 and λ^*, $\|u\| \geqslant \eta$ for $u \in \Lambda_\lambda^-$. Indeed, if for some $u_n \in \Lambda_\lambda^-$, $u_n \to 0$, then we have, by

Lemma 4.3,

$$0 < \sigma \leqslant J_\lambda(v_n) = \alpha \int h|v_n|^q \cdot \|u_n\|^{q-p} + \int f|v_n|^{p^*} \cdot \|u_n\|^{p^*-p} \to 0,$$

a contradiction, where $v_n = u_n/\|u_n\|$.

Using Lemma 4.3 we get

$$\begin{aligned} -\langle \Psi'_\lambda(u), u\rangle &= (p^* - p)J_\lambda(u) - \alpha(p^* - q)\int h|u|^q \\ &\geqslant (p^* - p)J_\lambda(u) - \alpha(p^* - q)\|h\|_Q \cdot \|u\|^q \\ &\geqslant (p^* - p)\sigma\eta^p - \alpha(p^* - q)\|h\|_Q(2R)^q > c > 0, \end{aligned}$$

for α small enough. The lemma is proved. □

Lemma 4.6 implies that $\Lambda_\lambda^- \cap U_{2R}$ is a closed set (in fact one can prove that Λ_λ^- is a closed set). Replacing Λ_λ^- by $\Lambda_\lambda^- \cap U_{2R}$, and noting that any minimizing sequence in $\Lambda_\lambda^- \cap U_{2R}$ will be a positive distance from the boundary $\|u\| = 2R$, we can check straightforwardly that the proofs of Lemmas 3.2, 3.4, 3.5, and 3.6 remain valid. So we can state our result.

Theorem 4.7. *Assume that* (f0), (f1), (f2), (f3) *or* (f3)′, (g0), (g1), (h0), (h1), *and* (h2)′ *hold. Then there exist* $\lambda^* > \lambda_1^+$ *and* $\alpha^* > 0$, *such that for any* $\lambda \in [\lambda_1^+, \lambda^*)$ *and* $\alpha \in (0, \alpha^*)$, *the problem*

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u + \alpha h(x)|u|^{q-2}u + f(x)|u|^{p^*-2}u$$

has at least one positive solution in V.

R e m a r k 4.8. As we remarked earlier, for $\lambda \in (0, \lambda_1^+)$, Theorems 3.8 and 4.7 hold without the integral conditions (f2), (h2) and (h2)′, and can be proved via Mountain Pass argument. Cf. [1], [2], [7] and [12].

R e m a r k 4.9. We note that the proofs are applicable to Dirichlet problems on bounded domains and similar results hold. We can also deal with

$$-\Delta_p u + a(x)|u|^{p-2}u = \lambda g(x)|u|^{p-2}u + \alpha h(x)|u|^{q-2}u + f(x)|u|^{p^*-2}u$$

in $\mathbb{R}^N$, where $a(x) \in L^\infty_{\mathrm{loc}}(\mathbb{R}^N)$, $a(x) \geqslant 0$.

R e m a r k 4.10. Similarly, one can consider the negative principal eigenvalue $\lambda_1^- < 0$ given by

$$-\Delta_p u = \lambda g(x)|u|^{p-2}u \quad \text{in } \mathbb{R}^N, \qquad \int g(x)|u|^p < 0.$$

Existence of positive solutions of $(1.1)_\lambda$ for $\lambda < 0$ can be obtained provided conditions similar to (f2), (h2) and (h2)′ hold.

References

[1] *C. O. Alves*: Multiple positive solutions for equations involving critical Sobolev exponent in $\mathbb{R}^N$. Electron. J. Differential Equations *13* (1997), 1–10.

[2] *C. O. Alves, J. V. Gonçalves, O. H. Miyagaki*: Remarks on multiplicity of positive solutions for nonlinear elliptic equations in $\mathbb{R}^N$ with critical growth. Preprint.

[3] *H. Brezis, L. Nirenberg*: Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents. Comm. Pure Appl. Math. *36* (1983), 437–477.

[4] *P. Drábek, Y. X. Huang*: Multiple positive solutions of quasilinear elliptic equations in $\mathbb{R}^N$. Nonlinear Anal. To appear.

[5] *P. Drábek, Y. X. Huang*: Multiplicity of positive solutions for some quasilinear elliptic equation in $\mathbb{R}^N$ with critical Sobolev exponent. J. Differential Equations *140* (1997), 106–132.

[6] *P. Drábek, Y. X. Huang*: Bifurcation problems for the p-Laplacian in $\mathbb{R}^N$. Trans. Amer. Math. Soc. *349* (1997), 171–188.

[7] *J. V. Gonçalves, C. O. Alves*: Existence of positive solutions for m-Laplacian equations in $\mathbb{R}^N$ involving critical Sobolev exponents. Nonlinear Anal. *32* (1998), 53–70.

[8] *P. L. Lions*: The concentration-compactness principle in the calculus of variations, the limit case, Part I, II. Rev. Mat. Iberoamericana *1* (1985), no. 2, 3, 109–145, 45–121.

[9] *J. Mawhin, M. Willem*: Critical Point Theory and Hamiltonian Systems. Appl. Math. Sci. Vol. 74, Springer-Verlag, New York, 1989.

[10] *E. S. Noussair, C. A. Swanson*: Multiple finite energy solutions of critical semilinear field equations. J. Math. Anal. Appl. *195* (1995), 278–293.

[11] *E. S. Noussair, C. A. Swanson, J. Yang*: Quasilinear elliptic problems with critical exponents. Nonlinear Anal. *20* (1993), 285–301.

[12] *C. A. Swanson, L. S. Yu*: Critical p-Laplacian problems in $\mathbb{R}^N$. Ann. Mat. Pura Appl. *169* (1995), 233–250.

[13] *G. Tarantello*: On nonhomogeneous elliptic equations involving critical Sobolev exponent. Ann. Inst. H. Poincaré Anal. Non Linéaire *9* (1992), 281–304.

[14] *J. Yang*: Positive solutions of quasilinear elliptic obstacle problems with critical exponents. Nonlinear Anal. *25* (1995), 1283–1306.

Authors' addresses: *Paul A. Binding*, Department of Mathematics & Statistics, University of Calgary, Calgary, Alberta, Canada, T2N 1N4, e-mail: `binding@acs.ucalgary.ca`; *Pavel Drábek*, Department of Mathematics, University of West Bohemia, P.O. Box 314, 306 14 Plzeň, Czech Republic, e-mail: `pdrabek@kma.zcu.cz`; *Yin Xi Huang*, Department of Mathematical Sciences, University of Memphis, Memphis, TN 38152, U.S.A., e-mail: `huangy@mathsci.msci.memphis.edu`.