МАТЕМАТИЧКИ ВЕСНИК
45 (1993), 29–33

UDC 517.982.2
оригинални научни рад
research paper

AN INEQUALITY RELATED TO THE UNIFORM CONVEXITY IN BANACH SPACES

Miroslav Pavlović

Abstract. We prove an inequality that implies that a 2-concave and p-convex Banach lattice is "more" uniformly convex than L^p.

1. Introduction

In this note we prove the following

THEOREM. *Let X be a 2-concave Banach lattice with 2-concavity constant equal to one, and let $1 \leqslant p \leqslant 2$. Then*

$$\|(|x+y|^p + |x-y|^p)^{1/p}\| \geqslant \left\{(\|x\| + \|y\|)^p + \big|\|x\| - \|y\|\big|^p\right\}^{1/p}, \tag{1}$$

for all $x, y \in X$. In particular, inequality (1) *holds in an arbitrary L^q space with $1 \leqslant q \leqslant 2$.*

For the definition of the expression $(|u|^p + |v|^p)^{1/p}$ and other notions concerning abstract Banach lattices we refer to [3], Ch. 1 (especially Theorem 1.d.1). In the case where $X = L^p$ $(1 < p < 2)$ inequality (1) becomes

$$\|x+y\|^p + \|x-y\|^p \geqslant (\|x\| + \|y\|)^p + \big|\|x\| - \|y\|\big|^p, \tag{2}$$

which was used by Hanner [2] to calculate the precise value of the modulus of convexity of L^p. Moreover, it follows from [4] that the validity of (2) in some normed spaces X implies that X is "more" uniformly convex than L^p (where L^p is at least two-dimensional). An immediate consequence of Theorem is that (1) holds in a large class (denoted by $\Delta(p,2)$; see Section 2) containing, for example, L^q for $p \leqslant q \leqslant 2$ as well as certain Orlicz and mixed normed Lebesgue spaces. Note that, in [4], the validity of (2) in L^q $(p \leqslant q \leqslant 2)$ was deduced from the case $q = p$ by using the fact that L^q can be embedded into $L^p(0,1)$ isometrically (see [3], pp. 181–182). The proof in the present note is quite elementary and lies on the fact that (for $1 \leqslant p \leqslant 2$) the function

$$F_p(\xi, \eta) := \left\{(\xi^{1/2} + \eta^{1/2})^p + |\xi^{1/2} - \eta^{1/2}|^p\right\}^{2/p} \qquad (\xi \geqslant 0,\ \eta \geqslant 0) \tag{3}$$

is convex. Before proving the result we mention a generalization of F_p that could be of some independent interest. Let r_j $(j = 0, 1, 2, \dots)$ denote the Rademacher functions,

$$r_j(t) = \operatorname{sign}(\sin(2^j \pi t)) \qquad (t \text{ real}).$$

Define the functions Φ_p on the positive cone l^1_+ of the sequence space l^1 by

$$\Phi_p(\xi) = \left\{ \int_0^1 \left| \sum_{j=0}^{\infty} r_j(t) \xi_j^{1/2} \right|^p dt \right\}^{2/p} \qquad (\xi = (\xi_j)_0^\infty \geqslant 0).$$

That the definition is correct follows from the well known fact that if $(a_j)_0^\infty \in l^2$, then the series $\sum a_j r_j(t)$ converges almost everywhere, and from Khintchine's inequality [3], Theorem 2.b.3, which says that

$$A_p \|\xi\|_{l^1} \leqslant \Phi_p(\xi) \leqslant B_p \|\xi\|_{l^1} \qquad (A_p, B_p = \text{const} > 0).$$

Starting from the observation that $\Phi(\xi_1, \xi_2, 0, 0, \dots) = \text{const } F_p(\xi_1, \xi_2)$ we conjecture that Φ_p is a convex function on l^1_+ (for $1 \leqslant p \leqslant 2$). (We shall also prove that if $p > 2$, then F_p is concave, and we conjecture that Φ_p is concave if $p > 2$).

This would lead to the inequality

$$\|\Phi_p(x_1, x_2, \dots)\| \geqslant \Phi_p(\|x_1\|, \|x_2\|, \dots),$$

where x_1, x_2, … are elements of a Banach lattice whose 2-concavity constant is eual to one. Further remarks are at the end of the paper.

2. Definitions and examples

We denote by $\Delta(p, q)$, where $1 \leqslant p \leqslant q \leqslant +\infty$, the class of (real) Banach lattices X such that

$$\|(|u|^p + |v|^p)^{1/p}\| \leqslant (\|u\|^p + \|v\|^p)^{1/p} \tag{4}$$

and

$$\|(|u|^q + |v|^q)^{1/q}\| \geqslant (\|u\|^q + \|v\|^q)^{1/q} \tag{5}$$

for all $u, v \in X$. In other words, X is in $\Delta(p, q)$ if it is p-convex and q-concave and its p-convexity and q-concavity constants are equal to one. It is clear that $\Delta(1, \infty)$ is just the class of all Banach lattices. And by [3], Proposition 1.d.5, $\Delta(p, q)$ is contained in $\Delta(r, s)$ for $r \leqslant p \leqslant q \leqslant s$. In particular, $L^q \in \Delta(r, s)$ if $r \leqslant q \leqslant s$, a fact which can easily be verified by direct calculations.

It was proved by Figiel [1] (see also [3], pp. 80–81) that if $X \in \Delta(p, q)$ with $p > 1$ and $q < +\infty$, then X is uniformly convex in the sense that

$$\delta_X(\varepsilon) = \inf \left\{ 1 - \left\| \frac{x + y}{2} \right\| : \|x - y\| = \varepsilon,\ \|x\| = \|y\| = 1 \right\} > 0$$

for $\varepsilon > 0$. The function δ_X is called the modulus of convexity of X. Let δ_p denote the modulus of convexity of L^p, $\dim(L^p) \geqslant 2$. (It follows from [2] that δ_p is independent of a particular choice of L^p.) As noted in Introduction, the following fact follows immediately from (1) and (4).

COROLLARY 1. *If* $X \in \Delta(p,2)$ (*in particular,* $X = L^q$ *for* $2 \geqslant q \geqslant p$), *then inequality* (2) *holds.*

As noted in Introduction, this implies the following

COROLLARY 2. *If* $X \in \Delta(p,2)$, *then* $\delta_X(\varepsilon) \geqslant \delta_p(\varepsilon)$ $(\varepsilon > 0)$.

Mixed normed spaces. For technical reasons we define only sequence spaces. Let $1 \leqslant r, s \leqslant 2$. The space $X = l^{r,s}$ consists of those scalar sequences $x = \{x_{j,k}\}_{j,k=0}^{\infty}$ such that

$$\|x\| = \left\{ \sum_{j=0}^{\infty} \left[\sum_{k=0}^{\infty} |x_{j,k}|^s \right]^{r/s} \right\}^{1/r} < \infty.$$

It is not hard to show that $l^{r,s} \in \Delta(p,q)$, where $p = \min(r,s)$ and $q = \max(r,s)$. Hence, by Corollary 2, $\delta_X(\varepsilon) \geqslant \delta_p(\varepsilon)$. Since $l^{r,s}$ contains an isometric copy of l^p, we conclude that $\delta_X = \delta_p$.

Orlicz spaces. Let M be a convex, strictly increasing function on the interval $[0,\infty)$ with $M(0) = 0$. The space l^M consists of the scalar sequnces $x = \{x_j\}_0^{\infty}$ for which

$$\|x\| = \|x\|_M = \inf \left\{ \lambda > 0 : \sum_{j=0}^{\infty} M\left(\frac{|x_j|}{\lambda}\right) \leqslant 1 \right\} < \infty.$$

One can prove that $l^M \in \Delta(p,q)$ provided that the function $M(t^{1/p})$ is convex and the function $M(t^{1/q})$ is concave. Therefore, inequality (1) holds in l^M if the function $M(t^{1/q})$ is concave. Estimates for the moduli of convexity of Orlicz spaces can be found in [1].

3. Proofs

Our proof is based on the following lemma.

LEMMA. *Let* F_p *be defined by* (3). *Then, if* $1 \leqslant p \leqslant 2$, *the function* F_p *is convex, and if* $p > 2$, *it is concave. In all the cases* $F_p(\xi,\eta)$ *increases with* ξ *and* η.

Before proving the lemma we use it to prove the theorem. Let $x, y \in X$, where $X \in \Delta(1,2)$, and $1 \leqslant p \leqslant 2$. Then

$$(|x+y|^p + |x-y|^p)^{1/p} = \left((|x|+|y|)^p + \big||x|-|y|\big|^p\right)^{1/p}$$

(this is deduced from the case where x, y are real scalars, by using Theorem 1.d.1 of [3]) and we may assume that $x \geqslant 0$, $y \geqslant 0$. Assuming this we have

$$(|x+y|^p + |x-y|^p)^{1/p} = F_p(x^2, y^2)^{1/2}$$

(see [3], Theorem 1.d.1). Since F_p is convex, homogeneous and "increasing", there is a set $A \subset \{\,(\alpha,\beta) : \alpha \geqslant 0, \beta \geqslant 0\,\}$ such that

$$F_p(\xi,\eta) = \sup\{\,\alpha\xi + \beta\eta : (\alpha,\beta) \in A\,\},$$

whence $F_p(x^2,y^2)^{1/2} \geqslant (\alpha x^2 + \beta y^2)^{1/2}$, $(\alpha,\beta) \in A$, and hence, by (5) with $q = 2$,

$$\|F_p(x^2,y^2)^{1/2}\| \geqslant (\alpha\|x\|^2 + \beta\|y\|^2)^{1/2}$$

for all $(\alpha,\beta) \in A$. Taking the supremum over $(\alpha,\beta) \in A$ we obtain

$$\|F_p(x^2,y^2)^{1/2}\| \geqslant F_p(\|x\|^2,\|y\|^2)^{1/2},$$

which concludes the proof. ∎

Proof of Lemma. Let $1 < p \leqslant 2$. (The case $p = 1$ is similar.) Since $F_p(\lambda\xi,\lambda\eta) = \lambda F_p(\xi,\eta)$ for $\lambda \geqslant 0$, the convexity of F_p will follow from the convexity of the function $f(t) = F_p(1,t)$, $t > 0$. To prove that f is convex observe first that $f(t) = tf(1/t)$, whence $f''(t) = t^{-3}f''(1/t)$ for $t \neq 1$. And since $f'(1)$ exists, it remains to prove that $f''(t) \geqslant 0$ for $0 < t < 1$. To prove this write f as

$$f(t) = g(t^{1/2})^{2/p}, \quad g(t) = (1+t)^p + (1-t)^p \qquad (0 < t < 1).$$

We have

$$\begin{aligned} 2pf''(t) = t^{-2/3}g(t^{1/2})^{(2/p)-2}\Big[\Big(\frac{2}{p}-1\Big)\, & g'(t^{1/2})^2t^{1/2} \\ & + g(t^{1/2})g''(t^{1/2})t^{1/2} - g(t^{1/2})g'(t^{1/2})\Big]. \end{aligned}$$

Hence, $f''(t) > 0$ if and only if $A(t) > 0$, where

$$\begin{aligned} A(t) &= \frac{1}{p}\Big[\Big(\frac{2}{p}-1\Big)\, g'(t)^2t + g(t)g''(t)t - g(t)g'(t)\Big] \\ &= 4(p-1)t(1-t^2)^{p-2} - [(1+t)^{2p-2} - (1-t)^{2p-2}]. \end{aligned}$$

If $3/2 \leqslant p \leqslant 2$, the function $\varphi(t) = (1+t)^{2p-2} - (1-t)^{2p-2}$ is concave and therefore

$$\varphi(t) \leqslant \varphi(0) + \varphi'(0)t = 4(p-1)t \leqslant 4(p-1)t(1-t^2)^{p-2},$$

which implies $A(t) > 0$. If $1 < p \leqslant 3/2$, then

$$A'(t) = 4(p-1)(1-t^2)^{p-3}[1 + (3-2p)t^2] - 2(p-1)[(1+t)^{2p-3} + (1-t)^{2p-3}].$$

Since $0 \leqslant 3-2p \leqslant 1$, the function $t \mapsto t^{3-2p}$ is concave, hence

$$\frac{(1+t)^{2p-3} + (1-t)^{2p-3}}{2} = \frac{1}{2}\left[\left(\frac{1}{1+t}\right)^{3-2p} + \left(\frac{1}{1-t}\right)^{3-2p}\right] \leqslant (1-t^2)^{2p-3}.$$

Hence

$$A'(t) \geqslant 4(p-1)(1-t^2)^{2p-3}[1 + (3-2p)t^2 - 1] \geqslant 0.$$

This implies $A(t) \geqslant A(0) = 0$, which concludes the proof in the case $1 < p \leqslant 2$. If $p > 2$, proving that F_p is concave reduces to proving that $A(t) \leqslant 0$ $(0 < t < 1)$. In this case the function φ is convex which implies that

$$\varphi(t) \geqslant \varphi(0) + \varphi'(0)t = 4(p-1)t \geqslant 4(p-1)t(1-t^2)^{p-2},$$

and this completes the proof. ■

Remark. The discussion of the case $1 < p \leqslant 2$ can be made simplier. Namely, it is easy to see that the function $g(t^{1/2})$ is convex $(0 < t < 1)$, which implies that $f(t) = g(t^{1/2})^{2/p}$ is convex (since $2/p > 1$). This trick can also be used if $2 < p < 3$, because then the function $g(t^{1/2})$ is concave. However, if $p > 3$, $g(t^{1/2})$ is convex.

4. Dual results

Using the case $p > 2$ of Lemma one proves that if $x, y \in X$, where $X \in \Delta(2, \infty)$ (which means that X satisfies (4) with $p = 2$), then there holds the reverse of (1). A consequence is that the reverse of (2) is valid in every lattice of class $\Delta(2, p)$ $(p > 2)$ and, in particular, in L^q for $2 \leqslant q \leqslant p$. (The latter was proved in [4] by using the Riesz-Thorin interpolation theorem.) Combining this with Hanner's results we see that if $X \in \Delta(2, p)$, then X is "more" uniformly convex than L^p (dimension $\geqslant 2$) in the sense that $\rho_X(\tau) \leqslant \rho_p(\tau)$, where

$$\rho_X(\tau) = \sup\left\{ \frac{\|x+\tau y\| + \|x-\tau y\|}{2} - 1 : \|x\| = 1,\ \|y\| = 1 \right\},$$

and $\rho_p = \rho_{L^p}$. The function ρ_X is called the modulus of smoothness of X (see [3], Ch. 1, for further information).

REFERENCES

[**1**] T. Figiel, *On the moduli of convexity and smoothness*, Studia math. **56** (1976), 121–155.

[**2**] O. Hanner, *On the uniform convexity of L^p and l^p*, Arkiv f. Math **3**:3 (1956), 239–244.

[**3**] J. Lindenstaruss and L. Tzafriri, *Classical Banach Spaces II. Function Spaces*, Ergebnisse **97**, Berlin-Heidelberg-New York, Springer-Verlag 1979.

[**4**] M. Pavlović, *Some inequalities in L^p spaces II*, Mat. vesnik **10(23)(38)** (1986), 321–326.

(received 05.03.1993)

Matematički fakultet, Studentski trg 16, 11000 Beograd, YUGOSLAVIA