PUBLICATIONS DE L'INSTITUT MATHÉMATIQUE
Nouvelle série, tome 33 (47), 1983, pp. 143–155

EXTRAPOLATION OF MOVING AVERAGE AND AUTOREGRESSIVE PROCESSES WHEN THE ENTIRE PAST OF THE PROCESSES IS KNOWN

Pavle Mladenović

1. Introduction. Let $X(t) = (X_1(t), \ldots, X_n(t))$, $t \in R$, be a multidimensional and wide sense stationary random process with the mean value zero, the correlation matrix $\|B^X_{jk}(\tau)\|$ and the spectral process $Z^X(\lambda) = (Z^X_1(\lambda), \ldots, Z^X_n(\lambda))$ $\lambda \in R$. Suppose that all elements $B_{jk}(\tau)$ of the correlation matrix $\|B_{jk}(\tau)\|$ fall off sufficiently rapidly at infinity so that they can be represented in the form:

$$B^X_{jk}(\lambda) = \int_{-\infty}^{+\infty} e^{i\tau\lambda} f^X_{jk}(\lambda) d\lambda, \quad i,k = 1,2,\ldots,n \tag{1.1}$$

(i.e. there exist the spectral densities $f^X_{jk}(\lambda)$).

Let H denote the Hilbert space of all second-order random functions, where a scalar product is defined by $(\xi, \eta) = E\xi\overline{\eta}$, and let $H(X)$ be the Hilbert space generated by $\{X_k(t) : t \in R, k = 1,2,\ldots,n\}$. Let $L^2(F)$ be the space of functions $\Phi(\lambda) = (\Phi_1(\lambda), \ldots, \Phi_n(\lambda))$ satisfying the following condition:

$$\sum_{k=1}^{n}\sum_{j=1}^{n} \int_{-\infty}^{+\infty} \Phi_k(\lambda)\overline{\Phi_j(\lambda)} f^X_{j,k}(\lambda) d\lambda < \infty \tag{1.2}$$

The space $L^2(F)$ becomes a Hilbert space if we define the scalar product $(\Phi(\lambda), \Psi(\lambda))$ by:

$$(\Phi(\lambda), \Psi(\lambda)) = \int_{-\infty}^{+\infty} \sum_{k=1}^{n}\sum_{j=1}^{n} \Phi_k(\lambda)\overline{\Psi_j(\lambda)} f^X_{j,k}(\lambda) d\lambda < \infty \tag{1.3}$$

AMS Subject Classification (1980): 60G10, 60G25

A random function ξ belongs to $H(X)$ if and only if there exists $\Phi(\lambda) \in L^2(F)$ so that

$$\xi = \sum_{k=1}^{n} \int_{-\infty}^{+\infty} \Phi_k(\lambda) dZ_k(\lambda) \tag{1.4}$$

The mapping of $H(X)$ to $L^2(F)$ given by $\xi = \sum_{k=1}^{n} \int \Phi_k(\lambda) dZ_k(\lambda) \to (\Phi_1(\lambda), \ldots, \Phi_n(\lambda))$ is an isometry.

A stationary random process $X(t)$, $t \in R$, is said to be a process with a rational spectrum if all the functions $f_{jk}^X(\lambda)$ are rational functions, i.e. $f_{jk}(\lambda) = Q_{jk}(\lambda)/P_{jk}(\lambda)$, where $P_{jk}(\lambda)$ and $Q_{jk}(\lambda)$ are polynomials in λ. The spectral densities $f_{jj}(\lambda)$ can be represented in the form:

$$B_0|\lambda^M + B_1\lambda^{M-1} + \cdots + B_N|^2 |\lambda^P + A_1\lambda^{P-1} + \cdots + A_P|^{-2} \tag{1.5}$$

where $B_0 > 0$ and $M < P$. Denote the determinant of the spectral matrix $\|f_{jk}(\lambda)\|$ by $D(\lambda)\,(= Q(\lambda)/P(\lambda))$. Let $2N_{jj}$, $2(N_{jj} - m_j)$, $2K$, $2L$ be the degrees of $P_{jj}(\lambda), Q_{jj}(\lambda), P(\lambda), Q(\lambda)$ respectively. Then $2K - 2L \geq 2\sum_{k=1}^{n} m_k$.

A stationary random process with a rational spectrum is nonsingular if the determinant $D(\lambda)$ has no real zeros and $2K - 2L \geq 2\sum_{k=1}^{n} m_k$. The polynomial $Q(\lambda)$ can be represented in the form $B(\lambda - \theta_1)\ldots(\lambda - \theta_L)(\lambda - \overline{\theta}_1)\ldots(\lambda - \overline{\theta}_L)$ where $\Im\theta_i > 0$, $i = 1, 2, \ldots, L$.

Now, suppose we know the values $X_k(s)$, $s \leq t$, $k = 1, 2, \ldots, n$. The simplest problem of linear extrapolation of a stationary random process $X(t)$ is the following: Find a random variable

$$\tilde{X}_1(t, T) = \sum_{k=1}^{n} \int_{-\infty}^{+\infty} \Phi_k(\lambda) dZ_k(\lambda) \in H(X)$$

which is the best approximation of $X_1(t + \tau)$, depending linearly on $X_k(s)$, $s \leq t$, $k = 1, 2, \ldots, n$. As an index of approximation quality we shall use the mean square error. We shall call $\tilde{X}_1(t, \tau)$ the linear least-squares estimator of $X_1(t + \tau)$. We see that $X_1(t, \tau)$ is the projection of $X_1(t + \tau)$ into the smallest Hilbert space $H(X, t)$ spaned by $\{X_k(s), s \leq t, k = 1, 2, \ldots, n\}$.

In the paper [**7**] the following theorem was proved:

THEOREM 1. (Yaglom): *If $X(t) = (X_1(t), \ldots, X_n(t))$ is a nonsingular stationary random process with a rational spectrum, then the linear least-squares estimator of $X_1(t + \tau)$ is given by:*

$$\tilde{X}_1(t, \tau) = \sum_{k=1}^{n} \left\{ \sum_{j=0}^{m_k - 1} wk^{(j)}(t) + \int_0^{\infty} w_k(s, \tau) X_k(t - s) ds \right\} \tag{1.6}$$

where the coefficients $w_k^{(j)}(\tau)$ *and the functions* $w_k(s,\tau)$ *can be found (in a unigue way) from the condition: the functions*

$$\Phi_k(\lambda) = \sum_{j=0}^{m_k-1} w_k^{(j)}(\tau)(i\lambda)^j + \int_0^\infty e^{-is\lambda} w_k(s,\tau)ds \tag{1.7}$$

have the form

$$\Phi_k(\lambda) = \omega_k(\lambda)\{(\lambda-\theta_1)\dots(\lambda-\theta_l)\}^{-1}, \quad k=1,2,\dots,n. \tag{1.8}$$

where $w_k(\lambda)$ *is a polynomial of degree* $L+m_k-1$ *(but not of a greater degree) such that the functions*

$$\Psi_k(\lambda) = (e^{i\tau\lambda} - \Phi_1(\lambda))f_{1k}(\lambda) - \sum_{j=2}^{n} \Phi_j(\lambda) f_{jk}(\lambda) \tag{1.9}$$

are analytic in the upper half plane.

The functions $\Phi_k(\lambda)$, $k=1,2,\dots,n$, will be called the spectral characteristic for extrapolation of the random process $X(t)$ at the point $t+\tau$.

2. Moving average and autoregressive processes

Let $X(t) = (X_1(t),\dots,X_n(t))$ and $Y(t) = (Y_1(t)),\dots,Y_n(t))$ be two multidimensional stationary random processes satisfying the following equation

$$Y(t) = \sum_{\nu=0}^{N} a_\nu X(t-\nu\theta), \quad a_0 = 1, \quad \theta > 0 \tag{2.1}$$

The process $Y(t)$ is called the moving average process of order N associated with the process $X(t)$. We shall denote it by $MA_X(N,\theta)$. The process $X(t)$ is called the autoregressive process of order N associated with the process $Y(t)$. We shall denote it by $AR_Y(N,\theta)$.

It the roots of

$$t^N + a_1 t^N + \dots + a_N = 0 \tag{2.2}$$

are less than one in absolute value, then the following equations also hold:

$$X_k(t) = \sum_{\nu=0}^{\infty} c_\nu Y_k(t-\nu\theta), \quad k=1,2,\dots,n. \tag{2.3}$$

The series on the right side of (2.3) converges in quadratic mean and the coefficients c_ν satisfy the homogeneous difference equation

$$a_0 c_k + a_1 c_{k-1} + \dots + a_N c_{k-N} = 0, \quad k > N \tag{2.4}$$

and the initial conditions

$$\begin{aligned} &a_0 = 1 \\ &a_0 c_k + a_1 c_{k-1} + \dots + a_k c_0 = 0, \quad O < k < N. \end{aligned} \tag{2.5}$$

THEOREM 2. *Let $X(t)$ and $Y(t)$ be a multidimensional stationary random processes with spectral processes $(Z_1^X(\lambda),\dots,Z_n^X(\lambda))$ and $(Z_1^Y(\lambda),\dots,Z_n^Y(\lambda))$ and let $X(t)$ and $Y(t)$ satisfy (2.1). Then:*

$$\text{a)}\quad dZ_k^Y(\lambda)=\left\{\sum_{\nu=0}^{N}a_\nu e^{-i\lambda\nu\theta}\right\}dZ_k^X(\lambda),\quad k=1,2,\dots,n \tag{2.6}$$

$$\text{b)}\quad f_{jk}(\lambda)=\left|\sum_{\nu=0}^{N}a_\nu e^{-i\nu\theta\lambda}\right|^2 f_{jk}(\lambda),\quad j,k=1,2,\dots,n \tag{2.7}$$

c) *If the roots of* (2.2) *are less than one in absolute value then*

$$H(X,t)=H(Y,t). \tag{2.8}$$

Proof: The equation (2.6) follows from

$$\int_{-\infty}^{+\infty}e^{it\lambda}dZ_k^Y(\lambda)=Y(t)=\sum_{\nu=0}^{N}a_\nu\int_{-\infty}^{+\infty}e^{i(t-\nu\theta)\lambda}dZ_k^X(\lambda)=\int_{-\infty}^{+\infty}e^{it\lambda}\sum_{\nu=0}^{N}a_\nu e^{-i\nu\theta\lambda}dZ_k^X(\lambda).$$

The equation (2.7) follows from

$$\int_{-\infty}^{+\infty}e^{i\tau\lambda}f_{jk}^Y(\lambda)d\lambda=B_{jk}^Y(\tau)=EY_j(t+\tau)\overline{Y_k(t)}=$$

$$=E\sum_{\nu=0}^{N}a_\nu X_j(t+\tau-\nu\theta)\sum_{\mu=0}^{N}\overline{a}_\mu\overline{X_k(t-\mu\theta)}=\sum_{\nu=0}^{N}\sum_{\mu=0}^{N}a_\nu\overline{a}_\mu B_{jk}^X(\tau-(\nu-\mu)\theta)$$

$$=\sum_{\nu=0}^{N}\sum_{\mu=0}^{N}a_\nu\overline{a}_\mu\int_{-\infty}^{+\infty}e^{i\tau\lambda}e^{i(\nu-\mu)\theta\lambda}f_{jk}^X(\lambda)=\int_{-\infty}^{+\infty}e^{i\tau\lambda}\left|\sum_{nu=0}^{N}a_\nu e^{-i\nu\theta\lambda}\right|^2 f_{jk}^X(\lambda)d\lambda.$$

The result (2.8) follows from (2.1) and (2.3).

In this paper, we shall solve the simplest extrapolation problem for the moving average and autoregressive processes associated with a nonsingular stationary process with a rational spectrum.

3. Linear extrapolation of moving average processes

THEOREM 3. *Let $X(s)$ and $Y(s)$ be two multidimensional stationary processes satisfying* (2.1) *and let the roots of the equation* (2.2) *be less than one in absolute value. If $\Phi_{k,\tau}^X(\lambda)$ and $\Phi_{k,\tau}^Y(\lambda)$ are the spectral characteristics for extrapolation of the processes $X(s)$ and $Y(s)$ at the point $t+\tau$ $(\tau>0)$, then:*

a) *If τ/θ is not an integer and $[\tau/\theta] = S < N$, then:*

$$\Phi^Y_{1,\tau}(\lambda) = \left\{ \sum_{\nu=0}^{S} a_\nu \Phi^X_{1,\tau-\nu\theta}(\lambda) + \sum_{\nu=S+1}^{N} a_\nu e^{i(\tau-\nu\theta)\lambda} \right\} \left\{ \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} \right\}^{-1} \tag{3.1}$$

$$\Phi^Y_{k,\tau}(\lambda) = \sum_{\nu=0}^{S} a_\nu \Phi^X_{1,\tau-\nu\theta}(\lambda) \left\{ \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} \right\}^{-1}, k = 2, \ldots, n. \tag{3.2}$$

b) *If $\tau = S\theta$ and $S \leq N$, it is only necessary to replace S by $S-1$ in the equations* (3.1) *and* (3.2).

c) *If τ/θ is not an integer and $S \geq N$ or if $\tau = S\theta$ and $S > N$, then:*

$$\Phi^Y_{k,\tau}(\lambda) = \sum_{\nu=0}^{N} a_\nu \Phi^X_{1,\tau-\nu\theta}(\lambda) \left\{ \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} \right\}^{-1}, k = 2, \ldots, n. \tag{3.3}$$

Proof: a) If τ/θ is not an integer and $S < N$, we have $\tau - \nu\theta > 0$ for $\nu = 0, 1, 2, \ldots, S$ and $\tau - \nu\theta \leq 0$ for $\nu = S+1, \ldots, N$.

It follows from (2.8) that

$$\tilde{Y}_1(t,\tau) = \operatorname{Proj}_{H(Y,t)} Y_1(t+\tau) = \sum_{\nu=0}^{N} \operatorname{Proj}_{H(X,t)} X_1(t+\tau-\nu\theta) =$$

$$= \sum_{\nu=0}^{S} a_\nu \tilde{X}_1(t, \tau-\nu\theta) + \sum_{\nu=S+1}^{N} a_\nu X_1(t+\tau-\nu\theta).$$

If we apply (1.6) and (2.6), we obtain

$$\tilde{Y}_1(t,\tau) = \int_{-\infty}^{+\infty} e^{it\lambda} \left\{ \sum_{\nu=0}^{S} a_\nu \Phi^X_{k,\tau-\nu\theta}(\lambda) + \sum_{\nu=S+1}^{N} a_\nu e^{i(\tau-\nu\theta)\lambda} \right\} \left\{ \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} \right\}^{-1} dZ^X_1(\lambda) +$$

$$+ \sum_{k=2}^{n} \int_{-\infty}^{+\infty} e^{it\lambda} \sum_{\nu=0}^{S} a_\nu \Phi_{k,\tau-\nu\theta}(\lambda) \left\{ \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} \right\}^{-1} dZ^X_k(\lambda)$$

and the desired result follows.

b) and c) The proof is similar.

COROLLARY: *If $X(s), s \in R$, is a nonsingular stationary random process with rational spectrum, then the spectral characteristics $\Phi^Y_{k,\tau}(\lambda)$, $k = 1, 2, \ldots, n$, can be represented in the form:*

a) *If τ/θ is not an integer and $S < N$ then:*

$$\Phi^Y_{1,\tau}(\lambda) R_1(\lambda) \sum_{\nu=0}^{\infty} c^{(1)}_\nu e^{-i\nu\theta\lambda} + \sum_{\nu=S+1}^{\infty} c^{(2)}_\nu e^{i(\tau-\nu\theta)\lambda} \tag{3.4}$$

$$\Phi^Y_{k,\tau}(\lambda) R_k(\lambda) \sum_{\nu=0}^{\infty} c_\nu e^{-i\nu\theta\lambda}, \quad k = 2, \ldots, n. \tag{3.5}$$

b) *If $\tau = s\theta$ and $S \leq N$, it is necessary to replace S by $S-1$ in the formulae* (3.4) *and* (3.5).

c) *If τ/θ is not an integer and $S \geq N$ or if $\tau = S\theta$ and $S > N$ then:*

$$\Phi^{Y}_{k,\tau}(\lambda) = R_k(\lambda) \sum_{\nu=0}^{\infty} c_\nu e^{-i\nu\theta\lambda}, \quad k = 1, 2, \ldots, n. \tag{3.6}$$

The functions $R_k(\lambda) = \sum_{\nu} a_\nu \Phi_{k,\tau-\nu\theta}(\lambda)$ are rational functions.

Proof: If α_ν, $\nu = 1, 2, \ldots, n$, are the roots of the equation (2.2), we have

$$\left\{ \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} \right\}^{-1} = \prod_{nu=0}^{N} (1 - \alpha e^{-i\nu\theta\lambda})^{-1} = \sum_{\nu=0}^{\infty} c_\nu e^{-i\nu\theta\lambda}$$

and the proof follows.

THEOREM 4. *Let $X(s)$ and $Y(s)$ satisfy the assumptions of Theorem* 3, *and let $X(s)$ be a nonsingular stationary process with rational spectrum. Suppose we know the values $Y_k(s)$, $s \leq t$, $k = 1, 2, \ldots, n$. If τ/θ is not an integer and $S < N$, then the linear least-squares estimator of $Y_1(t+\tau)$ is given by:*

$$\begin{aligned} \tilde{Y}_1(t,\tau) = \sum_{k=1}^{n} \left\{ \sum_{j=-}^{m_k-1} \sum_{\alpha=0}^{\infty} A^{(j)}_{k,\alpha} Y_k(j)(t - \alpha\theta) + \int_0^{\infty} B_k(u) Y_k(t-u)\,du \right\} + \\ + \sum_{\mu=0}^{\infty} D_\mu Y_1(t + \tau - (S+1)\theta - \mu\theta), \end{aligned} \tag{3.7}$$

where

$$\begin{aligned} A^{(j)}_{k,\alpha} &= c_\alpha \sum_{\nu=0}^{S} a_\nu w^{(j)}_k(\tau - \nu\theta), \\ B_k(u) &= \sum_{\alpha=0}^{\infty} c_\alpha D_k(u - \alpha\theta, \tau), \end{aligned} \tag{3.8}$$

$$D_k(u - \alpha\theta, \tau) = \begin{cases} 0 & 0 \leq u < \alpha\theta \\ \sum_{\nu=0}^{S} a_\nu w_k(u - \alpha\theta, \tau - \nu\theta) & u \geq \alpha\theta \end{cases}, \quad \alpha = 0, 1, \ldots, \tag{3.10}$$

$$D_\mu = \sum_{\nu=S+1}^{N} a_\mu F(\mu, \nu) \tag{3.11}$$

$$F(\mu,\nu) = \begin{cases} 0 & 0 \leq \mu < \nu - S - 1 \\ c_{\mu-\nu+S+1} & \mu \geq \nu - S - 1 \end{cases}, \quad \nu = S+1, \ldots, N \tag{3.12}$$

where the coefficients c_ν were given by (2.4) *and* (2.5) *and $w^{(j)}_k$, $w_k(s,\tau)$ were determined by Theorem* 1.

Proof: Using the formulae (1.6) and (2.3) we obtain

$$\tilde{Y}_1(t,\tau) = \sum_{\nu=0}^{N} a_\nu \tilde{X}_1(t,\tau-\nu\theta) + \sum_{\nu=S+1}^{N} a_\nu X(t+\tau-\nu\theta) =$$

$$= \sum_{\nu=0}^{N} a_\nu \left\{ \sum_{k=1}^{n} \left[\sum_{j=0}^{m_k-1} w_k^{(j)}(\tau-\nu\theta) X_k^{(j)}(t) + \int_0^\infty w_k(s,\tau-\nu\theta) X_k(t-s)ds \right] \right\} +$$

$$+ \sum_{\nu=S+1}^{N} a_\nu X_1(t+\tau-\nu\theta) = \sum_{k=1}^{n}\sum_{\alpha=0}^{\infty} c_\alpha \left\{ \sum_{j=0}^{m_k-1}\sum_{\nu=0}^{S} a_\nu w_k^{(j)}(\tau-\nu\theta) Y_k^{(j)}(t-\alpha\theta) + \right.$$

$$\left. + \int_0^\infty \sum_{\nu=0}^{S} a_\nu w_k(s,\tau-\nu\theta) Y_k(t-s-\alpha\theta)ds \right\} +$$

$$+ \sum_{\alpha=0}^{\infty} \left\{ c_\alpha \sum_{\nu=S+1}^{N} a_\nu a_\nu \right\} Y_1(t+\tau-\nu\theta-\alpha\theta),$$

and (3.7) follows from (3.8) – (3.12) and

$$\sum_{\alpha=0}^{\infty} c_\alpha \int_0^\infty \sum_{\nu=0}^{S} a_\nu w_k(s,\tau-\nu\theta) Y_k(t-s-\tau-\alpha\theta)ds =$$

$$= \sum_{\alpha=0}^{\infty} c_\nu \int_{\alpha\theta}^\infty \sum_{\nu=0}^{S} a_\nu w_k(u-\alpha\theta,\tau-\nu\theta)\cdot Y_k(t-u)du =$$

$$= \sum_{\alpha=0}^{\infty} c_\nu \int_0^\infty D_k(u-\alpha\theta,\tau) Y_k(t-u)du = \int_0^\infty B_k(u) Y_k(t-u)du,$$

$$\sum_{\nu=S+1}^{N} a_\nu \sum_{\alpha=0}^{\infty} c_\alpha Y_1(t+\tau-\nu\theta-\alpha\theta) Y_k(t-u)du =$$

$$= \sum_{\nu=S+1}^{N} a_\nu \sum_{\mu=\nu-S-1}^{\infty} c_{\mu-\nu+S+1} Y_1(t+\tau-(S+1)\theta-\mu\theta) =$$

$$= \sum_{\nu=S+1}^{N} a_\nu F(\mu,\nu) Y_1(t+\tau-(S+1)\theta-\mu\theta) = \sum_{\mu=0}^{\infty} D_\mu Y_1(t+\tau-(S+1)\theta-\mu\theta).$$

THEOREM 5. *If $T = S\theta$ and $S \le N$, we can obtain $\tilde{Y}_1(t,\tau)$ if we replace S by $S-1$ in the formulae* (3.7) – (3.12).

If τ/θ is not an integer and $S \ge N$, or if $\tau = S\theta$ and $S > N$, then $\tilde{Y}_1(t,\tau)$ is given by:

$$\tilde{Y}(t,\tau) = \sum_{k=1}^{n} \left\{ \sum_{j=0}^{m_k-1}\sum_{\alpha=0}^{\infty} A_{k,\alpha}^{(j)} Y_k^{(j)}(t-\alpha\theta) + \int_0^\infty B_k(u) Y_k(t-u)du \right\} \qquad (3.13)$$

where

$$A_{k,\alpha}^{(j)} = c_\alpha \sum_{\nu=0}^{N} a_\nu w_k^{(j)}(\tau - \nu\theta), \tag{3.14}$$

$B_k(u)$ *is given by* (3.9) *and*

$$D_k(u-\alpha\theta,\tau) = \begin{cases} 0 & 0 \le u < \alpha\theta \\ \sum_{\nu=0}^{S} a_\nu w_k(u-\alpha\theta, \tau-\nu\theta) & u \ge \alpha\theta \end{cases}, \quad \alpha = 0,1,\ldots \tag{3.15}$$

4. Linear extrapolation of autoregressive processes

THEOREM 6. *Let $X(s)$ and $Y(s)$ satisfy the assumptions of Theorem* 3. *Then:*
a) *If $\tau\theta$ is not an integer and $[\tau/\theta] = S$, then*

$$\Phi_{1,\tau}^{X}(\lambda) = \sum_{\nu=0}^{S} c_\nu \Phi_{1,\tau-\nu\theta}^{Y}(\lambda) \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} + \sum_{\mu=S+1}^{S+N} D_\mu e^{i(\tau-\mu\theta)\lambda} \tag{4.1}$$

$$D_\mu = \sum_{\nu=0}^{\mu-S-1} a_\nu c_{\mu-\nu}, \quad \mu = S+1,\ldots,S+N. \tag{4.2}$$

$$\Phi_{k,\tau}^{X}(\lambda) = \sum_{\nu=0}^{S} c_\nu \Phi_{k,\tau-\nu\theta}^{Y}(\lambda) \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda}, \quad k = 2,\ldots,n. \tag{4.3}$$

b) *If $\tau = S\theta$, then it is necessary to replace S by $S-1$ in the equations* (4.1) – (4.3).

Proof. a) It follows from (2.3), (2.6) and (2.8) that

$$\begin{aligned}
\tilde{X}_1(t,\tau) &= \mathrm{Proj}_{H(X,t)} X_1(t+\tau) = \mathrm{Proj}_{H(Y,t)} \sum_{\nu=0}^{\infty} c_\nu Y_1(t+\tau-\nu\theta) \\
&= \sum_{\nu=0}^{S} c_\nu \tilde{Y}_1(t,\tau-\nu\theta) + \sum_{\nu=S+1}^{\infty} c_\nu Y_1(t+\tau-\nu\theta) \\
&= \sum_{\nu=0}^{S} c_\nu \sum_{k=1}^{n} \int_{-\infty}^{+\infty} e^{it\lambda} \Phi_{k,\tau-\nu\theta}(\lambda) dZ_k^Y(\lambda) + \sum_{\nu=S+1}^{\infty} c_\nu \int_{-\infty}^{+\infty} e^{i(t+\tau-\nu\theta)\lambda} dZ_1^Y(\lambda) \\
&= \int_{-\infty}^{+\infty} e^{it\lambda} \left\{ \sum_{\nu=0}^{S} c_\nu \Phi_{1,\tau-\nu\theta}(\lambda) + \sum_{\nu=S+1}^{\infty} c_\nu e^{i(\tau-\nu\theta)\lambda} \right\} \{ \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} dZ_1^X(\lambda) \\
&+ \sum_{k=2}^{n} \int_{-\infty}^{+\infty} e^{it\lambda} \sum_{\nu=0}^{S} c_\nu \Phi_{k,\tau-\nu\theta}(\lambda) \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} dZ_k^X(\lambda).
\end{aligned}$$

The term $\sum_{\nu=0}^{\infty} c_\nu e^{i(\tau-\nu\theta)\lambda} \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda}$ can be represented in the form

$$\sum_{\mu=S+1}^{S+N} \sum_{\nu=0}^{\mu-S-1} a_\nu c_{\mu-\nu} e^{i(\tau-\mu\theta)\lambda} + \sum_{\mu=S+N+1}^{\infty} \sum_{\nu=0}^{N} a_\nu c_{\mu-\nu} e^{i(\tau-\mu\theta)\lambda}$$

and the desired result follows if we use (2.4) and (2.5).

COROLLARY: *If $Y(s)$, $s \in R$, is a nonsingular stationary process with a rational spectrum and $X = AR_Y(N,\theta)$, then the spectral characteristics $\Phi_{k,\tau}(\lambda)$ can be represented in the form:*

$$\Phi^X_{1,\tau}(\lambda = R_1(\lambda) \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda} + \sum_{\mu=S'}^{S'+N-1} D_\mu e^{i(\tau-\mu\theta)\lambda}$$

$$\Phi^X_{k,\tau}(\lambda = R_k(\lambda) \sum_{\nu=0}^{N} a_\nu e^{-i\nu\theta\lambda}, \quad k = 2,\dots,n$$

where $S' = S+1$ if τ/θ is not an integer and $S' = S$ if $\tau = S\theta$..

The functions $R_k(\lambda)$, $k = 1,2,\dots,n$, are rational functions.

THEOREM 7. *Let $X(s)$ and $Y(s)$ be two multidimensional stationary processes satisfying* (2.1) *and let the roots of the equation* (2.2) *be less then one in absolute value. Let $Y(s)$ be a nonsingular random process with a rational spectrum. Suppose we know the values of the stationary process $X(s) = AR_{Y(s)}(N,\theta)$ for $s \le t$. Then, the linear least-squares estimator of $X_1(t+\tau)$, $\tau > 0$, is given by:*

a) *If τ/θ is not an integer and $[\tau/\theta] = S$, then:*

$$\begin{aligned}\tilde{X}_1(t,\tau) &= \sum_{k=1}^{n} \left\{ \sum_{j=0}^{m_k-1} \sum_{\alpha=0}^{N} B^{(j)}_{k,\alpha} X^{(j)}_k(t-\alpha\theta) + \int_0^\infty B_k(u) X_k(t-u)du \right\} \\ &= \sum_{\mu=0}^{N-1} D_\mu X_1(t+\tau-(S+1)-\mu\theta),\end{aligned} \tag{4.4}$$

where

$$B^{(j)}_{k,\alpha} = a_\alpha \sum_{\nu=0}^{S} c_\nu w^{(j)}_k(\tau-\nu\theta), \tag{4.5}$$

$$B_k(u) = \sum_{\alpha=0}^{N} a_\alpha D_k(u-\alpha\theta,\tau), \tag{4.6}$$

$$D_k(u-\alpha\theta,\tau)=\begin{cases}0 & 0\le u<\alpha\theta\\ \sum_{\nu=0}^{S}c_\nu w_k(u-\alpha\theta,\tau-\nu\theta) & u\ge\alpha\theta\end{cases} \tag{4.7}$$

$$D_\mu=\sum_{\alpha=0}^{N}a_\alpha F(\mu,\alpha),\quad \mu=0,1,\dots,N-1 \tag{4.8}$$

$$F(\mu,\nu)=\begin{cases}0 & 0\le\mu<\alpha\\ c_{S+1+\mu-\alpha} & \mu\ge\alpha\end{cases},\quad \alpha=0,1,2\dots,N \tag{4.9}$$

b) *If $\tau=S\theta$, it is necessary to replace S by $S-1$ in the equations* (4.4) – (4.9).

Proof: a) Since $Y(s)$ is a nonsingular stationary process with a rational spectrum, we have

$$\tilde{Y}_1(t,\tau)=\sum_{k=1}^{n}\left\{\sum_{j=0}^{m_k-1}w_k^{(j)}(\tau)Y_k^{(j)}(t)+\int_0^\infty w_k(s,\tau)Y_k(t-s)ds\right\}$$

and

$$\tilde{X}_1(t,\tau)=\sum_{\nu=0}^{S}c_\nu\left\{\sum_{k=1}^{n}\left[\sum_{j=0}^{m_k-1}w_k^{(j)}(\tau-\nu\theta)Y_k^{(j)}(t)+\int_0^\infty w_k(s,\tau-\nu\theta)Y_k(t-s)ds\right]+\right.$$

$$\left.+\sum_{\nu=S1}^{\infty}c_\nu Y_1(t+\tau-\nu\theta)\right\}=$$

$$=\sum_{k=1}^{n}\left\{\sum_{j=0}^{m_k-1}\sum_{\alpha=0}^{N}\left(a_\alpha\sum_{\nu=0}^{S}c_\nu w_k^{(j)}(\tau-\nu\theta)\right)X_k^{(j)}(t-\alpha\theta)+\right.$$

$$\left.+\int_0^\infty\sum_{\nu=0}^{S}c_\nu w_k(s,\tau-\nu\theta)\sum_{\alpha=0}^{N}a_\alpha X_k(t-s-\alpha\theta)ds\right\}+$$

$$+\sum_{\nu=S+1}^{\infty}c_\nu\sum_{\alpha=0}^{N}a_\alpha X_1(t+\tau-\nu\theta-\alpha\theta)$$

and (4.4) follows from (4.5) – (4.9) and

$$\sum_{\alpha=0}^{N}a_\alpha\int_0^\infty\sum_{\nu=0}^{S}c_\nu w_k(s,\tau-\nu\theta)X_k(t-s-\alpha\theta)ds=\sum_{\alpha=0}^{N}a_\alpha\int_0^\infty\sum_{\nu=0}^{S}c_\nu w_k(u-\alpha\theta,\tau-\nu\theta)\cdot$$

$$\cdot X_k(t-u)du=\sum_{\alpha=0}^{N}a_\alpha\int_0^\infty D_k(u-\alpha\theta,\tau)X_k(t-u)du=\int_0^\infty B_k(u)X_k(t-u)du,$$

$$\sum_{\nu=S+1}^{\infty}c_\nu\sum_{\alpha=0}^{N}a_\alpha X_1(t+\tau-\nu\theta-\alpha\theta)=\sum_{\alpha=0}^{N}\sum_{\mu=\alpha}^{\infty}c_{S+1+\mu-\alpha}X_1(t+\tau-(S+1)\theta-\mu\theta)$$

$$=\sum_{\alpha=0}^{N}a_\alpha\sum_{\mu=\alpha}^{\infty}F(\mu,\alpha)X_1(t+\tau-(S+1)\theta-\mu\theta)=\sum_{\mu=0}^{N-1}D_\mu X_1(t+\tau-(S+1)\theta-\mu\theta).$$

The last equation follows from (2.4).

5. Let us now give some examples.

Example 1. Let $x_1(s)$ and $x_2(s)$ be two independent stationary random processes with the rational spectral densities

$$f_1(\lambda) = c_1(\lambda^2 + \alpha_1^2)^{-1},\ f_2(\lambda) = c_2(\lambda^2 + \alpha_2^2)^{-1},\ \ c_1, c_2, \alpha_1, \alpha_2 > 0,\ \ \alpha_1 \neq, \alpha_2,$$

and let $X_1(s) = x_1(s) + x_2(s)$, $X_2(s) = x_1(s) - x_2(s)$. It is easy to see that the spectral density matrix $\|f_{ij}^k(\lambda)\|_{2\times 2}$ of the stationary random process $X(s) = (X_1(s), X_2(s))$ is given by

$$\begin{aligned} &M^X(\lambda) = \|f_{ij}^{(X)}(\lambda)\| = \\ &= \left\| \begin{matrix} c_1(\lambda^2 + \alpha_1^2)^{-1} + c_2(\lambda^2 + \alpha_2^2)^{-1} & c_1(\lambda^2 + \alpha_1^2)^{-1} - c_2(\lambda^2 + \alpha_2^2)^{-1} \\ c_1(\lambda^2 + \alpha_1^2)^{-1} - c_2(\lambda^2 + \alpha_2^2)^{-1} & c_1(\lambda^2 + \alpha_1^2)^{-1} + c_2(\lambda^2 + \alpha_2^2)^{-1} \end{matrix} \right\| \end{aligned} \tag{5.1}$$

In this case $D(\lambda) = 4c_1c_2(\lambda^2 + \alpha_1^2)^{-1}(\lambda^2 + \alpha_2^2)^{-1}$ has no real zeros, $n = 2$, $2L = 0$, $2K = 4$, $m_1 = m_2 = 1$, $2K - 2L = 2(m_1 + m_2)$, and thus $X(s)$ is a nonsingular stationary random process with a rational spectrum.

Suppose we know the values of the process $X(s)$ for $s \leq t$. Using Theorem 1 we have:

$$\begin{aligned} &\Phi_1(\lambda) = \omega_1(\lambda) = K_1,\ \ \Phi_2(\lambda) = \omega_2(\lambda) = K_2, \\ &\Psi_1(\lambda) = (e^{i\tau\lambda} - K_1)f_{11}(\lambda) - K_2 f_{21}(\lambda), \\ &\Psi_2(\lambda) = (e^{i\tau\lambda} - K_1)f_{12}(\lambda) - K_2 f_{22}(\lambda). \end{aligned}$$

From the fact that the functions $\Psi_1(\lambda)$ and $\Psi_2(\lambda)$ are analytic in the upper half-plane, it follows that

$$2K_1 = 2K_1(T) = e^{-\tau\alpha_1} + e^{-\tau\alpha_2},\ \ 2K_2 = K2(\tau) = e^{-\tau\alpha_1} - e^{-\tau\alpha_2} \tag{5.2}$$

Then, we have $w_1^{(0)}(\tau) = K_1(\tau)$, $w_2^{(0)}(\tau) = K_2(\tau)$, $w_1(s,\tau) = w_2(s,\tau) = 0$, and

$$2X_1(t,\tau) = (e^{-\tau\alpha_1} + e^{-\tau\alpha_2})X_1(t) + e^{-\tau\alpha_1} - e^{-\tau\alpha_2}X_2(t). \tag{5.3}$$

Example 2. Let $X(s)$ be the process given in the example 1, $Y(s) = X(s) - \beta X(s-1)$, $|\beta| < 1$, and suppose we know the values of the process $X(s)$ for $s \leq t$.

In this case: $N = 1$, $a_0 = 1$, $a_1 = -\beta$, $Y(s) = MA_X(1,\theta)$, $X(s) = \sum_{\alpha=0}^{\infty} \beta^\alpha Y(s - \alpha\theta)$.

a) If $0 < \tau < \theta$ $(0 = S < N = 1)$, using the formulae (3.8) – (3.12) we obtain

$$\begin{aligned} &A_{1,\alpha}^{(0)} = \beta^\alpha a_0 w_1^{(0)}(\tau) = \beta^\alpha K_1(\tau),\ \ A_{2,\alpha}^{(0)} = \beta^\alpha a_0 w_2^{(0)}(\tau) = \beta^\alpha K_2(\tau), \\ &B_1(u) = B_2(u) = 0,\ \ D_\mu = a_1 F(\mu, 1) = a_1 c_\mu = a_1\beta^\mu = -\beta^{\mu+1}, \end{aligned}$$

where $K_1(\tau)$ and $K_2(\tau)$ were given in the example 1, and finally

$$\tilde{Y}_1(t,\tau) = \frac{1}{2}(e^{-\tau\alpha_1} + e^{-\tau\alpha_2}) \sum_{\alpha=0}^{\infty} \beta^\alpha Y_1(t-\alpha\theta) + \frac{1}{2}(e^{-\tau\alpha_1} - e^{-\tau\alpha_2}) \sum_{\alpha=0}^{\infty} \beta^\alpha Y_2(t-\alpha\theta) - \sum_{\mu=0}^{\infty} \beta^{\mu+1} Y_1(t+\tau-\theta-\mu\theta). \tag{5.4}$$

b) If $\tau = \theta$, $(S = N = 1)$, then replacing τ by θ, $\tilde{Y} - 1(t,\theta)$ can be obtained from (5.4).

c) If $\tau = \theta$, using the formulae (3.9) and (3.13) – (3.15) we obtain:

$$B_1(u) = B_2(u) = 0,$$
$$2A^{(0)}_{1,\alpha} = 2\beta^\alpha(a_0 K_1(\tau-\theta)) = \beta^\alpha(e^{-\tau\alpha_1} + e^{-\tau\alpha_2} - \beta e^{-(\tau-\theta)\alpha_1} - \beta e^{-(\tau-\theta)\alpha_2}),$$
$$2A^{(0)}_{2,\alpha} = 2\beta^\alpha(a_0 K_2(\tau-\theta)) = \beta^\alpha(e^{-\tau\alpha_1} - e^{-\tau\alpha_2} - \beta e^{-(\tau-\theta)\alpha_1} - \beta e^{-(\tau-\theta)\alpha_2}),$$

and finally

$$2\tilde{Y}_1(t,\tau) = (e^{\tau\alpha_1} + e^{-\tau\alpha_2} - \beta e^{\tau\alpha_1} - \beta e^{-\tau\alpha_2} \sum_{\alpha=0}^{\infty} \beta^\alpha Y_1(t-\alpha\theta) + (e^{\tau\alpha_1} - e^{-\tau\alpha_2} - \beta e^{-(\tau-\theta)\alpha_1} - \beta e^{-(\tau-\theta)\alpha_2}) \sum_{\alpha=0}^{\infty} \beta^\alpha Y_2(t-\alpha\theta) \tag{5.5}$$

Example 3. Let $Y(s)$ be a process with the spectral density matrix (5.1) and $X(s) - \beta X(s-1) = Y(s)$, $|\beta| < 1$. Then $X(s) = AR_Y(1,\theta)$, $N = 1$, $a_0 = 1$, $a_1 = -\beta$ and using the formulae (4.4) – (4.9) we obtain:

$$2B^{(0)}_{1,\alpha} = 2a_\alpha \sum_{\nu=0}^{S} \beta^\nu K_1(\tau-\nu\theta) = a_\alpha \sum_{\nu=0}^{S} \beta^\nu (e^{-(\tau-\nu\theta)\alpha_1} + e^{-(\tau-\nu\theta)\alpha_2})$$
$$2B^{(0)}_{2,\alpha} = 2a_\alpha \sum_{\nu=0}^{S} \beta^\nu K_2(\tau-\nu\theta) = a_\alpha \sum_{\nu=0}^{S} \beta^\nu (e^{-(\tau-\nu\theta)\alpha_1} + e^{-(\tau-\nu\theta)\alpha_2})$$

$D_\mu = a_0 F(\mu,0) + a_1 F(\mu,1) = a_0 c_{S+1-\mu} + a_1 c_{S+\mu} = \beta^{S+1+\mu} - \beta\beta^{S+\mu} = 0$, $\mu \geq 1$
$D_0 = a_0 c_{S+1} = \beta^{S+1}$ and

$$\tilde{X}_1(t,\tau) = \frac{1}{2}\sum_{\nu=0}^{S} \beta^\nu \{e^{-(\tau-\nu\theta)\alpha_1} + e^{-(\tau-\nu\theta)\alpha_2}\}(X_1(t) - \beta X_1(t-\theta)) + \frac{1}{2}\sum_{\nu=0}^{S} \beta^\nu \{e^{-(\tau-\nu\theta)\alpha_1} + e^{-(\tau-\nu\theta)\alpha_2}\}(X_2(t) - \beta X_2(t-\theta)) + \beta^{S+1} X_1(t+\tau-(S+1)\theta). \tag{5.6}$$

REFERENCES

[1] A. W. Fuller, *Introduction to Statistical Time Series*, New York, 1976.

[2] P. Mladenović, *O linearnoj prognozi višedimenzionalnih stacionarnih procesa sa neracionalnim spektrom*, Magistarski rad, PMF, Beograd, 1981.

[3] A. M. Yaglom, *Stationary Random Functions*, New York, 1962.

[4] Й. Малишич, *Об экспортровании по всему прошлому для некоторых стационарных процессов с нерациональными плотностьями*, Мат. Весник 2, (Београд) (1978).

[5] А. Ю. Розанов, *Стационарные случайные процессы*, Москва, 1963.

[6] А. М. Яглом, *Экстраполирование, интерполирование и фильтрация стационарных случайных процессов с рациональной спектральной плотностью*, Труды ММО 4, (1955), 237–278.

[7] А. М. Яглом, *Эффективные решения линейных аппроксимационных задач для многомерных стационарных процессов с рациональным спектром*, Теория вероят. и ее прим. V, 3, 256–292. 1960.

Odsek za matematiku
Prirodno-matematički fakultet
11000 Beograd
Jugoslavija

(Received 23 12 1981)