Electronic Journal: Southwest Journal of Pure and Applied Mathematics
Internet: http://rattler.cameron.edu
ISSN 1083-0464
Vol. No. 01, 1995, pp. 13–16.
Submitted: March 16, 1995. Published: June 30, 1995.

DIFFERENTIATED SHIFT-INVARIANT MULTIVARIATE INTEGRAL OPERATORS

George A. Anastassiou

Abstract. In a recent paper [1] the author, together with H. Gonska, introduced some wavelet-type, very general multivariate integral operators over $\mathbb{R}^d$, $d \geq 1$, and studied their various properties such as shift-invariance, global smoothness preservation and optimality, convergence to the unit, differentiation, and preservation of continuous multivariate probability distribution functions. These operators are introduced through a convolution-like iteration of another very general multivariate operator with a general multivariate scaling type function. In this paper the author gives sufficient conditions, so that the *partial derivatives of the above described multivariate operators* possess most of the above listed nice properties of their originals. In particular, a sufficient condition is given so that a corresponding "global smoothness preservation" sharp multivariate inequality is attained. The corresponding convergence of differentiated operators to the differentiated unit operator is given with rates of convergence.

(1991) A.M.S. (MOS) Subject Classification Codes. 60, 62, 28.

Key Words and Phrases. Wavelet integral operators, Probability distribution functions.

Main Results.

Let $X := C_U(\mathbb{R}^d)$, $d \geq 1$ be the space of uniformly continuous functions from $\mathbb{R}^d$ into $\mathbb{R}$.

For $f \in X$, define the first order modulus of continuity of f by

$$\omega_1(f, \delta) := \sup_{\substack{x,y \in \mathbb{R}^d \\ \|x-y\| \leq \delta}} |f(x) - f(y)|, \delta > 0,$$

where $\|\cdot\|$ is an arbitrary norm in $\mathbb{R}^d$. It is known [1] that $\omega_1(f, \delta) < \infty$, $\forall \delta > 0$. Here $C(\mathbb{R}^d)$ denotes the space of continuous functions from $\mathbb{R}^d$ into $\mathbb{R}$. Let $\{\ell_k\}_{k \in \mathbb{Z}}$ be a sequence of linear operators from $C(\mathbb{R}^d)$ into itself such that

$$\ell_k(f, x) = \ell_0(f(2^{-k}\cdot), x), \quad \text{all } x \in \mathbb{R}^d, \ f \in C(\mathbb{R}^d). \tag{1}$$

Assume that

$$\ell_0 : C^{(\rho)}(\mathbb{R}^d) \to C^{(\rho)}(\mathbb{R}^d), \quad \rho \in \mathbb{N} \quad \text{be fixed}, \tag{2}$$

where $C^{(\rho)}(\mathbb{R}^d)$ is the space of ρ-times continuously differentiable functions.

In this article we consider *only* $f \in C^{(\rho)}(\mathbb{R}^d)$ such that $\frac{\partial^\rho f}{\partial x_{i_1}^{j_1} \cdots \partial x_{i_m}^{j_m}} \in X$, where fixed $i_1, \ldots, i_m \in \{1, \ldots, d\}$ be such that $i_1 < \cdots < i_m$, and fixed $j_1, \ldots, j_m \in \mathbb{N}$ be such that $\sum_{r=1}^m j_r = \rho$.

From now on we will denote by

$$f^\partial := \frac{\partial^\rho f}{\partial x_{i_1}^{j_1} \cdots \partial x_{i_m}^{j_m}}, \quad f \in C^{(\rho)}(\mathbb{R}^d), \tag{3}$$

Department of Mathematics, Memphis State University, Memphis, TN
Email: anastasg@ kermes.msci.memst.edu
Copyright ©1995 by Cameron University

Typeset by $\mathcal{AMS}$-TEX

where the mixed partial is defined as above.

Let φ be a bounded real valued function of compact support $\subseteq \times_{i=1}^{d}[-a_i, a_i]$, $a_i > 0$. We assume that $\varphi \geq 0$, φ is Lebesgue measurable and

$$\underbrace{\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty}}_{d-\text{fold}} \varphi(x_1 - u_1, x_2 - u_2, \ldots, x_d - u_d)\, du_1 \ldots du_d = 1,$$

$$\text{all } (x_1, \ldots, x_d) \in \mathbb{R}^d \tag{4}$$

((4) same as

$$\int_{\mathbb{R}^d} \varphi(x - u)\, du = 1, \text{all } x \in \mathbb{R}^d;$$

where $x = (x_1, \ldots, x_d)$, $u = (u_1, \ldots, u_d)$). One can easily prove that

$$\int_{\mathbb{R}^d} \varphi(u) du = 1. \tag{5}$$

Example. For $i = 1, \ldots, d$ take the characteristic function

$$\varphi_i(x_i) := \chi_{[-\frac{1}{2}, \frac{1}{2})}(x_i) = \begin{cases} 1, & x_i \in \left[-\frac{1}{2}, \frac{1}{2}\right) \\ 0, & \text{else.} \end{cases}$$

Define

$$\varphi^*(x) := \prod_{i=1}^{d} \varphi_i(x_i), \quad \text{all } x = (x_1, \ldots, x_d) \in \mathbb{R}^d.$$

Note that φ^* is bounded, supp $\varphi^* \subseteq \left[-\frac{1}{2}, \frac{1}{2}\right]^d$, $\varphi^* \geq 0$ and φ^* is Lebesgue measurable. Also

$$\underbrace{\int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty}}_{d-\text{fold}} \prod_{i=1}^{d} \varphi_i(x_i - u_i)\, du_1 \ldots du_d = 1,$$

i.e.,

$$\int_{\mathbb{R}^d} \varphi^*(x - u) du = 1, \quad \text{all } x \in \mathbb{R}^d.$$

Let $\{\mathcal{L}_k\}_{k \in \mathbb{Z}}$ be the sequence of linear operators acting on $C^{(\rho)}(\mathbb{R}^d)$ and defined as follows

$$\mathcal{L}_0(f, x) := \int_{\mathbb{R}^d} \ell_0(f, u) \varphi(x - u)\, du \tag{6}$$

and

$$\mathcal{L}_k(f, x) := \mathcal{L}_0(f(2^{-k}\cdot); 2^k x), \tag{7}$$

all $x \in \mathbb{R}^d$, $k \in \mathbb{Z}$. I.e.,

$$\mathcal{L}_k(f, x) = \int_{\mathbb{R}^d} \ell_k(f, u) \varphi(2^k x - u)\, du. \tag{8}$$

For our convenience we shall use the notation

$$\int_{-\vec{a}}^{\vec{a}} \bullet = \int_{\times_{i=1}^{d}[-a_i, a_i]} \bullet. \tag{9}$$

Furthermore, we obtain that

$$\begin{aligned} \mathcal{L}_k^{\partial}(f, x) &= 2^{k\rho} \cdot \int_{-\vec{a}}^{\vec{a}} \ell_k^{\theta}(f, 2^k x - u) \varphi(u)\, du \\ &= 2^{k\rho} \cdot \int_{\mathbb{R}^d} \ell_k^{\partial}(f, 2^k x - u) \varphi(u)\, du \\ &= 2^{k\rho} \cdot \int_{\mathbb{R}^d} \ell_k^{\partial}(f, u) \varphi(2^k x - u)\, du, \text{all } x \in \mathbb{R}^d, \ k \in \mathbb{Z}. \end{aligned} \tag{10}$$

Assumption. Take $f \in C^{(\rho)}(\mathbb{R}^d)$, $\rho \in \mathbb{N}$ fixed, such that $f^\partial \in X$. For fixed $\alpha > 0$ we assume that

$$\sup_{\substack{u,y\in\mathbb{R}^d \\ \|u-y\|_\infty \le \alpha}} |\ell_0^\partial(f,u) - f^\partial(y)| \le \omega_{1,\infty}\left(f^\partial, \frac{m\alpha+n}{2^r}\right), \tag{11}$$

is true for f as above, $m \in \mathbb{N}$, $n \in \mathbb{Z}_+$, $r \in \mathbb{Z}$, where $\omega_{1,\infty}$ is the modulus of continuity ω_1 defined with respect to $\|\cdot\|_\infty$.

Here we present our results without proofs due to their length. We intend to show proofs elsewhere.

Definition 1. Let $f_\alpha(\cdot) := f(\cdot+\alpha)$, $\alpha \in \mathbb{R}^d$ and ϕ be an operator. If $\phi(f_\alpha) = (\phi f)_\alpha$, then ϕ is called a *shift invariant operator.*

Proposition 1. *Assume that*

$$\ell_0^\partial\left(f(2^{-k}\cdot+\alpha); 2^k u\right) = \ell_0^\partial\left(f(2^{-k}\cdot); 2^k(u+\alpha)\right), \tag{12}$$

all $k \in \mathbb{Z}$, $\alpha \in \mathbb{R}^d$ fixed, all $u \in \mathbb{R}^d$; any $f \in C^{(\rho)}(\mathbb{R}^d)$; $\rho \in \mathbb{N}$ be given. Then $\mathcal{L}_k^\partial$, any $k \in \mathbb{Z}$ is a shift invariant operator.

Next we establish the property of "*global smoothness preservation*" of operators $\mathcal{L}_k^\partial$, $k \in \mathbb{Z}$.

Theorem 1. *For any $f \in C^{(\rho)}(\mathbb{R}^d)$, such that $f^\partial \in X$, $\rho \in \mathbb{N}$ fixed, and any $u \in \mathbb{R}^d$ assume that*

$$|\ell_0^\partial(f, x-u) - \ell_0^\partial(f, y-u)| \le \omega_1(f^\partial, \|x-y\|), \quad \text{all } x, y \in \mathbb{R}^d, \tag{13}$$

where $\|\cdot\|$ is an arbitrary norm in $\mathbb{R}^d$. Then

$$\omega_1((\mathcal{L}_k f)^\partial, \delta) \le \omega_1(f^\partial, \delta), \tag{14}$$

any $\delta > 0$.

Optimality of (14) follows.

Theorem 2. *Let $i_1, \dots, i_m \in \{1, \dots, d\}$ be such that $i_1 < \cdots < i_m$, and $j_1, \dots, j_m \in \mathbb{N}$ be such that $\sum_{r=1}^m j_r = \rho$. Consider $g_{j_1}(x) := \frac{x^{j_1+1}}{(j_1+1)!}$, and $g_{j_r}(x) := \frac{x^{j_r}}{j_r!}$, $r = 2, \dots, m$; all $x \in \mathbb{R}$. Denote by $f^\partial := \frac{\partial^\rho f}{\partial x_{i_1}^{j_1} \dots \partial x_{i_m}^{j_m}}$, for any $f \in C^{(\rho)}(\mathbb{R}^d)$. Denote also by $pr_{i_r} : \mathbb{R}^d \ni (x_1, \dots, x_d) \to x_{i_r}$, $r = 1, \dots, m$ the projection onto the i_r coordinate. For all $x, y, u \in \mathbb{R}^d$ assume that*

$$\left(\ell_0\left(\left(\sum_{r=1}^m g_{j_r} \circ pr_{i_r}\right)(2^{-k}\cdot)\right)\right)^\partial (2^k x - u)$$

$$-\left(\ell_0\left(\left(\prod_{r=1}^m g_{j_r} \circ pr_{i_r}\right)(2^{-k}\cdot)\right)\right)^\partial (2^k y - u) = 2^{-k\rho}\cdot(x_{i_1} - y_{i_1}). \tag{15}$$

Then

$$\omega_1\left(\left(\mathcal{L}_k\left(\prod_{r=1}^m g_{j_r} \circ pr_{i_r}\right)\right)^\partial; \delta\right) = \omega_1\left(\left(\prod_{r=1}^m g_{j_r} \circ pr_{i_r}\right)^\partial; \delta\right), \tag{16}$$

for any $\delta > 0$; $k \in \mathbb{Z}$. That is establishing that (14) is sharp!

The convergence of $\frac{\partial^\rho \mathcal{L}_k}{\partial x_{i_1}^{j_1} \dots \partial x_{i_m}^{j_m}} (\sum_{r=1}^m j_r = \rho)$ to $\frac{\partial^\rho}{\partial x_{i_1}^{j_1} \dots \partial x_{i_m}^{j_m}}$ operator, as $k \to +\infty$, with rates given in the following.

Theorem 3. *For $f \in C^{(\rho)}(\mathbb{R}^d)$, $d \ge 1$, such that $f^\partial \in C_U(\mathbb{R}^d)$, under the assumption (11), it holds*

$$|\mathcal{L}_k^\partial(f, x) - f^\partial(x)| \le \omega_{1,\infty}\left(f^\partial, \frac{ma+n}{2^{k+r}}\right), \tag{17}$$

where $m \in \mathbb{N}$, $n \in \mathbb{Z}_+$, $k, r \in \mathbb{Z}$, and $a := \max(a_i)$, $i = 1, \dots, d$.

Again, due to their length we intend to present applications of the above results elsewhere.

References

1. G.A. Anastassiou and H.H. Gonska, *On some shift-invariant integral operators, multivariate case*, Proceedings of International Conference in Approximation Theory and Probability, Santa Barbara, California, May 20-22, 1993 (1994), Plenum, New York, 41–64.
2. H. Bauer, *Maß-und Integrationstheorie* (1990), de Gruyter, Berlin.
3. C. Cottin and H.H. Gonska, *Simultaneous approximation and global smoothness preservation*, Proceedings of the Second International Conference in Functional Analysis and Approximation Theory Acquafredda di Maratea-Potenza-Italy, 1992. *Supplemento ai Rendiconti del Circolo Matematico di Palermo*, Serie II, Numero 33, (1993), 259–279..
4. A.N. Shiryayev, *Probability*, Springer-Verlag, New York, Berlin, 1984.