Electronic Journal: Southwest Journal of Pure and Applied Mathematics
Internet: http://rattler.cameron.edu/swjpam.html
ISSN 1083-0464
Issue 1, June 2001, pp. 38–53.
Submitted: February 3, 2001. Published: June 30, 2001.

INTERPOLATION AND THE LAGUERRE–PÓLYA CLASS

Andrew Bakan, Thomas Craven and George Csordas

Abstract. A long-standing open problem in the theory of distribution of zeros of real entire functions requires the characterization of all real sequences $T = \{\gamma_k\}_{k=0}^{\infty}$ such that for any real polynomial $p(x) := \sum_0^n a_k x^k$, the polynomial $\sum_0^n \gamma_k a_k x^k$ has no more nonreal zeros than $p(x)$ has. The main result of this paper shows that under a mild growth restriction, an entire function of exponential type $f(z)$ for which the sequence $T = \{f(k)\}_{k=0}^{\infty}$ satisfies the above condition must have only real zeros. The paper concludes with some applications to the Riemann Hypothesis.

1991 A.M.S. Subject Classification Codes:. 41A05, 26C10, 30D15

Keywords and Phrases:. Laguerre-Pólya class, multiplier sequences, distribution of zeros of entire functions, interpolation, Riemann Hypothesis

Andrew Bakan, Institute of Mathematics, National Academy of Sciences of Ukraine
E-mail: bakan@imat.gluk.apc.org
Thomas Cravan, Department of Mathematics, University of Hawaii, Honolulu, HI
E-mail: tom@math.hawaii.edu
George Csordas, Department of Mathematics, University of Hawaii, Honolulu, HI
E-mail: george@math.hawaii.edu
©2001 Cameron University

Typeset by $\mathcal{AMS}$-TEX