Electronic Journal: Southwest Journal of Pure and Applied Mathematics
Internet: http://rattler.cameron.edu/swjpam.html
ISBN 1083-0464
Issue 1 July 2004, pp. 10 – 32
Submitted: September 10, 2003. Published: July 1, 2004

ORLICZ-SOBOLEV SPACES WITH ZERO BOUNDARY VALUES ON METRIC SPACES

NOUREDDINE AÏSSAOUI

Abstract. In this paper we study two approaches for the definition of the first order Orlicz-Sobolev spaces with zero boundary values on arbitrary metric spaces. The first generalization, denoted by $M_{\Phi}^{1,0}(E)$, where E is a subset of the metric space X, is defined by the mean of the notion of the trace and is a Banach space when the N-function satisfies the Δ_2 condition. We give also some properties of these spaces. The second, following another definition of Orlicz-Sobolev spaces on metric spaces, leads us to three definitions that coincide for a large class of metric spaces and N-functions. These spaces are Banach spaces for any N-function.

A.M.S. (MOS) Subject Classification Codes.46E35, 31B15, 28A80.
Key Words and Phrases. Orlicz spaces, Orlicz-Sobolev spaces, modulus of a family of paths, capacities.

Ecole Normale Supérieure, B.P 5206 Ben Souda, Fès, Maroc
n.aissaoui@caramail.com
Copyright ©2004 by Cameron University