Electronic Journal: Southwest Journal of Pure and Applied Mathematics
Internet: http://rattler.cameron.edu/swjpam.html
ISSN 1083-0464
Issue 2, December 2001, pp. 19–23.
Submitted: October 25, 2001. Published: December 31 2001.

CONVERGENCE THEOREMS FOR NEWTON-LIKE METHODS USING DATA FROM A SET OR A SINGLE POINT AND OUTER INVERSES

Ioannis K. Argyros

Abstract. Semilocal convergence theorems for Newton-like methods are provided using information from a set or a single point and outer inverses.

A.M.S. (MOS) Subject Classification Codes. 65H10, 47H17, 49M15.

Key Words and Phrases. Newton-like methods, Banach space, semilocal convergence theorem, outer inverse.

1. Introduction

In this study we are concerned with the problem of approximating a locally unique solution x^* of the equation

$$LF(x) = 0, \tag{1}$$

where F is a Fréchet-differentiable operator defined on an open convex subset D of a Banach space X with values in a Banach space Y, $L \in L(X,Y)$ the space of bounded linear operators from X into Y.

We consider Newton-like methods of the form

$$x_{n+1} = x_n - A(x_n)^{\#} F(x_n) \quad (n \geq 0) \tag{2}$$

for generating a sequence converging to x^*. Here $A(x_n) \in L(X,Y)$ $(n \geq 0)$ is an approximation of the Fréchet-derivative $F'(x_n)$ $(n \geq 0)$ and $A(x_n)^{\#}$ denotes an outer inverse of $A(x_n)$, i.e., $A(x_n)^{\#} A(x_n) A(x_n)^{\#} = A(x_n)^{\#}$. This setting includes generalized Newton methods for undetermined systems [1], [3].

Semilocal convergence theorems for Newton-like methods of the form (2) are provided using information from a set or a single point.

Department of Mathematics, Cameron University, Lawton, OK 73505, U.S.A.
E-mail Address: ioannisa@cameron.edu
©2001 Cameron University

Typeset by $\mathcal{AMS}$-TEX

2. Convergence Analysis

Define parameters r_i, $i = 0, 1, \ldots, 5$, h_0 and γ by

$$
\begin{aligned}
&r_0 = \frac{1}{\gamma}\left(1 - \sqrt[3]{\frac{2\gamma}{M+L}}\right), \quad M + L \neq 0, \quad r_1 = \frac{1-\ell}{L}, \quad L \neq 0,\\
&r_2 = \frac{1}{\gamma}, \quad \gamma \neq 0, \quad r_3 = \min\{r_0, r_1, r_2\},\\
&r_3 = r_0 \text{ if } L = \gamma = 0, \quad r_3 = \min\{r_0, r_2\} \text{ if } L = 0,\\
&r_3 = \min\{r_0, r_1\} \text{ if } \gamma = 0, \quad r_4 = \min\{r_3, r^{**}\},\\
&r_5 = \min\{r_1, r_2\}, \quad r_5 = r_1 \text{ if } \gamma = 0,\\
&r_5 = r_2 \text{ if } L = 0, \quad r_5 = \infty \text{ if } L = \gamma = 0,\\
&h_0 = (M+L)\eta - \frac{1}{2}(1-b)^2, \quad \text{for } L \geq 0,\ M \geq 0\\
&\gamma = \sup_{k>1}\left\|\frac{1}{k!}A^{\#}F^{(k)}(x_0)\right\|, \quad x_0 \in X \quad \text{(if it exists)}
\end{aligned}
$$

where $A^{\#}$ is a bounded outer inverse of A $(= A(x_0))$ and functions f, g, f_1, f_2 by

$$
\begin{aligned}
f(r) &= \frac{\sigma}{2}r^2 - (1-b)r + \eta, \quad \sigma = \max\{K, M+L\},\\
g(r) &= Lr + \ell - 1,\\
f_1(r) &= \frac{M+L}{2}r^2 - (1-b)r + \eta,\\
f_2(r) &= \frac{\gamma r^2}{(1-\gamma r)^3} - (1-b)r + \eta,
\end{aligned}
$$

where r^*, r^{**} are the zeros of function f_1.

Assume:

(C) $r_0 \geq 0$, $h_0 \leq 0$ and $r^* \leq r_3$

Note that under condition (C), $0 \leq r^* \leq r^{**}$, $f_1(r) \geq 0$, $g(r) < 0$ and $\frac{2\gamma}{(1-\gamma r)^3} \equiv K_0 \leq M + L$ on $[0, r^*]$.

or

(C_1) $r_0 \leq 0$ (or $M + L = 0$) and there exists $r_6 \in (0, r_5)$ or $r_6 \in [r_0, r_5))$ such that $f_2(r_6) \leq 0$.

Denote by r_7 the zero of function f_2 in $(0, r_6)$. (Such a zero exists by the intermediate value theorem.) Note that under condition (C_1), $f_2(r) \geq 0$, $g(r) < 0$ and $M + L \leq K_0$ on $[0, r_6]$.

We can state and prove the following semilocal convergence theorem for Newton-like methods using information from a set and outer inverses.

Theorem 1. *Let $F: D \subseteq X \to Y$ be twice-Fréchet-differentiable and let $A(x) \in L(X, Y)$ $(x \in D)$ be an approximation of $F'(x)$. Assume*:

(a) *there exists an open convex subset D_0 of D, $x_0 \in D_0$, a bounded outer inverse $A^{\#}$ of A $(= A(x_0))$ and constants η, $K > 0$, M, L, μ, $\ell \geq 0$ such that for al $x \in D_0$*

the following conditions hold:

$$\|A^{\#}F(x_0)\| \leq \eta, \tag{3}$$

$$\|A^{\#}F''(x)\| \leq K, \tag{4}$$

$$\|A^{\#}(F'(x) - A(x))\| \leq M\|x - x_0\| + \mu \tag{5}$$

$$\|A^{\#}(A(x) - A)\| \leq L\|x - x_0\| + \ell, \tag{6}$$

$$b = \mu + \ell < 1, \tag{7}$$

$$h = \sigma\eta \leq \frac{1}{2}(1-b)^2 \tag{8}$$

and

$$\overline{U}(x_0, t^*) \subseteq D_0, \tag{9}$$

where, t^* is the small zero of function f.

Then,

(1) *Newton-like method $\{x_n\}$ $(n \geq 0)$ generated by (2) is well defined, remains in $U(x_0, t^*)$ for all $n \geq 0$ and converges to a solution $x^* \in \overline{U}(x_0, t^*)$ of equation $A^{\#}F(x) = 0$, with*

$$A(x_n)^{\#} = (I + A^{\#}(A(x_n) - A))^{-1}A^{\#};$$

(2) *equation $A^{\#}F(x) = 0$ has a unique solution x^* in $\widetilde{U} \cap \{R(A^{\#}) + x_0\}$, where*

$$\widetilde{U} = \begin{cases} \overline{U}(x_0, t^*) \cap D_0, & \text{if } h = \frac{1}{2}(1-b)^2 \\ U(x_0, t^{**}) \cap D_0, & \text{if } h < \frac{1}{2}(1-b)^2, \end{cases}$$

where t^{**} is the large zero of function f and

$$R(A^{\#}) + x_0 = \{x + x_0; x \in R(A^{\#})\};$$

(3) *the following error bounds hold for all $n \geq 0$*

$$\|x_{n+1} - x_n\| \leq t_{n+1} - t_n$$

and

$$\|x_n - x^*\| \leq t^* - t_n,$$

where,

$$t_0 = 0, \quad t_{n+1} = t_n - \frac{f(t_n)}{g(t_n)}.$$

Proof. Lipschitz condition (3.2) in [3] follows from (4). The rest of the theorem follows exactly as Theorem 3.1 in [3].

That completes the proof of the theorem.

We can show the following semilocal result for Newton-like methods using information from a set of a single point and outer inverses.

Theorem 2. *Let $F: X \to Y$ be analytic on some open convex subset D of X and consider $A(x) \in L(X,Y)$ $(x \in D)$ to be an approximation of $F'(x)$. Assume*:

(a) *there exists an open convex subset D_0 of D, $x_0 \in D_0$, a bounded outer inverse $A\#$ of A $(= A(x_0))$ and constants $\eta > 0$, M, L, μ, $\ell \geq 0$ such that for all $x \in D_0$ conditions (3), (5), (6), and (7) hold*;

(b) *conditions (C) (or (C_1)) hold*;

(c) $\overline{U}(x_0, r^*) \subseteq D_0$ *(or* $\overline{U}(x_0, r_1) \subseteq D_0$*)*.

Then,

(1) *Newton's method $\{x_n\}$ $(n \geq 0)$ generated by (2) is well defined, remains in $U(x_0, r^*)$ (or $U(x_0, r_7)$) for all $n \geq 0$ and converges to a solution $x^* \in \overline{U}(x_0, r^*)$ (or $\overline{U}(x_0, r_7)$) of equation $A^{\#}F(x) = 0$ with $A(x_n)^{\#} = (I + A^{\#}(A(x_n) - A))^{-1} A^{\#}$*;

(2) *the equation $A^{\#}F(x) = 0$ has a unique solution in $\widetilde{U}_1 \cap \{R(A^{\#}) + x_0\}$ (or $\widetilde{U}_2 \cap \{R(A^{\#} + x_0\})$, where*

$$\widetilde{U}_1 = \begin{cases} \overline{U}(x_0, r^*) \cap D_0 & \text{if } h_0 = 0 \\ U(x_0, r_4) \cap D_0 & \text{if } h_0 < 0, \end{cases}$$

$$\widetilde{U}_2 = \begin{cases} \overline{U}(x_0, r_5) \cap D_0 & r_6 < r_5 \\ U(x_0, r_6) \cap D_0 & r_5 = r_6; \end{cases}$$

(3) *the following error bound hold for all $n \geq 0$*:

$$\|x_{n+1} - x_n\| \leq r_{n+1} - r_n \quad (\text{or } r^1_{n+1} - r^1_n), \tag{10}$$

$$\|x_n - x^*\| \leq r^* - r_n \quad (\text{or } r_6 - r^1_n), \tag{11}$$

where,

$$r_0 = 0, \quad r_{n+1} = r_n - \frac{f_1(r_n)}{g(r_n)}$$

and

$$r^1_0 = 0, \quad r^1_{n+1} = r^1_n - \frac{f_2(r^1_n)}{g(r^1_n)}.$$

Proof. Since $\gamma r^* < 1$ under (C) (or $\gamma r_6 < 1$ under (C_1)), we find for all $x \in U(x_0, r^*)$ (or r_6) that

$$\begin{aligned} \|A^{\#}F''(x)\| &= \left\| \sum_{n=0}^{\infty} \frac{1}{k!} A^{\#} F^{(k+2)}(x_0)(x - x_0)^k \right\| \\ &\leq \gamma \sum_{k=0}^{\infty} (k+2)(k+2)(\|x - x_0\|)^k \leq K_0. \end{aligned}$$

Estimates (10) and (11) are shown exactly as in Theorem 1 since under condition (C) $r^* \leq r^{**}$, $\frac{2\gamma}{(1-\gamma r)^3} = K_0 \leq M + L$, $f_1(r) \geq 0$, $g(r) < 0$ on $[0, r^*]$ (or under (C_1): $f_2(r) \geq 0$, $g(r) < 0$ and $M + L < K_0$ on $[0, r_6]$).

The rest follows exactly as in Theorem 1.

That completes the proof of the theorem.

REFERENCES

[1] Argyros, I.K., Convergence domains for some iterative processes in Banach space using outer and generalized inverses, *Journ. Computational Anal. and Applic.* **1**, 1 (1999), 87–104.

[2] Argyros, I.K. and Szidarovszky, F., *The Theory and Application of Iteration Methods*, CRC Press, Inc., Boca Raton, Florida, U.S.A., 1993.

[3] Chen, X. and Nashed, M.Z., Convergence of Newton-like methods for singular operator equations using outer inverses, *Numer. Math.* **66** (1993), 235–257.

[4] Kantorovich, L.V. and Akilov, G.P., *Functional Analysis*, Pergamon Press, Oxford, 1982.