Theory and Applications of Categories, Vol. 21, No. 10, 2008, pp. 182–190.

DESCENT FOR COMPACT 0-DIMENSIONAL SPACES

Dedicated to Walter Tholen on the occasion of his 60th birthday

GEORGE JANELIDZE[1] AND MANUELA SOBRAL[2]

Abstract. Using the reflection of the category $\mathcal{C}$ of compact 0-dimensional topological spaces into the category of Stone spaces we introduce a concept of a fibration in $\mathcal{C}$. We show that: (i) effective descent morphisms in $\mathcal{C}$ are the same as the surjective fibrations; (ii) effective descent morphisms in $\mathcal{C}$ with respect to the fibrations are all surjections.

Introduction

Our original intention was to describe effective descent morphisms in the category $\mathcal{C}$ of compact 0-dimensional topological spaces by combining the following well-known facts:

- A compact 0-dimensional space is nothing but a set equipped with a surjection into a Stone space (see Theorem 2.1 for the precise formulation).
- The effective descent morphisms in the categories of sets and of Stone spaces are just surjections.

It is still the main purpose of the paper, although it turned out that:

- Not all pullbacks exist in $\mathcal{C}$. Therefore the definition of an effective descent morphism p in $\mathcal{C}$ should include the requirement: *all pullbacks along p must exist* (see Definition 3.2).
- When p is surjective, that requirement holds if and only if p is a fibration in a suitable sense (see Definition 2.2), which is very different from what is happening in the situations studied by H. Herrlich [1], and which makes the descent problem much easier. In a somewhat different situation, this is made clear in [3].
- The surjectivity requirement does not create any problem since it is independently forced by the reflection of isomorphisms by the pullback functor along an effective descent morphism.

1. Partially supported by South African NRF.
2. Partially supported by FCT/Centro de Matemática da Universidade de Coimbra.
Received by the editors 2008-07-10 and, in revised form, 2008-11-12.
Published on 2008-11-15 in the Tholen Festschrift.
2000 Mathematics Subject Classification: 18B30, 18A25, 18A20.
Key words and phrases: comma categories, effective descent, effective $\mathbb{F}$-descent.
© George Janelidze[1] and Manuela Sobral[2], 2008. Permission to copy for private use granted.

- Therefore the problem of describing effective descent morphisms in $\mathcal{C}$ has an easy solution: Theorem 3.3 says that they are the same as the surjective fibrations.

- However, this suggests a new question, namely, what are the effective descent morphisms with respect to fibrations? Fortunately there is a complete answer again: they are all surjections (Theorem 3.1).

- In particular, even though the spaces we consider are not necessarily Hausdorff spaces, which prevents their convergence relations to be maps, our characterization of their effective descent morphisms avoids using the Reiterman-Tholen characterization of effective descent morphisms in the category of all topological spaces [4].

Accordingly, the paper is organized as follows:

Section 1 contains preliminary categorical observations with no topology involved. The ground category $\mathcal{C}$ there is constructed as a full subcategory in the comma category $(\mathcal{S} \downarrow U)$, where $U : \mathcal{X} \to \mathcal{S}$ is a pullback preserving functor between categories with pullbacks, using also a distinguished class $\mathbb{E}$ of morphisms in $\mathcal{S}$. This class is also used to define what we call fibrations in $\mathcal{C}$. The sufficient conditions for a morphism to be an effective descent morphism (globally or with respect to the class of fibrations) given in Section 1 will become also necessary in the topological context of Sections 2 and 3.

Section 2 begins by recalling relevant topological concepts, presents the category of compact 0-dimensional spaces as a special case of $\mathcal{C}$ above, introduces fibrations of 0-dimensional spaces accordingly, and ends by proving that a surjective morphism in $\mathcal{C}$ admits all pullbacks along morphisms with the same codomain if and only if it is a fibration.

The purpose of Section 3 is to formulate and prove the two main results, namely the above mentioned Theorems 3.1 and 3.3.

1. Categorical framework

We fix the following data: categories $\mathcal{S}$ and $\mathcal{X}$ with pullbacks, a pullback preserving functor $U : \mathcal{X} \to \mathcal{S}$ and a class $\mathbb{E}$ of morphisms in $\mathcal{S}$ that has the following properties:

- contains all isomorphisms;

- is pullback stable;

- is closed under composition;

- forms a *stack* (=coincides with its *localization*), which means that if



is a pullback diagram with w being an effective descent morphism, then $u \in \mathbb{E} \Rightarrow v \in \mathbb{E}$.

Let $\mathcal{C} = \mathcal{C}[\mathcal{X}, \mathcal{S}, U, \mathbb{E}]$ be the full subcategory in the comma category $(\mathcal{S} \downarrow U)$ with objects all triples $A = (A_1, e_A, A_0)$, in which $e_A : A_1 \to U(A_0)$ is in $\mathbb{E}$; accordingly, a morphism $A \to B$ in $\mathcal{C}$ is a pair $f = (f_1, f_0)$, in which $f_1 : A_1 \to B_1$ and $f_0 : A_0 \to B_0$ are morphisms in $\mathcal{S}$ and $\mathcal{X}$ respectively,

$$\begin{array}{ccc} A_1 & \xrightarrow{e_A} & U(A_0) \\ {\scriptstyle f_1}\downarrow & & \downarrow{\scriptstyle U(f_0)} \\ B_1 & \xrightarrow[e_B]{} & U(B_0) \end{array}$$

such that $U(f_0)e_A = e_B f_1$.

1.1. Definition. *A morphism $f : A \to B$ in $(\mathcal{S} \downarrow U)$ is said to be a fibration if the morphism*

$$< f_1, e_A >: A_1 \to B_1 \times_{U(B_0)} U(A_0)$$

is in $\mathbb{E}$.

1.2. Observation. *If $f : A \to B$ is a fibration, and B is in $\mathcal{C}$, then, since the class $\mathbb{E}$ is pullback stable, A also is in $\mathcal{C}$.*

1.3. Proposition. *Let*

(1.1)
$$\begin{array}{ccc} D & \xrightarrow{q} & A \\ {\scriptstyle g}\downarrow & & \downarrow{\scriptstyle f} \\ E & \xrightarrow[p]{} & B \end{array}$$

be a pullback diagram in $(\mathcal{S} \downarrow U)$ with $p : E \to B$ in $\mathcal{C}$. Then:

(a) If f is a fibration, then so is g.

(b) If g is a fibration, and p_1 is an effective descent morphism in $\mathcal{S}$, then f also is a fibration.

(c) If p is a fibration and A is in $\mathcal{C}$, then D is in $\mathcal{C}$.

(d) If $\mathbb{E}$ has the (weak left) cancellation property (e', $e \cdot e' \in \mathbb{E} \Rightarrow e \in \mathbb{E}$) and p_1 and $U(p_0)$ are in $\mathbb{E}$ and D is in $\mathcal{C}$, then A is in $\mathcal{C}$.

PROOF. Consider the diagram



in which:

- the enveloping cube represents the diagram (1.1);
- $e_E s = U(g_0)s'$ and $e_B t = U(f_0)t'$ are pullbacks;
- $d =< g_1, e_D >$, $a =< f_1, e_A >$, and $h = p_1 \times U(q_0)$ are the suitable induced morphisms.

Since the front square $U(p_0)U(g_0) = U(f_0)U(q_0)$ and the quadrilaterals $e_E s = U(g_0)s'$ and $e_B t = U(f_0)t'$ are pullbacks, so is the quadrilateral $p_1 s = th$. Next, since $p_1 g_1 = f_1 q_1$ and $p_1 s = th$ are pullbacks, so is $hd = aq_1$. This proves (a).

(b): Since p_1 is an effective descent morphism and $p_1 s = th$ is a pullback, h also is an effective descent morphism ([5], Theorem 3.1). Since $hd = aq_1$ is a pullback, this proves (b).

For (c) and (d), in order to use the same observations, let us "turn the diagram (1.1) around the diagonal connecting D and B", i.e. let us reformulate (c) and (d) as follows:

(c′) If f is a fibration and E is in $\mathcal{C}$, then D is in $\mathcal{C}$.

(d′) If f_1 and $U(f_0)$ are in $\mathbb{E}$ and D is in $\mathcal{C}$, then E is in $\mathcal{C}$.

Proof of (c′):

- Since f is a fibration, a is in $\mathbb{E}$.
- Since E is in $\mathcal{C}$ and $e_E s = U(g_0)s'$ is a pullback, s' is in $\mathbb{E}$.
- Since a and s' are in $\mathbb{E}$, so is e_D, i.e. D is in $\mathcal{C}$.

Proof of (d′):

- Since f_1 and $U(f_0)$ are in $\mathbb{E}$, so are g_1 and $U(g_0)$.
- Since $g_1, U(g_0)$ and e_D are in $\mathbb{E}$, the cancellation property of (d′) implies that e_E is in $\mathbb{E}$, as desired.

■

From Observation 1.2 and Proposition 1.3(a) we obtain:

1.4. Corollary. *The category $\mathcal{C}$ is closed in $(\mathcal{S} \downarrow U)$ under pullbacks along fibrations; that is, if (1.1) is a pullback diagram in $(\mathcal{S} \downarrow U)$ with f in $\mathcal{C}$ and p being a fibration in $\mathcal{C}$, then it is a pullback diagram in $\mathcal{C}$.*

When $\mathcal{S}$ has coequalizers of equivalence relations, all effective descent morphisms in $\mathcal{S}$ are regular epimorphisms. Using this fact it is easy to show that if $p : E \to B$ is a morphism in $(\mathcal{S} \downarrow U)$, for which p_0 and p_1 are effective descent morphisms in $\mathcal{X}$ and in $\mathcal{S}$ respectively, then p itself is an effective descent morphism. After that, using Proposition 1.3 and Corollary 1.4, we obtain:

1.5. Proposition. *If $\mathcal{S}$ has coequalizers of equivalence relations and $p : E \to B$ is a morphism in $\mathcal{C}$, for which p_0 and p_1 are effective descent morphisms in $\mathcal{X}$ and in $\mathcal{S}$ respectively, then*

(a) p is an effective $\mathbb{F}$-descent morphism in $\mathcal{C}$, where $\mathbb{F}$ is the class of all fibrations (in $\mathcal{C}$).

(b) if p is a fibration, then it is an effective descent morphism in $\mathcal{C}$.

2. The category of compact 0-dimensional spaces

For a topological space A, we shall write $Open(A)$ for the set of open subsets in A and $Clopen(A)$ for the set of those subsets in A that are *clopen*, i.e. closed and open at the same time. Let us recall the definitions of the following full subcategories of the category $\mathcal{T}op$ of topological spaces:

– $\mathcal{T}op_0$, the category of T_0-spaces; a space A is a T_0-space if, for every two distinct points a and a' in A, either there exists $U \in Open(A)$ with $a \in U$ and $a' \notin U$, or there exists $U \in Open(A)$ with $a' \in U$ and $a \notin U$. Note that $\mathcal{T}op_0$ is a reflective subcategory in $\mathcal{T}op$, with the reflection given by

$$A \mapsto A_0 = A/\sim, \text{ where } a \sim a' \Leftrightarrow \forall_{U \in Open(A)}(a \in U \Leftrightarrow a' \in U). \tag{2.1}$$

– 0-$\mathcal{D}im\mathcal{T}op$, the category of 0-dimensional spaces; a space is 0-dimensional if it has a basis of clopen subsets, i.e. if every open subset in it can be presented as a union of clopen subsets.

– The category of compact 0-dimensional spaces, which is the category of interest in this paper, will be simply denoted by $\mathcal{C}$; hence

$$\mathcal{C} = \mathcal{C}ompTop \cap 0\text{-}\mathcal{D}imTop$$

where $\mathcal{C}ompTop$ is the category of compact spaces.

– $\mathcal{S}tone$, the category of Stone spaces = spaces that occur as spectra of Boolean algebras = spaces that occur as limits of finite discrete spaces = compact Hausdorff 0-dimensional spaces = compact spaces A, such that for every two distinct points a and a' in A, there exists $U \in Clopen(A)$ with $a \in U$ and $a' \notin U$. The T_0-reflection (2.1) of course induces a reflection

$$\mathcal{C} \mapsto \mathcal{S}tone, A \mapsto A_0. \tag{2.2}$$

The following theorem is a reformulation of well-known results (see also Example 3.3 in [2] for the same result for arbitrary topological spaces, which, together with other similar results was mentioned already in [1]):

2.1. Theorem. *The category $\mathcal{C}$ of compact 0-dimensional spaces is equivalent to the category $\mathcal{C}[\mathcal{X}, \mathcal{S}, U, \mathbb{E}]$ (see Section 1), for $\mathcal{X} = \mathcal{S}tone$, $\mathcal{S} = \mathcal{S}et$, $U : \mathcal{S}tone \to \mathcal{S}et$, U being the usual forgetful functor into the category of sets and $\mathbb{E}$ being the class of all surjective maps. Under this equivalence a space A corresponds to the triple (A_1, e_A, A_0), in which A_1 is the underlying set of A, A_0 is the T_0-reflection of A, and $e_A : A_1 \to U(A_0)$ is the canonical map (and we write again $A = (A_1, e_A, A_0)$). Conversely, the space corresponding to a triple (A_1, e_A, A_0) has A_1 as its underlying set and the inverse e_A-images of open sets in A_0 as its open subsets.*

According to this theorem and Definition 1.1, we introduce:

2.2. Definition. *A morphism $f : A \to B$ in $\mathcal{C}$ is said to be a fibration if so is the corresponding morphism in $\mathcal{C}[\mathcal{X}, \mathcal{S}, U, \mathbb{E}]$ of Theorem 2.1, i.e. if for every a in A and b in B with $f(a) \sim b$ there exists a' in A with $a' \sim a$ and $f(a') = b$.*

After that Proposition 1.3 helps to prove:

2.3. Theorem. *Let $p : E \to B$ be a morphism in $\mathcal{C}$. If p is surjective, then the following conditions are equivalent:*

(a) every morphism $f : A \to B$ in $\mathcal{C}$ admits pullback along p;

(b) p is a fibration.

Proof. (a)⇒(b): Suppose p is not a fibration. This means that there are e in E and b in B with

$$p(e) \sim b \text{ and } (x \in p^{-1}(b) \Rightarrow \exists_{U_x \in Clopen(E)}(x \in U_x \text{ and } e \notin U_x)). \tag{2.3}$$

We choose U_x as in (2.3) for each x in $p^{-1}(b)$, and consider two cases:

Case 1. There exists a finite subset Y in X, for which

$$p^{-1}(b) \subseteq \bigcup_{x \in Y} U_x.$$

Case 2. There is no such Y.

In *Case 1* we take

$$V = \bigcap_{x \in Y} (E \setminus U_x),$$

and observe that since Y is finite, V is clopen; and of course V contains e and has empty intersection with $p^{-1}(b)$. After that we take

$$A = \{n^{-1} | n = 1, 2, 3, \cdots\} \cup \{0\}$$

with the topology induced from the real line, and define $f : A \to B$ by $f(n^{-1}) = b$ and $f(0) = p(e)$. Suppose the pullback of p and f does exist, and let us write it as the diagram (1.1). Using the universal property of this pullback with respect to maps from a one-point space, we easily conclude that it is preserved by the forgetful functor into the category of sets. In particular, since V contains e and has empty intersection with $p^{-1}(b)$, we have

$$q(g^{-1}(E \setminus V)) = \{n^{-1} | n = 1, 2, 3, \cdots\}.$$

This is a contradiction since $g^{-1}(E \setminus V)$ being clopen in D must be compact in it, while $\{n^{-1} | n = 1, 2, 3, \cdots\}$ is not compact in A.

In *Case* 2 we take $A = \{a\}$ to be a one-point space, and define $f : A \to B$ by $f(a) = b$. Then, using (1.1) as above, we observe that $g(D) = p^{-1}(b)$ – which is again a contradiction because now $p^{-1}(b)$ is not compact.

That is, whenever p is not a fibration, there exists a morphism $f : A \to B$ in $\mathcal{C}$ that has no pullback along p.

(b)⇒(a) follows from Corollary 1.4 and Theorem 2.1. ∎

3. $\mathbb{F}$-Descent and global descent

Let $\mathcal{C}$ be as in Section 2.

3.1. Theorem. *The following conditions on a morphism $p : E \to B$ in $\mathcal{C}$ are equivalent:*

(a) p is an effective $\mathbb{F}$-descent morphism in $\mathcal{C}$;

(b) p is a surjective map.

Proof. (a)⇒(b): Suppose p is not surjective, and choose $b \in B \setminus p(E)$. Let A be the equivalence class of b with respect to the equivalence relation $\sim$ (see (2.1)). We take

$$A' = (A \setminus \{b\}) \cup \{b\} \times \{1, 2\}$$

equipped with the indiscrete topology, and define $\alpha : A' \to A$ by $\alpha(a) = a$, for $a \in A$, and $\alpha(b, 1) = b = \alpha(b, 2)$; then α becomes a morphism $(A', \alpha f) \to (A, f)$, where $f : A \to B$ is

the inclusion map, in the category $\mathbb{F}(B)$ of fibrations over B (in $\mathcal{C}$). Since the image of this morphism under the pullback functor $p^* : F(B) \to F(E)$ is an isomorphism, p cannot be an effective $\mathbb{F}$-descent morphism in $\mathcal{C}$.

(b)⇒(a): Let $(p_1, p_0) : (E_1, e_E, E_0) \to (B_1, e_B, B_0)$ be the morphism in $\mathcal{C}[\mathcal{X}, \mathcal{S}, U, \mathbb{E}]$ corresponding to p under the category equivalence of Theorem 2.1, where $X = \mathcal{S}tone, \mathcal{S} = \mathcal{S}et$, $U : \mathcal{S}tone \to \mathcal{S}et$ being the usual forgetful functor into the category of sets, and $\mathbb{E}$ being the class of all surjective maps. Then p_1 is surjective and this makes p_0 surjective too. Since in both $\mathcal{S}tone$ and $\mathcal{S}et$ surjections are effective descent morphisms, this makes p an effective $\mathbb{F}$-descent morphism by Proposition 1.5(a). ∎

Since $\mathcal{C}$ does not admit some pullbacks, we define effective (global-)descent morphisms in $\mathcal{C}$ as follows:

3.2. Definition. *A morphism $p : E \to B$ in $\mathcal{C}$ is said to be an effective descent morphism if every morphism $f : A \to B$ in $\mathcal{C}$ admits pullback along p, and the pullback functor*

$$p* : (\mathcal{C} \downarrow B) \to (\mathcal{C} \downarrow E)$$

is monadic.

3.3. Theorem. *The following conditions on a morphism $p : E \to B$ in $\mathcal{C}$ are equivalent:*
(a) p is an effective descent morphism;
(b) p is a surjective fibration.

Proof. (a)⇒(b): Surjectivity can be proved in the same way as in the proof of Theorem 3.1 (or even much simpler by considering the empty and one-point space instead of A' and A there). The fact that p must be a fibration follows from the implication (a)⇒(b) of Theorem 2.3.

(b)⇒(a) can be deduced from Proposition 1.5(b) and Theorem 2.1 with the same arguments as in the proof of Theorem 3.1(b)⇒(a). ∎

References

[1] H. Herrlich, *Topological functors*, General Topology and Appl. 4 (1974) 125-142.

[2] G. Janelidze, *Light morphisms for generalized T_0-reflections*, submitted.

[3] G. Janelidze and M. Sobral, *Finite preorders and topological descent I*, J. Pure Appl. Algebra 175(1-3)(2001) 187-205.

[4] J. Reiterman and W. Tholen, *Effective descent maps of topological spaces*, Topology Appl. 57 (1994) 53-69.

[5] M. Sobral and W. Tholen, *Effective descent morphisms and effective equivalence relations*, in: Category Theory 1991, CMS Conference Proceedings (Amer. Math. Soc., Providence, R. I. 1992), 421-432.

Dept. Math. and Appl. Math.
University of Cape Town
Rondebosh 7701, Cape Town
South Africa

Departamento de Matemática
Universidade de Coimbra
Ap. 3008, 3001-454 Coimbra
Portugal

Email: janelidg@maths.uct.ac.za
sobral@mat.uc.pt

This article may be accessed at http://www.tac.mta.ca/tac/ or by anonymous ftp at ftp://ftp.tac.mta.ca/pub/tac/html/volumes/21/10/21-10.{dvi,ps,pdf}

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.
Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.
Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at `http://www.tac.mta.ca/tac/` and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION. Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to `tac@mta.ca` including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, `rrosebrugh@mta.ca`.

INFORMATION FOR AUTHORS. The typesetting language of the journal is TEX, and LATEX2e strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at `http://www.tac.mta.ca/tac/`.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: `rrosebrugh@mta.ca`

TEXNICAL EDITOR. Michael Barr, McGill University: `barr@math.mcgill.ca`

ASSISTANT TEX EDITOR. Gavin Seal, McGill University: `gavin_seal@fastmail.fm`

TRANSMITTING EDITORS.
Richard Blute, Université d' Ottawa: `rblute@uottawa.ca`
Lawrence Breen, Université de Paris 13: `breen@math.univ-paris13.fr`
Ronald Brown, University of North Wales: `ronnie.profbrown (at) btinternet.com`
Aurelio Carboni, Università dell Insubria: `aurelio.carboni@uninsubria.it`
Valeria de Paiva, Cuill Inc.: `valeria@cuill.com`
Ezra Getzler, Northwestern University: `getzler(at)northwestern(dot)edu`
Martin Hyland, University of Cambridge: `M.Hyland@dpmms.cam.ac.uk`
P. T. Johnstone, University of Cambridge: `ptj@dpmms.cam.ac.uk`
Anders Kock, University of Aarhus: `kock@imf.au.dk`
Stephen Lack, University of Western Sydney: `s.lack@uws.edu.au`
F. William Lawvere, State University of New York at Buffalo: `wlawvere@acsu.buffalo.edu`
Jean-Louis Loday, Université de Strasbourg: `loday@math.u-strasbg.fr`
Ieke Moerdijk, University of Utrecht: `moerdijk@math.uu.nl`
Susan Niefield, Union College: `niefiels@union.edu`
Robert Paré, Dalhousie University: `pare@mathstat.dal.ca`
Jiri Rosicky, Masaryk University: `rosicky@math.muni.cz`
Brooke Shipley, University of Illinois at Chicago: `bshipley@math.uic.edu`
James Stasheff, University of North Carolina: `jds@math.unc.edu`
Ross Street, Macquarie University: `street@math.mq.edu.au`
Walter Tholen, York University: `tholen@mathstat.yorku.ca`
Myles Tierney, Rutgers University: `tierney@math.rutgers.edu`
Robert F. C. Walters, University of Insubria: `robert.walters@uninsubria.it`
R. J. Wood, Dalhousie University: `rjwood@mathstat.dal.ca`