Theory and Applications of Categories, Vol. 21, No. 12, 2009, pp. 210–230.

RELATIVE INJECTIVITY AS COCOMPLETENESS FOR A CLASS OF DISTRIBUTORS

Dedicated to Walter Tholen on the occasion of his sixtieth birthday

MARIA MANUEL CLEMENTINO AND DIRK HOFMANN

Abstract. Notions and techniques of enriched category theory can be used to study topological structures, like metric spaces, topological spaces and approach spaces, in the context of topological theories. Recently in [D. Hofmann, Injective spaces via adjunction, arXiv:math.CT/0804.0326] the construction of a Yoneda embedding allowed to identify injectivity of spaces as cocompleteness and to show monadicity of the category of injective spaces and left adjoints over Set. In this paper we generalise these results, studying cocompleteness with respect to a given class of distributors. We show in particular that the description of several semantic domains presented in [M. Escardó and B. Flagg, Semantic domains, injective spaces and monads, Electronic Notes in Theoretical Computer Science 20 (1999)] can be translated into the V-enriched setting.

Introduction

This work continues the research line of previous papers, aiming to use categorical tools in the study of topological structures. Indeed, the perspective proposed in [3, 7] of looking at topological structures as (Eilenberg-Moore) lax algebras and, simultaneously, as a monad enrichment of V-enriched categories, has shown to be very effective in the study of special morphisms – like effective descent and exponentiable ones – at a first step [4, 5], and recently in the study of (Lawvere/Cauchy-)completeness and injectivity [6, 12, 11]. The results we present here complement this study of injectivity. More precisely, in the spirit of Kelly-Schmitt [13] we generalise the results of [11], showing that injectivity and cocompleteness – when considered relative to a class of distributors – still coincide. Suitable choices of this class of distributors allow us to recover, in the V-enriched setting, results on injectivity of Escardó-Flagg [8].

The starting point of our study of injectivity is the notion of distributor (or bimodule, or profunctor), which allowed the study of weighted colimit, presheaf category, and the

The authors acknowledge partial financial assistance by Centro de Matemática da Universidade de Coimbra/FCT and Unidade de Investigação e Desenvolvimento Matemática e Aplicações da Universidade de Aveiro/FCT.

Received by the editors 2008-07-14 and, in revised form, 2009-02-12.

Published on 2009-02-15 in the Tholen Festschrift.

2000 Mathematics Subject Classification: 18A05, 18D15, 18D20, 18B35, 18C15, 54B30, 54A20.

Key words and phrases: Quantale, V-category, monad, topological theory, distributor, Yoneda lemma, weighted colimit.

© Maria Manuel Clementino and Dirk Hofmann, 2009. Permission to copy for private use granted.

Yoneda embedding. It was then a natural step to 'relativize' these ingredients and to consider *cocompleteness with respect to a class of distributors* Φ. Namely, we introduce the notion of Φ-cocomplete category, we construct the Φ-presheaf category, and we prove that Φ-cocompleteness is equivalent to the existence of a left adjoint of the Yoneda embedding into the Φ-presheaf category. Furthermore, the class Φ determines a class of embeddings so that the injective $\mathcal{T}$-categories with respect to this class are precisely the Φ-cocomplete categories. This result links our work with [8], where the authors study systematically semantic domains and injectivity characterisations with the help of Kock-Zöberlein monads.

1. The Setting

The topological structures we study throughout are those which are describable as lax (Eilenberg-Moore) algebras, or as $(\mathbb{T}, \mathsf{V})$-enriched categories, for a suitable (thin) category V and a suitable monad $\mathbb{T}$ in Set, with a lax extension to V-Rel. Recall from [1] that topological spaces viewed as convergence structures provide the prime example of such a situation, where $\mathbb{T} = \mathbb{U}$ is the ultrafilter monad and $\mathsf{V} = \mathsf{2}$ the two-element Boolean algebra. Our study could be based on the setting described by Clementino-Tholen in [7], but we chose to use the slightly different approach of Hofmann [10] – the so-called topological theories –, which encodes the lax extension of $\mathbb{T}$ in a $\mathbb{T}$-algebra structure on V.

Throughout this paper we consider a (strict) *topological theory* as introduced in [10]. Such a theory $\mathcal{T} = (\mathbb{T}, \mathsf{V}, \xi)$ consists of:

(1). a cocomplete monoidal closed ordered set V, with tensor $\otimes$ and unit k, and we denote by hom the right adjoint to $\otimes$ (that is, the internal hom),

(2). a Set-monad $\mathbb{T} = (T, e, m)$, where T and m satisfy (BC); that is, T sends pullbacks to weak pullbacks and each naturality square of m is a weak pullback, and

(3). a $\mathbb{T}$-algebra structure $\xi : T\mathsf{V} \longrightarrow \mathsf{V}$ on V such that:

(a) $\otimes : \mathsf{V} \times \mathsf{V} \longrightarrow \mathsf{V}$ and $k : 1 \longrightarrow \mathsf{V}$, $* \longmapsto k$, are $\mathbb{T}$-algebra homomorphisms making (V, ξ) a monoid in $\mathsf{Set}^{\mathbb{T}}$; that is, the following diagrams

$$\begin{array}{ccc} T1 & \xrightarrow{Tk} & T\mathsf{V} \\ {\scriptstyle !}\downarrow & & \downarrow{\scriptstyle \xi} \\ 1 & \xrightarrow[k]{} & \mathsf{V} \end{array} \qquad \begin{array}{ccc} T(\mathsf{V}\times\mathsf{V}) & \xrightarrow{T(\otimes)} & T\mathsf{V} \\ {\scriptstyle \langle \xi\cdot T\pi_1, \xi\cdot T\pi_2\rangle}\downarrow & & \downarrow{\scriptstyle \xi} \\ \mathsf{V}\times\mathsf{V} & \xrightarrow[\otimes]{} & \mathsf{V} \end{array}$$

are commutative;

(b) For each set X, $\xi_X : \mathsf{V}^X \longrightarrow \mathsf{V}^{TX}$, $(X \xrightarrow{\varphi} \mathsf{V}) \longmapsto (TX \xrightarrow{T\varphi} T\mathsf{V} \xrightarrow{\xi} \mathsf{V})$, defines a natural transformation $(\xi_X)_X : P \longrightarrow PT : \mathsf{Set} \longrightarrow \mathsf{Ord}$.

Here $P : \mathsf{Set} \longrightarrow \mathsf{Ord}$ is the V-powerset functor defined as follows. We put $PX = \mathsf{V}^X$ with the pointwise order. Each map $f : X \longrightarrow Y$ defines a monotone map $\mathsf{V}^f : \mathsf{V}^Y \longrightarrow \mathsf{V}^X$, $\varphi \longmapsto \varphi \cdot f$. Since V^f preserves all infima and all suprema, it has a left adjoint Pf. Explicitly, for $\varphi \in \mathsf{V}^X$ we have $Pf(\varphi)(y) = \bigvee\{\varphi(x) \mid x \in X,\ f(x) = y\}$.

1.1. Examples. Throughout this paper we will keep in mind the following topological theories:

(1). The identity theory $\mathcal{I} = (\mathbb{1}, \mathsf{V}, 1_{\mathsf{V}})$, for each quantale V, where $\mathbb{1} = (\mathrm{Id}, 1, 1)$ denotes the identity monad.

(2). $\mathcal{U}_2 = (\mathbb{U}, 2, \xi_2)$, where $\mathbb{U} = (U, e, m)$ denotes the ultrafilter monad and ξ_2 is essentially the identity map.

(3). $\mathcal{U}_{\mathsf{P}_+} = (\mathbb{U}, \mathsf{P}_+, \xi_{\mathsf{P}_+})$ where $\mathsf{P}_+ = ([0, \infty]^{\mathrm{op}}, +, 0)$ and

$$\xi_{\mathsf{P}_+} : U\mathsf{P}_+ \longrightarrow \mathsf{P}_+, \quad \mathfrak{x} \longmapsto \inf\{v \in \mathsf{P}_+ \mid [0, v] \in \mathfrak{x}\}.$$

(4). The word theory $(\mathbb{L}, \mathsf{V}, \xi_\otimes)$, for each quantale V, where $\mathbb{L} = (L, e, m)$ is the word monad and

$$\begin{gathered} \xi_\otimes : L\mathsf{V} \longrightarrow \mathsf{V}. \\ (v_1, \ldots, v_n) \longmapsto v_1 \otimes \ldots \otimes v_n \\ () \longmapsto k \end{gathered}$$

As we mentioned at the beginning of this section, every theory $\mathcal{T} = (\mathbb{T}, \mathsf{V}, \xi)$ encompasses several interesting ingredients.

I. *The category* V-Rel, with *sets* as objects and V-*relations* (also called V-*matrices*, see [2]) $r : X \times Y \longrightarrow \mathsf{V}$ as morphisms, is a locally complete locally ordered bicategory. We use the usual notation for relations, denoting the V-relation $r : X \times Y \longrightarrow \mathsf{V}$ by $r : X \relbar\!\!\!\rightarrow Y$. Since every map $f : X \longrightarrow Y$ can be thought of as a V-relation $f : X \times Y \longrightarrow \mathsf{V}$ through its graph, there is an injective on objects and faithful functor

$$\mathsf{Set} \longrightarrow \mathsf{V}\text{-}\mathsf{Rel},$$

unless V is degenerate (that is, k is the bottom element). Moreover, V-Rel has an involution

$$(-)^\circ : \mathsf{V}\text{-}\mathsf{Rel} \longrightarrow \mathsf{V}\text{-}\mathsf{Rel},$$

assigning to $r : X \relbar\!\!\!\rightarrow Y$ the V-relation $r^\circ : Y \relbar\!\!\!\rightarrow X$, with $r^\circ(y, x) := r(x, y)$. For each V-relation $r : X \relbar\!\!\!\rightarrow Y$, the maps

$$(-) \cdot r : \mathsf{V}\text{-}\mathsf{Rel}(Y, Z) \longrightarrow \mathsf{V}\text{-}\mathsf{Rel}(X, Z) \text{ and } r \cdot (-) : \mathsf{V}\text{-}\mathsf{Rel}(Z, X) \longrightarrow \mathsf{V}\text{-}\mathsf{Rel}(Z, Y)$$

preserve suprema; hence they have right adjoints, the so-called *extensions* and *liftings*, respectively,

$$(-) \bullet\!\!- r : \mathsf{V\text{-}Rel}(X, Z) \longrightarrow \mathsf{V\text{-}Rel}(Y, Z) \text{ and } r \multimap\!\!\bullet\ (-) : \mathsf{V\text{-}Rel}(Z, Y) \longrightarrow \mathsf{V\text{-}Rel}(Z, X) :$$

II. *The* Set*-functor* T *extends to a 2-functor* $T_\xi : \mathsf{V\text{-}Rel} \longrightarrow \mathsf{V\text{-}Rel}$. To each V-relation $r : X \times Y \longrightarrow \mathsf{V}$, T_ξ assigns a V-relation $T_\xi r : TX \times TY \longrightarrow \mathsf{V}$, such that, for every (order-preserving) map $s : TX \times TY \longrightarrow \mathsf{V}$,

$$\xi \cdot Tr \leq s \cdot \langle T\pi_1, T\pi_2 \rangle \ \Leftrightarrow\ T_\xi r \leq s :$$



In other words, regarding TX, TY and $TX \times TY$ as discrete ordered sets, $T_\xi r$ is the left Kan extension in Ord of $\xi \cdot Tr$ along $\langle T\pi_1, T\pi_2 \rangle$. Hence, for $\mathfrak{x} \in TX$ and $\mathfrak{y} \in TY$,

$$T_\xi r(\mathfrak{x}, \mathfrak{y}) = \bigvee \left\{ \xi \cdot Tr(\mathfrak{w}) \;\middle|\; \mathfrak{w} \in T(X \times Y), T\pi_1(\mathfrak{w}) = \mathfrak{x}, T\pi_2(\mathfrak{w}) = \mathfrak{y} \right\}.$$

This 2-functor T_ξ preserves the involution in the sense that $T_\xi(r^\circ) = T_\xi(r)^\circ$ (and we write $T_\xi r^\circ$) for each V-relation $r : X \relbar\!\!\!\rightarrow Y$, m becomes a natural transformation $m : T_\xi T_\xi \longrightarrow T_\xi$ and e an op-lax natural transformation $e : \mathrm{Id} \longrightarrow T_\xi$, that is, $e_Y \cdot r \leq T_\xi r \cdot e_X$ for all $r : X \relbar\!\!\!\rightarrow Y$ in V-Rel.

III. A V-relation of the form $\alpha : TX \relbar\!\!\!\rightarrow Y$, called a $\mathfrak{T}$*-relation* and denoted by $\alpha : X \relbar\!\!\!\rightharpoonup Y$, will play an important role here. Given two $\mathfrak{T}$-relations $\alpha : X \relbar\!\!\!\rightharpoonup Y$ and $\beta : Y \relbar\!\!\!\rightharpoonup Z$, their *Kleisli convolution* $\beta \circ \alpha : X \relbar\!\!\!\rightharpoonup Z$ is defined as

$$\beta \circ \alpha = \beta \cdot T_\xi \alpha \cdot m_X^\circ.$$

This operation is associative and has the $\mathfrak{T}$-relation $e_X^\circ : X \relbar\!\!\!\rightharpoonup X$ as a lax identity:

$$a \circ e_X^\circ = a \text{ and } e_Y^\circ \circ a \geq a,$$

for any $a : X \relbar\!\!\!\rightharpoonup Y$.

IV. $\mathcal{T}$-relations satisfying the usual unit and associativity categorical rules define $\mathcal{T}$-categories: a $\mathcal{T}$-*category* is a pair (X, a) consisting of a set X and a $\mathcal{T}$-relation $a : X \nrightarrow X$ on X such that

$$e_X^\circ \leq a \qquad \text{and} \qquad a \circ a \leq a.$$

Expressed elementwise, these conditions become

$$k \leq a(e_X(x), x) \qquad \text{and} \qquad T_\xi a(\mathfrak{X}, \mathfrak{x}) \otimes a(\mathfrak{x}, x) \leq a(m_X(\mathfrak{X}), x)$$

for all $\mathfrak{X} \in TTX$, $\mathfrak{x} \in TX$ and $x \in X$. A function $f : X \longrightarrow Y$ between $\mathcal{T}$-categories (X, a) and (Y, b) is a $\mathcal{T}$-*functor* if $f \cdot a \leq b \cdot Tf$, which in pointwise notation reads as

$$a(\mathfrak{x}, x) \leq b(Tf(\mathfrak{x}), f(x))$$

for all $\mathfrak{x} \in TX$, $x \in X$. The category of $\mathcal{T}$-categories and $\mathcal{T}$-functors is denoted by

$$\mathcal{T}\text{-}\mathsf{Cat}.$$

V. In particular, the internal hom in V, combined with the $\mathbb{T}$-algebra structure ξ, induces a $\mathcal{T}$-category structure in V,

$$\hom_\xi : T\mathsf{V} \times \mathsf{V} \longrightarrow \mathsf{V},\ (\mathfrak{v}, v) \longmapsto \hom(\xi(\mathfrak{v}), v).$$

VI. *The forgetful functor* $\mathrm{O} : \mathcal{T}\text{-}\mathsf{Cat} \longrightarrow \mathsf{Set},\ (X, a) \longmapsto X$, *is topological*, hence it has a left and a right adjoint. In particular, the free $\mathcal{T}$-category on X is given by (X, e_X°).

VII. A V-relation $\varphi : X \nrightarrow Y$ between $\mathcal{T}$-categories $X = (X, a)$ and $Y = (Y, b)$ is a $\mathcal{T}$-*distributor*, denoted as $\varphi : X \multimap Y$, if $\varphi \circ a \leq \varphi$ and $b \circ \varphi \leq \varphi$. Note that we always have $\varphi \circ a \geq \varphi$ and $b \circ \varphi \geq \varphi$, so that the $\mathcal{T}$-distributor conditions above are in fact equalities. $\mathcal{T}$-categories and $\mathcal{T}$-distributors form a 2-category, denoted by

$$\mathcal{T}\text{-}\mathsf{Mod},$$

with Kleisli convolution as composition and with the 2-categorical structure inherited from V-Rel.

VIII. *Each* $\mathcal{T}$-*functor* $f : (X, a) \longrightarrow (Y, b)$ *induces an adjunction*

$$f_* \dashv f^*$$

in $\mathcal{T}$-Mod, with $f_* : X \nrightarrow Y$ and $f^* : Y \nrightarrow X$ defined as $f_* = b \cdot Tf$ and $f^* = f^\circ \cdot b$ respectively. In fact, these assignments are functorial and therefore define two functors:

$$\begin{array}{rcl} (-)_* : \mathcal{T}\text{-}\mathsf{Cat}^{\mathrm{co}} & \longrightarrow & \mathcal{T}\text{-}\mathsf{Mod} \\ X & \longmapsto & X_* = X \\ f & \longmapsto & f_* = b \cdot Tf \end{array} \qquad \text{and} \qquad \begin{array}{rcl} (-)^* : \mathcal{T}\text{-}\mathsf{Cat}^{\mathrm{op}} & \longrightarrow & \mathcal{T}\text{-}\mathsf{Mod}, \\ X & \longmapsto & X^* = X \\ f & \longmapsto & f^* = f^\circ \cdot b \end{array}$$

A $\mathcal{T}$-functor $f : X \longrightarrow Y$ is called *fully faithful* if $f^* \circ f_* = 1_X^*$, while it is called *dense* if $f_* \circ f^* = 1_Y^*$. Note that f is fully faithful if and only if, for all $\mathfrak{x} \in TX$ and $x \in X$, $a(\mathfrak{x}, x) = b(Tf(\mathfrak{x}), f(x))$.

IX. For a $\mathcal{T}$-distributor $\alpha : X \multimap Y$, *the composition function* $- \circ \alpha$ *has a right adjoint*

$$(-) \circ \alpha \dashv (-) \circ\!\!- \alpha$$

where, for a given $\mathcal{T}$-distributor $\gamma : X \multimap Z$, the extension $\gamma \circ\!\!- \alpha : Y \multimap Z$ is constructed in V-Rel as the extension $\gamma \circ\!\!- \alpha = \gamma \bullet\!\!- (T_\xi \alpha \cdot m_X^\circ)$.



The following rules are easily checked.

1.2. Lemma.

(1). If α *is a right adjoint, then* $\alpha \circ (\varphi \circ\!\!- \psi) = (\alpha \circ \varphi) \circ\!\!- \psi$.

(2). If $\gamma \dashv \delta$, *then* $(\alpha \circ\!\!- \beta) \circ \gamma = \alpha \circ\!\!- (\delta \circ \beta)$.

(3). If $\gamma \dashv \delta$, *then* $(\alpha \circ \gamma) \circ\!\!- \beta = \alpha \circ\!\!- (\beta \circ \delta)$.

X. It is also important the interplay of several functors relating these structures: *Eilenberg-Moore algebras*, $\mathcal{T}$*-categories* and V*-categories*. The inclusion functor $\mathsf{Set}^{\mathbb{T}} \hookrightarrow \mathcal{T}$-$\mathsf{Cat}$, given by regarding the structure map $\alpha : TX \longrightarrow X$ of an Eilenberg-Moore algebra (X, α) as a $\mathcal{T}$-relation $\alpha : X \nrightarrow X$, has a left adjoint, constructed *à la Čech-Stone compactification* in [3].



We denote by $|X|$ the free Eilenberg-Moore algebra (TX, m_X) considered as a $\mathcal{T}$-category.

Making use of the identity $e : \mathrm{Id} \longrightarrow T$ of the monad, to each $\mathcal{T}$-category $X = (X, a)$ we assign a V-category structure on X, $a \cdot e_X : X \nrightarrow X$. This correspondence defines a functor $\mathrm{S} : \mathcal{T}$-$\mathsf{Cat} \longrightarrow \mathsf{V}$-$\mathsf{Cat}$, which has also a left adjoint $\mathrm{A} : \mathsf{V}$-$\mathsf{Cat} \longrightarrow \mathcal{T}$-$\mathsf{Cat}$, with $\mathrm{A}(X, a) := (X, e_X^\circ \cdot T_\xi r)$.



Furthermore, making now use of the multiplication $m : T^2 \longrightarrow T$ of the monad, one can define a functor



which sends a $\mathcal{T}$-category (X, a) to the V-category $(TX, T^{\circ}_{\xi} a \cdot m^{\circ}_X)$.

We can now define the process of *dualizing a $\mathcal{T}$-category* as the composition of the following functors

$$\begin{array}{ccc} \mathcal{T}\text{-}\mathsf{Cat} & \xrightarrow{(\ )^{\mathrm{op}}} & \mathcal{T}\text{-}\mathsf{Cat} \\ {\scriptstyle M}\downarrow & & \uparrow{\scriptstyle A} \\ \mathsf{V}\text{-}\mathsf{Cat} & \xrightarrow[(\ )^{\mathrm{op}}]{} & \mathsf{V}\text{-}\mathsf{Cat} \end{array}$$

that is, the *dual of a $\mathcal{T}$-category* (X, a) is defined as

$$X^{\mathrm{op}} = \mathrm{A}(\mathrm{M}(X)^{\mathrm{op}}),$$

which is a structure on TX. If $\mathbb{T}$ is the identity monad, then X^{op} is indeed the dual V-category of X.

XI. The tensor product on V can be transported to $\mathcal{T}$-Cat by putting

$$(X, a) \otimes (Y, b) = (X \times Y, c),$$

with

$$c(\mathfrak{w}, (x, y)) = a(T\pi_1(\mathfrak{w}), x) \otimes b(T\pi_2(\mathfrak{w}), y),$$

where $\mathfrak{w} \in T(X \times Y)$, $x \in X$, $y \in Y$. The $\mathcal{T}$-category $E = (1, k)$ is a $\otimes$-neutral object, where 1 is a singleton set and $k : T1 \times 1 \longrightarrow \mathsf{V}$ the constant relation with value $k \in \mathsf{V}$. For each set X, the functor $|X| \otimes (-) : \mathcal{T}\text{-}\mathsf{Cat} \longrightarrow \mathcal{T}\text{-}\mathsf{Cat}$ has a right adjoint $(-)^{|X|} : \mathcal{T}\text{-}\mathsf{Cat} \longrightarrow \mathcal{T}\text{-}\mathsf{Cat}$. Explicitly, the structure $[\![-, -]\!]$ on $\mathsf{V}^{|X|}$ is given by the formula

$$[\![\mathfrak{p}, \psi]\!] = \bigwedge_{\mathfrak{q} \in T(|X| \times \mathsf{V}^{|X|}) \atop \mathfrak{q} \longmapsto \mathfrak{p}} \hom(\xi \cdot T\,\mathrm{ev}(\mathfrak{q}), \psi(m_X \cdot T\pi_1(\mathfrak{q}))),$$

for each $\mathfrak{p} \in T\mathsf{V}^{|X|}$ and $\psi \in \mathsf{V}^{|X|}$.

1.3. Theorem. [6] *For $\mathcal{T}$-categories (X, a) and (Y, b), and a $\mathcal{T}$-relation $\psi : X \nrightarrow Y$, the following assertions are equivalent.*

(i). $\psi : (X, a) \multimap (Y, b)$ is a $\mathcal{T}$-distributor.

(ii). Both $\psi : |X| \otimes Y \longrightarrow \mathsf{V}$ and $\psi : X^{\mathrm{op}} \otimes Y \longrightarrow \mathsf{V}$ are $\mathcal{T}$-functors.

XII. Hence, *each $\mathcal{T}$-distributor $\varphi : X \multimap Y$ provides a $\mathcal{T}$-functor*

$$\ulcorner \varphi \urcorner : Y \longrightarrow \mathsf{V}^{|X|}$$

which factors through the embedding $\mathcal{P}X \hookrightarrow \mathsf{V}^{|X|}$, where

$$\mathcal{P}X = \{\psi \in \mathsf{V}^{|X|} \mid \psi : X \multimap (1, e^{\circ}_1)\}$$

is the $\mathcal{T}$-category of *contravariant presheafs on* X:



In particular, for each $\mathcal{T}$-category $X = (X, a)$, the V-relation $a : TX \times X \longrightarrow \mathsf{V}$ is a $\mathcal{T}$-distributor $a : X \multimap X$, and therefore we have the *Yoneda functor*

$$y_X = \ulcorner a \urcorner : X \longrightarrow \mathcal{P}X.$$

1.4. Theorem. [11] *Let* $\psi : X \multimap Z$ *and* $\varphi : X \multimap Y$ *be* $\mathcal{T}$*-distributors. Then, for all* $\mathfrak{z} \in TZ$ *and* $y \in Y$,

$$[\![T \ulcorner \psi \urcorner(\mathfrak{z}), \ulcorner \varphi \urcorner(y)]\!] = (\varphi \circ\!\!- \psi)(\mathfrak{z}, y).$$

1.5. Corollary. [11] *For each* $\varphi \in \hat{X}$ *and each* $\mathfrak{x} \in TX$, $\varphi(\mathfrak{x}) = [\![T\, y_X(\mathfrak{x}), \varphi]\!]$, *that is,* $(y_X)_* : X \multimap \hat{X}$ *is given by the evaluation map* $\mathrm{ev} : TX \times \hat{X} \longrightarrow \mathsf{V}$. *As a consequence,* $y_X : X \longrightarrow \hat{X}$ *is fully faithful.*

XIII. Transporting the order-structure on hom-sets from $\mathcal{T}$-Mod to $\mathcal{T}$-Cat via the functor $(-)^* : \mathcal{T}\text{-}\mathsf{Cat}^{\mathrm{op}} \longrightarrow \mathcal{T}\text{-}\mathsf{Mod}$, $\mathcal{T}$-Cat *becomes a 2-category.* That is, for $\mathcal{T}$-functors $f, g : X \longrightarrow Y$ we define

$$f \leq g \text{ in } \mathcal{T}\text{-}\mathsf{Cat} \;:\Leftrightarrow\; f^* \leq g^* \text{ in } \mathcal{T}\text{-}\mathsf{Mod} \;\Leftrightarrow\; g_* \leq f_* \text{ in } \mathcal{T}\text{-}\mathsf{Mod}.$$

We call $f, g : X \longrightarrow Y$ *equivalent*, and write $f \cong g$, if $f \leq g$ and $g \leq f$. Hence, $f \cong g$ if and only if $f^* = g^*$ if and only if $f_* = g_*$. A $\mathcal{T}$-category X is called *separated* (see [12] for details) whenever $f \cong g$ implies $f = g$, for all $\mathcal{T}$-functors $f, g : Y \longrightarrow X$ with codomain X. One easily verifies that the $\mathcal{T}$*-category* $\mathsf{V} = (\mathsf{V}, \hom_\xi)$ *is separated*, and so is each $\mathcal{T}$-category of the form $\mathcal{P}X$ for a $\mathcal{T}$-category X. The full subcategory of $\mathcal{T}$-Cat consisting of all separated $\mathcal{T}$-categories is denoted by

$$\mathcal{T}\text{-}\mathsf{Cat}_{\mathrm{sep}}.$$

The 2-categorical structure on $\mathcal{T}$-Cat allows us to consider adjoint $\mathcal{T}$-functors: $\mathcal{T}$-functor $f : X \longrightarrow Y$ is *left adjoint* if there exists a $\mathcal{T}$-functor $g : Y \longrightarrow X$ such that $1_X \leq g \cdot f$ and $1_Y \geq f \cdot g$. Considering the corresponding $\mathcal{T}$-distributors, f is left adjoint to g if and only if $g_* \dashv f_*$, that is, if and only if $f_* = g^*$:

$$f \dashv g \text{ in } \mathcal{T}\text{-}\mathsf{Cat} \;\Leftrightarrow\; g_* \dashv f^* \text{ in } \mathcal{T}\text{-}\mathsf{Mod} \;\Leftrightarrow\; f_* = g^*.$$

A more complete study of this subject can be found in [10, 11].

2. The results

In the sequel we consider a class Φ of $\mathcal{T}$-distributors subject to the following axioms.

(Ax 1). For each $\mathcal{T}$-functor f, $f^* \in \Phi$.

(Ax 2). For all $\varphi \in \Phi$ and all $\mathcal{T}$-functors $f : A \longrightarrow X$ we have

$$f^* \circ \varphi \in \Phi, \qquad \varphi \circ f^* \in \Phi, \qquad f_* \in \Phi \Rightarrow \varphi \circ f_* \in \Phi;$$

whenever the compositions are defined.

(Ax 3). For all $\varphi : X \nrightarrow Y \in \mathcal{T}\text{-}\mathsf{Mod}$,

$$(\forall y \in Y \,.\, y^* \circ \varphi \in \Phi) \Rightarrow \varphi \in \Phi$$

where y^* is induced by $y : 1 \longrightarrow Y$, $* \longmapsto y$.

Condition (Ax 2) requires that Φ is closed under certain compositions. In fact, in most examples Φ will be closed under arbitrary compositions. Furthermore, there is a largest and a smallest such class of $\mathcal{T}$-distributors, namely the class $\mathcal{P}$ of all $\mathcal{T}$-distributors and the class $\mathcal{R} = \{f^* \mid f : X \longrightarrow Y\}$ of all representable $\mathcal{T}$-distributors.

We call a $\mathcal{T}$-functor $f : X \longrightarrow Y$ Φ-*dense* if $f_* \in \Phi$. Certainly, if f is a left adjoint $\mathcal{T}$-functor, with $f \dashv g$, then $f_* = g^* \in \Phi$ and therefore f is Φ-dense. A $\mathcal{T}$-category X is called Φ-*injective* if, for all $\mathcal{T}$-functors $f : A \longrightarrow X$ and fully faithful Φ-dense $\mathcal{T}$-functors $i : A \longrightarrow B$, there exists a $\mathcal{T}$-functor $g : B \longrightarrow X$ such that $g \cdot i \cong f$. Furthermore, X is called Φ-*cocomplete* if each weighted diagram

$$\begin{array}{ccc} Y & \xrightarrow{h} & X \\ {\scriptstyle\varphi}\,\vdots & & \\ Z & & \end{array}$$

with $\varphi \in \Phi$ has a colimit $g \cong \mathrm{colim}(\varphi, h) : Z \longrightarrow X$. A $\mathcal{T}$-functor $f : X \longrightarrow Y$ is Φ-*cocontinuous* if f preserves all existing Φ-weighted colimits. Note that in both cases it is enough to consider diagrams where $h = 1_X$. We denote by

$$\mathcal{T}\text{-}\mathsf{Cocont}^{\Phi}$$

the 2-category of all Φ-cocomplete $\mathcal{T}$-categories and Φ-cocontinuous $\mathcal{T}$-functors, and by $\mathcal{T}\text{-}\mathsf{Cocont}^{\Phi}_{\mathrm{sep}}$ its full subcategory of all Φ-cocomplete and separated $\mathcal{T}$-categories.

If Φ is the class $\mathcal{P}$ of all $\mathcal{T}$-distributors, then $\mathcal{T}\text{-}\mathsf{Cocont}^{\Phi}$ is the category of cocomplete $\mathcal{T}$-categories and left adjoint $\mathcal{T}$-functors (as shown in [11, Prop. 2.12]).

2.1. Lemma. *Consider the (up to isomorphism) commutative triangle*



of $\mathfrak{T}$-functors. Then the following assertions hold.

(1). If g and f are Φ-dense, then so is h.

(2). If h is Φ-dense and g is fully faithful, then f is Φ-dense.

(3). If h is Φ-dense and f is dense, then g is Φ-dense.

Proof. The proof is straightforward: (1) $h_* = g_* \circ f_* \in \Phi$ by (Ax 2), since $g_*, f_* \in \Phi$; (2) $f_* = g^* \circ g_* \circ f_* = g^* \circ h_* \in \Phi$ by (Ax 2), since $h_* \in \Phi$; (3) $g_* = g_* \circ f_* \circ f^* = h_* \circ f^* \in \Phi$ by (Ax 2), since $h_* \in \Phi$. ∎

We put now

$$\Phi X = \{\psi \in \mathcal{P}X \mid \psi \in \Phi\}$$

considered as a subcategory of $\mathcal{P}X$. We have the restriction

$$y^{\Phi}_X : X \longrightarrow \Phi X$$

of the Yoneda map, and each $\psi \in \Phi X$ is a Φ-weighted colimit of representables (see [11, Proposition 2.5]).

2.2. Lemma. *The following assertions hold.*

(1). $y^{\Phi}_X : X \longrightarrow \Phi X$ is Φ-dense.

(2). For each $\mathfrak{T}$-distributor $\varphi : X \multimap Y$, $\varphi \in \Phi$ if and only if $\ulcorner\varphi\urcorner : Y \longrightarrow \mathcal{P}X$ factors through the embedding $\Phi X \hookrightarrow \mathcal{P}X$.

Proof. By the Yoneda Lemma (Corollary 1.5), for any $\psi \in \Phi X$ we have $\psi^* \circ (y^{\Phi}_X)_* = \psi \in \Phi$, therefore $(y^{\Phi}_X)_* \in \Phi$ by (Ax 3) and the assertion (1) follows. To see (2), just observe that $\ulcorner\varphi\urcorner(y) = y^* \circ \varphi$, and use again (Ax 3). ∎

Our next result extends Theorem 2.6 of [11]. We omit its proof because it uses exactly the same arguments.

2.3. THEOREM. *The following assertions are equivalent, for a $\mathfrak{T}$-category X.*

(i). X is Φ-injective.

(ii). $y^\Phi_X : X \longrightarrow \Phi X$ has a left inverse $\mathrm{Sup}^\Phi_X : \Phi X \longrightarrow X$.

(iii). $y^\Phi_X : X \longrightarrow \Phi X$ has a left adjoint $\mathrm{Sup}^\Phi_X : \Phi X \longrightarrow X$.

(iv). X is Φ-cocomplete.

Recall from [11] that, for a given $\mathfrak{T}$-functor $f : X \longrightarrow Y$, we have an adjoint pair of $\mathfrak{T}$-functors $\mathcal{P}f \dashv f^{-1}$ where

$$\begin{array}{ccc} \mathcal{P}f : \mathcal{P}X \longrightarrow \mathcal{P}Y & \text{and} & f^{-1} : \mathcal{P}Y \longrightarrow \mathcal{P}X. \\ \psi \longmapsto \psi \circ f^* & & \psi \longmapsto \psi \circ f_* \end{array}$$

By (Ax 1) and (Ax 2), the $\mathfrak{T}$-functor $\mathcal{P}f : \mathcal{P}X \longrightarrow \mathcal{P}Y$ restricts to a $\mathfrak{T}$-functor

$$\Phi f : \Phi X \longrightarrow \Phi Y.$$

On the other hand, $f^{-1} : \mathcal{P}Y \longrightarrow \mathcal{P}X$ restricts to $f^{-1} : \Phi Y \longrightarrow \Phi X$ provided that f is Φ-dense.

2.4. PROPOSITION. *The following conditions are equivalent for a $\mathfrak{T}$-functor $f : X \longrightarrow Y$.*

(i). f is Φ-dense.

(ii). Φf is left adjoint.

(iii). Φf is Φ-dense.

PROOF. (i) $\Rightarrow$ (ii): If f is Φ-dense, then $\Phi f \dashv f^{-1} : \Phi Y \longrightarrow \Phi X$ defined above. (ii) $\Rightarrow$ (iii): If $\Phi f \dashv g$, then $(\Phi f)_* = g^* \in \Phi$ and Φf is Φ-dense. (iii) $\Rightarrow$ (i): Consider the diagram

$$\begin{array}{ccc} X & \xrightarrow{y^\Phi_X} & \Phi X \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle \Phi f} \\ Y & \xrightarrow[y^\Phi_Y]{} & \Phi Y \end{array}$$

If Φf is Φ-dense, then $y^\Phi_Y \cdot f = \Phi f \cdot y^\Phi_X$ is Φ-dense, and so by 2.1(2) f is Φ-dense because y^Φ_Y is fully faithful. ∎

In particular, for each $\mathfrak{T}$-category X, $\Phi\, y^\Phi_X : \Phi X \longrightarrow \Phi\Phi X$ has a right adjoint, $(y^\Phi_X)^{-1}$. We show next that $(y^\Phi_X)^{-1}$ has also a right adjoint, $y^\Phi_{\Phi X} : \Phi X \longrightarrow \Phi\Phi X$, so that:

$$\Phi\, y^\Phi_X \dashv (y^\Phi_X)^{-1} = \mathrm{Sup}^\Phi_{\Phi X} \dashv y^\Phi_{\Phi X}\,.$$

2.5. Proposition. *For each $\mathfrak{T}$-category X, ΦX is Φ-cocomplete where $\mathrm{Sup}^\Phi_{\Phi X} = (y^\Phi_X)^{-1}$.*

Proof. Since y^Φ_X is Φ-dense, we may define $\mathrm{Sup}^\Phi_{\Phi X} := (y^\Phi_X)^{-1}$. We have to show that $\mathrm{Sup}^\Phi_{\Phi X}$ is a left inverse for $y^\Phi_{\Phi X}$; that is, $(y^\Phi_X)^{-1} \cdot y^\Phi_{\Phi X} = 1_{\Phi X}$: for each $\psi \in \Phi X$, $((y^\Phi_X)^{-1} \cdot y^\Phi_{\Phi X})(\psi) = \psi^* \circ (y^\Phi_X)_* = \psi$. ∎

In [11] we constructed $\mathcal{P}f$ as the colimit $\mathcal{P}f \cong \mathrm{colim}((y_X)_*, y_Y \cdot f)$, and a straightforward calculation shows that also $\Phi f \cong \mathrm{colim}((y^\Phi_X)_*, y^\Phi_Y \cdot f)$, for each $\mathfrak{T}$-functor $f : X \longrightarrow Y$. To see this, we consider the commutative diagrams



and obtain

$$\begin{aligned}
(\Phi f)_* &= i^*_Y \circ i_{Y*} \circ (\Phi f)_* \\
&= i^*_Y \circ (\mathcal{P}f)_* \circ i_{X*} \\
&= i^*_Y \circ ((y_{Y*} \circ f_*) \circ\!\!- y_{X*}) \circ i_{X*} && \text{since } \mathcal{P}f \cong \mathrm{colim}((y_X)_*, y_Y \cdot f) \\
&= (i^*_Y \circ y_{Y*} \circ f_*) \circ\!\!- (i_X{}^* \circ y_{X*}) && \text{by Lemma 1.2} \\
&= (y^\Phi_{Y*} \circ f_*) \circ\!\!- y^\Phi_{X*}.
\end{aligned}$$

2.6. Proposition. *Let $f : X \longrightarrow Y$ a $\mathfrak{T}$-functor where X and Y are Φ-cocomplete.*

(1). The following assertions are equivalent.

(a) f is Φ-cocontinuous.

(b) We have $f \cdot \mathrm{Sup}^\Phi_X \cong \mathrm{Sup}^\Phi_Y \cdot \Phi f$.



(2). f is Φ-cocontinuous and Φ-dense if and only it is a left adjoint.

PROOF. (1) (a) $\Rightarrow$ (b): Recall that



Hence

$$\begin{aligned}(f \cdot \mathrm{Sup}_X^\Phi)_* &= f_* \circ\!\!- (y_X^\Phi)_* \\ &= ((\mathrm{Sup}_Y^\Phi)_* \circ (y_Y^\Phi)_* \circ f_*) \circ\!\!- (y_X^\Phi)_* \\ &= (\mathrm{Sup}_Y^\Phi)_* \circ ((y_Y^\Phi)_* \circ f_* \circ\!\!- (y_X^\Phi)_*) \\ &= (\mathrm{Sup}_Y^\Phi)_* \circ \Phi f_*.\end{aligned}$$

(b)$\Rightarrow$ (a): Consider



Then

$$\begin{aligned}(f \cdot \mathrm{Sup}_X^\Phi \cdot \ulcorner\varphi\urcorner) &= \mathrm{Sup}_Y^\Phi \cdot \Phi f \cdot \ulcorner\varphi\urcorner \\ &= \mathrm{Sup}_Y^\Phi \cdot \ulcorner\varphi \cdot f^*\urcorner \\ &\cong \mathrm{colim}(\varphi, f)\end{aligned}$$

(2) If f is Φ-cocontinuous and Φ-dense, from the commutative diagram of (1)(b) we have $f \dashv \mathrm{Sup}_X^\Phi \cdot f^{-1} \cdot y_Y^\Phi$ since $f \cdot \mathrm{Sup}_X^\Phi = \mathrm{Sup}_Y^\Phi \cdot \Phi f \dashv f^{-1} \cdot y_Y^\Phi$ and $\mathrm{Sup}_X^\Phi \cdot y_X^\Phi = 1_X$. The converse is trivially true. ∎

2.7. COROLLARY. *ΦX is closed in $\mathcal{P}X$ under Φ-weighted colimits.*

PROOF. We show that the inclusion functor $i : \Phi X \longrightarrow \mathcal{P}X$ is Φ-cocontinuous, which, by the proposition above, is equivalent to the commutativity of the diagram



In Proposition 2.5 we observed $\mathrm{Sup}_{\Phi X}^\Phi = (y_X^\Phi)^{-1}$, and from Theorem 2.3 and [11, Theorem 2.8] follows that $\mathrm{Sup}_{\mathcal{P}X}^\Phi$ is the restriction of $y_X^{-1} : \mathcal{P}\mathcal{P}X \longrightarrow \mathcal{P}X$ to $\Phi\mathcal{P}X$. Let $\Psi \in \Phi\Phi X$. Then

$$i \cdot (y_X^\Phi)^{-1}(\Psi) = \Psi \circ (y_X^\Phi)_*$$

and

$$y_X^{-1} \cdot \Phi i(\Psi) = y_X^{-1}(\Psi \circ i^*) = \Psi \circ i^* \circ (y_X)_* = \Psi \circ (y_X^\Phi)_*,$$

and the assertion follows. ∎

Theorem 2.3 says in particular that, for each $\mathcal{T}$-functor $f : A \longrightarrow X$, Φ-injective $\mathcal{T}$-category X and fully faithful Φ-dense $\mathcal{T}$-functor $i : A \longrightarrow B$, we have a canonical extension $g : B \longrightarrow X$ of f along i, namely $g \cong \mathrm{colim}(i_*, f)$, giving us an alternative description of Φf.

2.8. THEOREM. *Composition with $y^\Phi_X : X \longrightarrow \Phi X$ defines an equivalence*

$$\mathcal{T}\text{-}\mathsf{Cocont}^\Phi(\Phi X, Y) \longrightarrow \mathcal{T}\text{-}\mathsf{Cat}(X, Y)$$

of ordered sets, for each Φ-cocomplete $\mathcal{T}$-category Y.

The series of results above tell us that $\mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}}$ is actually a (non-full) reflective subcategory of $\mathcal{T}\text{-}\mathsf{Cat}$, with left adjoint $\Phi : \mathcal{T}\text{-}\mathsf{Cat} \longrightarrow \mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}}$. In fact, Φ is a 2-functor and one verifies as in [11] that the induced monad $\mathbb{I}^\Phi = (\Phi, y^\Phi, (y^\Phi)^{-1})$ on $\mathcal{T}\text{-}\mathsf{Cat}$ is of Kock-Zöberlein type. Theorem 2.3 and Proposition 2.6 imply that $\mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}}$ is equivalent to the category of Eilenberg-Moore algebras of $\mathbb{I}^\Phi$.

Finally, we wish to study monadicity of the canonical forgetful functor

$$\mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}} \xrightarrow{\ \mathrm{G}\ } \mathsf{Set}.$$

Certainly,

(a) G has a left adjoint given by the composite

$$\mathsf{Set} \xrightarrow{\ \mathrm{disc}\ } \mathcal{T}\text{-}\mathsf{Cat} \xrightarrow{\ \Phi\ } \mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}},$$

where $\mathrm{disc}(X) = (X, e^\circ_X)$, and $\mathrm{disc}(f) = f$.

In order to prove monadicity of G we will impose, in addition to (Ax 1)-(Ax 3),

(Ax 4). For each surjective $\mathcal{T}$-functor f, $f_* \in \Phi$.

Hence, any bijective $f : X \longrightarrow Y$ in $\mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}}$ is Φ-dense and therefore left adjoint. By [11, Lemma 2.16], f is invertible and we have seen that

(b) G reflects isomorphisms.

In order to conclude that G is monadic, it is left to show that

(c) $\mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}}$ has and G preserves coequaliser of G-equivalence relations

(see, for instance, [15, Corollary 2.7]). To do so, let $\pi_1, \pi_2 : R \rightrightarrows X$ in $\mathcal{T}\text{-}\mathsf{Cocont}^\Phi_{\mathrm{sep}}$ be an equivalence relation in Set, where π_1 and π_2 are the projection maps, and let $q : X \longrightarrow Q$ be its coequaliser in $\mathcal{T}\text{-}\mathsf{Cat}$. The proof in [11, Section 2.6] rests on the observation that

$$\mathcal{P}R \underset{\mathcal{P}\pi_2}{\overset{\mathcal{P}\pi_1}{\rightrightarrows}} \mathcal{P}X \xrightarrow{\ \mathcal{P}q\ } \mathcal{P}Q$$

is a split fork in $\mathcal{T}$-$\mathsf{Cat}_{\mathrm{sep}}$. Naturally, we wish to show that, in our setting,

$$\Phi R \underset{\Phi\pi_2}{\overset{\Phi\pi_1}{\rightrightarrows}} \Phi X \xrightarrow{\Phi q} \Phi Q$$

gives rise to a split fork in $\mathcal{T}$-$\mathsf{Cat}_{\mathrm{sep}}$ as well. Since π_1, π_2 and q are surjective, the $\mathcal{T}$-functors π_1, π_2 and q are Φ-dense and therefore we have $\mathcal{T}$-functors $q^{-1} : \Phi Q \longrightarrow \Phi X$ and $\pi_1^{-1} : \Phi X \longrightarrow \Phi R$. Furthermore, $\Phi q \cdot q^{-1} = 1_{\Phi X} = \Phi\pi_1 \cdot \pi_1^{-1}$. It is left to show that

$$q^{-1} \cdot \Phi q = \Phi\pi_2 \cdot \pi_1^{-1},$$

which can be shown with the same calculation as in [11], based on the following proposition.

2.9. Proposition. *Consider the following diagram in* $\mathcal{T}$-Cat

$$R \underset{\pi_2}{\overset{\pi_1}{\rightrightarrows}} X \xrightarrow{q} Q$$

with $\pi_1, \pi_2 : R \rightrightarrows X$ *in* $\mathcal{T}$-$\mathsf{Cocont}^{\Phi}_{\mathrm{sep}}$, (π_1, π_2) *an equivalence relation in* Set, *and* $q : X \longrightarrow Q$ *its coequaliser in* $\mathcal{T}$-Cat.

(1). If π_1, π_2 *are left adjoints, then* q *is proper.*

(2). The diagram

$$\Phi R \underset{\Phi\pi_2}{\overset{\Phi\pi_1}{\rightrightarrows}} \Phi X \xrightarrow{\Phi q} \Phi Q \qquad \pi_1^{-1} \text{ over } \Phi\pi_1, \quad q^{-1} \text{ over } \Phi q$$

is a split fork in $\mathcal{T}$-Cat.

Proof. (1) As in [11, Lemma 2.19 and Corollary 2.20].

(2) Analogous to the proof presented in [11, Section 2.6]. ∎

Finally, we conclude that:

2.10. Theorem. *Under (Ax 1)-(Ax 4), the forgetful functor* $G : \mathcal{T}$-$\mathsf{Cocont}^{\Phi}_{\mathrm{sep}} \longrightarrow \mathsf{Set}$ *is monadic.*

Proof. In order to show that $\mathcal{T}$-$\mathsf{Cocont}^{\Phi}_{\mathrm{sep}}$ has and G preserves coequaliser of G-equivalence relations, consider again the first diagram of Proposition 2.9. We have seen that

$$\Phi R \underset{\Phi\pi_2}{\overset{\Phi\pi_1}{\rightrightarrows}} \Phi X \xrightarrow{\Phi q} \Phi Q \qquad \pi_1^{-1} \text{ over } \Phi\pi_1, \quad q^{-1} \text{ over } \Phi q$$

is a split fork and hence a coequaliser diagram in $\mathcal{T}$-Cat. Since π_1 and π_2 are Φ-cocontinuous, there is a $\mathcal{T}$-functor $\mathrm{Sup}^{\Phi}_Q : \Phi Q \longrightarrow Q$ which, since $q : X \longrightarrow Q$ is the coequaliser of

$\pi_1, \pi_2 : R \rightrightarrows X$ in $\mathcal{T}$-Cat, satisfies $\mathrm{Sup}_Q^\Phi \cdot y_Q^\Phi = 1_Q$. The situation is depicted in the following diagram.



We conclude that Q is separated and Φ-cocomplete, and $q : X \longrightarrow Q$ is Φ-cocontinuous. Finally, to see that $q : X \longrightarrow Q$ is the coequaliser of $\pi_1, \pi_2 : R \rightrightarrows X$ in $\mathcal{T}$-$\mathsf{Cocont}^\Phi_{\mathrm{sep}}$, let $h : X \longrightarrow Y$ be in $\mathcal{T}$-$\mathsf{Cocont}^\Phi_{\mathrm{sep}}$ with $h \cdot \pi_1 = h \cdot \pi_2$. Then, since Φq is the coequaliser of $\Phi\pi_1, \Phi\pi_2 : \Phi R \rightrightarrows \Phi X$ in $\mathcal{T}$-$\mathsf{Cocont}^\Phi_{\mathrm{sep}}$, there is a Φ-cocontinuous $\mathcal{T}$-functor $f : \Phi Q \longrightarrow Y$ such that $f \cdot \Phi q = h \cdot \mathrm{Sup}_X^\Phi$. Then

$$f \cdot y_Q^\Phi \cdot q = f \cdot \Phi q \cdot y_X^\Phi = h \cdot \mathrm{Sup}_X^\Phi \cdot y_X^\Phi = h$$

and

$$\begin{aligned}\mathrm{Sup}_Y^\Phi \cdot \Phi f \cdot \Phi\, y_Q^\Phi \cdot \Phi q = f \cdot \mathrm{Sup}_{\Phi Q}^\Phi \cdot \Phi\, y_Q^\Phi \cdot \Phi q = f \cdot \Phi q = h \cdot \mathrm{Sup}_X^\Phi \\ = f \cdot y_Q^\Phi \cdot q \cdot \mathrm{Sup}_X^\Phi = f \cdot y_Q^\Phi \cdot \mathrm{Sup}_Q^\Phi \cdot \Phi q,\end{aligned}$$

hence $\mathrm{Sup}_Y \cdot \Phi(f \cdot y_Q^\Phi) = f \cdot y_Q^\Phi \cdot \mathrm{Sup}_Q^\Phi$, that is, $f \cdot y_Q^\Phi$ is Φ-cocontinuous. ∎

3. The examples

3.1. All distributors. The class $\Phi = \mathcal{P}$ of all distributors satisfies obviously all four axioms. In fact, this is the situation studied in [11].

3.2. Representable distributors. The smallest possible choice is $\Phi = \mathcal{R}$ being the class of all representable $\mathcal{T}$-distributors $\mathcal{R} = \{f^* \mid f \text{ is a } \mathcal{T}\text{-functor}\}$. Clearly, $\mathcal{R}$ satisfies (Ax 1), (Ax 2) and (Ax 3) but not (Ax 4). We have $\mathcal{R}(X) = \{x^* \mid x \in X\}$, each $\mathcal{T}$-category is $\mathcal{R}$-cocomplete and each $\mathcal{T}$-functor is $\mathcal{R}$-cocontinuous, and therefore $\mathcal{T}$-$\mathsf{Cocont}^{\mathcal{R}}_{\mathrm{sep}} = \mathcal{T}$-$\mathsf{Cat}_{\mathrm{sep}}$. This case is certainly not very interesting; however, our results tell us that the inclusion functor $\mathcal{T}$-$\mathsf{Cat}_{\mathrm{sep}} \hookrightarrow \mathcal{T}$-$\mathsf{Cat}$ is monadic. In particular, *the category* Top_0 *of topological* T_0*-spaces and continuous maps is a monadic subcategory of* Top.

3.3. Almost representable distributors. We can modify slightly the example above and consider $\Phi = \mathcal{R}_0$ the class of all almost representable $\mathcal{T}$-distributors, where a $\mathcal{T}$-distributor $\varphi : X \nrightarrow Y$ is called *almost representable* whenever, for each $y \in Y$, either $y^* \circ \varphi = \bot$ or $y^* \circ \varphi = x^*$ for some $x \in X$. As above, $\mathcal{R}_0$ satisfies (Ax 1), (Ax 2) and (Ax 3) but not (Ax 4).

By definition, for a $\mathcal{T}$-category X we have

$$\mathcal{R}_0(X) = \{\psi \in \mathcal{P}X \mid \psi \in \mathcal{R}_0\} = \{x^* \mid x \in X\} \cup \{\bot\},$$

with the structure inherited from $\mathcal{P}X$. Furthermore, a $\mathcal{T}$-functor $f : (X, a) \longrightarrow (Y, b)$ is $\mathcal{R}_0$-dense whenever, for each $y \in Y$,

$$\exists \mathfrak{x} \in TX \,.\, b(Tf(\mathfrak{x}), y) > \bot \;\Rightarrow\; \exists x \in X \,\forall \mathfrak{x} \in TX \,.\, b(Tf(\mathfrak{x}), y) = a(\mathfrak{x}, x).$$

Hence, with

$$Y_0 = \{y \in Y \mid \exists \mathfrak{x} \in TX \,.\, b(Tf(\mathfrak{x}), y) > \bot\}$$

we can factorise an $\mathcal{R}_0$-dense $\mathcal{T}$-functor $f : X \longrightarrow Y$ as

$$X \xrightarrow{f} Y_0 \hookrightarrow Y,$$

where $Y_0 \hookrightarrow Y$ is fully faithful and $X \xrightarrow{f} Y_0$ is left adjoint. If we consider $f : X \longrightarrow Y$ in Top, then $Y_0 = \overline{f(X)}$ is the closure of the image of f, so that each $\mathcal{R}_0$-dense continuous map factors as a left adjoint continuous map followed by a closed embedding. Consequently, *for a topological space X, the following assertions are equivalent:*

(i). X is injective with respect to $\mathcal{R}_0$-dense fully faithful continuous maps.

(ii). X is injective with respect to closed embeddings.

Note that in this example we are working with the dual order, compared with [8, Section 11].

3.4. Right adjoint distributors. Now we consider $\Phi = \mathcal{L}$ the class of all right adjoint $\mathcal{T}$-distributors. This class contains all distributors of the form f^*, for a $\mathcal{T}$-functor f, and it is closed under composition. Since adjointness of a $\mathcal{T}$-distributor $\varphi : X \multimap Y$ can be tested pointwise in Y, the axioms (Ax 1), (Ax 2) and (Ax 3) are satisfied. By definition, $\mathcal{L}(X) = \{\psi \in \mathcal{P}X \mid \psi \text{ is right adjoint}\}$, and a $\mathcal{T}$-category is $\mathcal{L}$-cocomplete if each pair $\varphi \dashv \psi$, $\varphi : Y \multimap X$, $\psi : X \multimap Y$, of adjoint $\mathcal{T}$-distributors is of the form $f_* \dashv f^*$, for a $\mathcal{T}$-functor $f : Y \longrightarrow X$. For V-categories, this is precisely the well-known notion of Cauchy-completeness as introduced by Lawvere in [14] as a generalisation of the classical notion for metric spaces. However, Lawvere never proposed the name "Cauchy-complete", and, while working on this notion in the context of $\mathcal{T}$-categories in [6] and [12], we used instead Lawvere-complete and L-complete, respectively. Furthermore, one easily verifies that each $\mathcal{T}$-functor is $\mathcal{L}$-cocontinuous, that is, (right adjoint)-weighted colimits are absolute, so that $\mathcal{T}\text{-}\mathsf{Cocont}^{\mathcal{L}}_{\mathrm{sep}} = \mathcal{T}\text{-}\mathsf{Cat}_{\mathrm{cpl}}$ is the full subcategory of $\mathcal{T}\text{-}\mathsf{Cat}$ consisting of all separated and Lawvere complete $\mathcal{T}$-categories.

On the other hand, for a surjective $\mathcal{T}$-functor f, f_* does not need to be right adjoint, so that (Ax 4) is in general not satisfied. This is not a surprise, since natural instances of this example fail Theorem 2.10. Indeed, in the category of ordered sets and monotone maps, any ordered set is Lawvere-complete, hence the category of Lawvere-complete and

separated ordered sets coincides with the category of anti-symmetric ordered sets. The canonical forgetful functor from this category to Set is surely not monadic. Also, the canonical forgetful functor from the category of Lawvere-complete and separated topological spaces (= sober spaces) and continuous maps to Set is also not monadic.

3.5. INHABITED DISTRIBUTORS. Another class of distributors considered in [11] is $\Phi = \mathcal{I}$ the class of all inhabited $\mathfrak{T}$-distributors. Here a $\mathfrak{T}$-distributor $\varphi : X \mathrel{-\!\!\circ\!\!\rightarrow} Y$ is called *inhabited* if

$$\forall y \in Y \,.\, k \leq \bigvee_{\mathfrak{x} \in TX} \varphi(\mathfrak{x}, y).$$

(Ax 3) is satisfied by definition, and in [11] we showed already the validity of (Ax 1) and (Ax 2). Furthermore, one easily verifies that (Ax 4) is satisfied. Hence, as already observed in [11], all results stated in Section 2 are available for this class of distributors. Let us recall that, specialised to Top, inhabited-dense continuous maps are precisely the topologically dense continuous maps, and the injective spaces with respect to topologically dense embeddings are known as *Scott domains* [9].

3.6. "CLOSED" DISTRIBUTORS. A further interesting class of distributors is given by

$$\Phi = \{\varphi : X \mathrel{-\!\!\circ\!\!\rightarrow} Y \mid \forall y \in Y,\, \mathfrak{x} \in TX \,.\, \varphi(\mathfrak{x}, y) \leq \bigvee_{x \in X} a(\mathfrak{x}, x) \otimes \varphi(e_X(x), y)\},$$

that is, $\varphi \in \Phi$ if and only if $\varphi \leq \varphi \cdot e_X \cdot a$. Clearly, (Ax 3) is satisfied. For each $\mathfrak{T}$-functor $g : (Y, b) \longrightarrow (X, a)$ we have

$$g^* \cdot e_X \cdot a = g^\circ \cdot a \cdot e_X \cdot a \geq g^\circ \cdot a = g^*,$$

hence $g^* \in \Phi$. Furthermore, given $\mathfrak{T}$-distributors $\varphi : X \mathrel{-\!\!\circ\!\!\rightarrow} Y$ and $\psi : Y \mathrel{-\!\!\circ\!\!\rightarrow} Z$ in Φ, then

$$\begin{aligned}\psi \circ \varphi = \psi \cdot T_\xi \varphi \cdot m_X^\circ &\leq \psi \cdot e_Y \cdot b \cdot T_\xi \varphi \cdot m_X^\circ = \psi \cdot e_Y \cdot \varphi \leq \psi \cdot e_Y \cdot \varphi \cdot e_X \cdot a \\ &\leq \psi \cdot T_\xi \varphi \cdot e_{TX} \cdot e_X \cdot a \leq \psi \cdot T_\xi \varphi \cdot m_X^\circ \cdot e_X \cdot a = (\psi \circ \varphi) \cdot e_X \cdot a\end{aligned}$$

and therefore also $\psi \circ \varphi \in \Phi$. We have seen that this class of distributors satisfies (Ax 1), (Ax 2) and (Ax 3). On the other hand, (Ax 4) is not satisfied.

By definition, a $\mathfrak{T}$-functor $f : (X, a) \longrightarrow (Y, b)$ is Φ-dense whenever, for all $\mathfrak{x} \in TX$ and $y \in Y$,

$$b(Tf(\mathfrak{x}), y) \leq \bigvee_{x \in X} a(\mathfrak{x}, x) \otimes b(e_Y(f(x)), y).$$

Hence, each proper $\mathfrak{T}$-functor (see [4]) is Φ-dense. In fact, Φ-dense $\mathfrak{T}$-functors can be seen as "proper over V-Cat", and the condition above states exactly properness of f if the underlying V-category $\mathrm{S}\,Y$ of $Y = (Y, b)$ is discrete. Furthermore, each surjective Φ-dense $\mathfrak{T}$-functor is final with respect to the forgetful functor $\mathrm{S} : \mathfrak{T}\text{-}\mathsf{Cat} \longrightarrow \mathsf{V}\text{-}\mathsf{Cat}$. To see this, let $f : (X, a) \longrightarrow (Y, b)$ be a surjective Φ-dense $\mathfrak{T}$-functor, $Z = (Z, c)$ a $\mathfrak{T}$-category and $g : \mathrm{S}\,Y \longrightarrow \mathrm{S}\,Z$ a V-functor such that gf is a $\mathfrak{T}$-functor. We have to show that g is a

$\mathfrak{T}$-functor. Let $\mathfrak{y} \in TY$ and $y \in Y$. Since Tf is surjective, there is some $\mathfrak{x} \in TX$ with $Tf(\mathfrak{x}) = \mathfrak{y}$. We conclude

$$\begin{aligned} b(\mathfrak{y}, y) &= b(Tf(\mathfrak{x}), y) \\ &\leq \bigvee_{x \in X} a(\mathfrak{x}, x) \otimes b(e_Y(f(x)), y) \\ &\leq \bigvee_{x \in X} c(T(gf)(\mathfrak{x}), gf(x) \otimes c(e_Z(gf(x)), g(y)) \\ &\leq c(Tg(\mathfrak{y}), g(y)). \end{aligned}$$

3.7. Further examples. A wide class of examples of injective topological spaces is described in [8], where the authors consider injectivity with respect to a class of embeddings $f : X \longrightarrow Y$ such that the induced frame morphism $f_* : \Omega X \longrightarrow \Omega Y$ preserves certain suprema. A similar construction can be done in our setting; to do so *we assume from now on* $T1 = 1$. For a $\mathfrak{T}$-category X, the V-*category of covariant presheafs* V^X is defined as

$$\mathsf{V}^X = \{\alpha : 1 \multimap X \mid \alpha \text{ is a } \mathfrak{T}\text{-distributor}\} = \{\alpha : X \longrightarrow \mathsf{V} \mid \alpha \text{ is a } \mathfrak{T}\text{-functor}\},$$

and the V-categorical structure $[\alpha, \beta] \in \mathsf{V}$ is given as the lifting



for all $\alpha, \beta \in \mathsf{V}^X$. Since $e_1 : 1 \longrightarrow T1$ is an isomorphism, this lifting of $\mathfrak{T}$-distributors does exist and can be calculated as the corresponding lifting of V-distributors



Each $\mathfrak{T}$-distributor $\varphi : X \multimap Y$ induces a V-functor

$$\varphi \circ (-) : \mathsf{V}^X \longrightarrow \mathsf{V}^Y,\ \alpha \longmapsto \varphi \circ \alpha,$$

which is right adjoint if φ is a right adjoint $\mathfrak{T}$-distributor. Given now a class Ψ of V-distributors, we may consider the class Φ of all those $\mathfrak{T}$-distributors φ for which $\varphi \circ (-)$ preserves Ψ-weighted limits. This class of $\mathfrak{T}$-distributors is certainly closed under composition, and contains all right adjoint $\mathfrak{T}$-distributors, hence it includes all representable ones. Finally, if Ψ-weighted limits are calculated pointwise in V^X, then also (Ax 3) is fulfilled. As particular examples we have the class Φ of all $\mathfrak{T}$-distributors $\varphi : X \multimap Y$ for which $\varphi \circ (-)$ preserves

(1). the top element of V^X, that is, for which $\varphi \circ \top = \top$. In pointwise notation, this reads as

$$\forall y \in Y \,.\, \top = \bigvee_{\mathfrak{x} \in TX} \varphi(\mathfrak{x}, y) \otimes \top.$$

If $k = \top$, then this class of $\mathfrak{T}$-distributors coincides with the class of inhabited $\mathfrak{T}$-distributors considered in 3.5.

(2). cotensors, that is, for each $u \in \mathsf{V}$ and each $\alpha \in \mathsf{V}^X$, $\varphi \circ \hom(u, \alpha) = \hom(u, \varphi \circ \alpha)$.

(3). finite infima (cf. [8, Section 6]).

(4). arbitrary infima (cf. [8, Section 7]).

(5). codirected infima (cf. [8, Section 8]).

References

[1] M. Barr, *Relational algebras*, in Reports of the Midwest Category Seminar, IV, Lecture Notes in Mathematics, Vol. 137. Springer, Berlin, 1970, pp. 39–55.

[2] R. Betti, A. Carboni, R. Street, and R. Walters, *Variation through enrichment*, J. Pure Appl. Algebra, 29 (1983), pp. 109–127.

[3] M. M. Clementino and D. Hofmann, *Topological features of lax algebras*, Appl. Categ. Structures, 11 (2003), pp. 267–286.

[4] ——, *Effective descent morphisms in categories of lax algebras*, Appl. Categ. Structures, 12 (2004), pp. 413–425.

[5] ——, *Exponentiation in V-categories*, Topology Appl., 153 (2006), pp. 3113–3128.

[6] ——, *Lawvere completeness in Topology*, Accepted for publication in Appl. Categ. Structures, (2008), arXiv:math.CT/0704.3976.

[7] M. M. Clementino and W. Tholen, *Metric, topology and multicategory—a common approach*, J. Pure Appl. Algebra, 179 (2003), pp. 13–47.

[8] M. Escardó and B. Flagg, *Semantic domains, injective spaces and monads.* Brookes, Stephen (ed.) et al., Mathematical foundations of programming semantics. Proceedings of the 15th conference, Tulane Univ., New Orleans, LA, April 28 - May 1, 1999. Amsterdam: Elsevier, Electronic Notes in Theoretical Computer Science. 20, electronic paper No.15 (1999)., 1999.

[9] G. Gierz, K. H. Hofmann, K. Keimel, J. D. Lawson, M. W. Mislove, and D. S. Scott, *A compendium of continuous lattices*, Springer-Verlag, Berlin, 1980.

[10] D. Hofmann, *Topological theories and closed objects*, Adv. Math., 215 (2007), pp. 789–824.

[11] ——, *Injective spaces via adjunction*, tech. rep., University of Aveiro, 2008, arXiv:math.CT/0804.0326.

[12] D. Hofmann and W. Tholen, *Lawvere completion and separation via closure*, *Appl. Categ. Structures*, to appear, arXiv:math.CT/0801.0199.

[13] G. M. Kelly and V. Schmitt, *Notes on enriched categories with colimits of some class*, Theory Appl. Categ., 14 (2005), pp. no. 17, 399–423 (electronic).

[14] F. W. Lawvere, *Metric spaces, generalized logic, and closed categories*, Rend. Sem. Mat. Fis. Milano, 43 (1973), pp. 135–166 (1974). Also in: Repr. Theory Appl. Categ. 1:1–37 (electronic), 2002.

[15] J. MacDonald and M. Sobral, *Aspects of monads*, in Categorical foundations, vol. 97 of Encyclopedia Math. Appl., Cambridge Univ. Press, Cambridge, 2004, pp. 213–268.

Departamento de Matemática
Universidade de Coimbra
3001-454 Coimbra
Portugal

Departamento de Matemática
Universidade de Aveiro
3810-193 Aveiro
Portugal

Email: mmc@mat.uc.pt
dirk@ua.pt

This article may be accessed at http://www.tac.mta.ca/tac/ or by anonymous ftp at ftp://ftp.tac.mta.ca/pub/tac/html/volumes/21/12/21-12.{dvi,ps,pdf}

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.
Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.
Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at `http://www.tac.mta.ca/tac/` and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION. Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to `tac@mta.ca` including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, `rrosebrugh@mta.ca`.

INFORMATION FOR AUTHORS. The typesetting language of the journal is TEX, and LATEX2e strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at `http://www.tac.mta.ca/tac/`.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: `rrosebrugh@mta.ca`

TEXNICAL EDITOR. Michael Barr, McGill University: `barr@math.mcgill.ca`

ASSISTANT TEX EDITOR. Gavin Seal, McGill University: `gavin_seal@fastmail.fm`

TRANSMITTING EDITORS.
Richard Blute, Université d' Ottawa: `rblute@uottawa.ca`
Lawrence Breen, Université de Paris 13: `breen@math.univ-paris13.fr`
Ronald Brown, University of North Wales: `ronnie.profbrown (at) btinternet.com`
Aurelio Carboni, Università dell Insubria: `aurelio.carboni@uninsubria.it`
Valeria de Paiva, Cuill Inc.: `valeria@cuill.com`
Ezra Getzler, Northwestern University: `getzler(at)northwestern(dot)edu`
Martin Hyland, University of Cambridge: `M.Hyland@dpmms.cam.ac.uk`
P. T. Johnstone, University of Cambridge: `ptj@dpmms.cam.ac.uk`
Anders Kock, University of Aarhus: `kock@imf.au.dk`
Stephen Lack, University of Western Sydney: `s.lack@uws.edu.au`
F. William Lawvere, State University of New York at Buffalo: `wlawvere@acsu.buffalo.edu`
Jean-Louis Loday, Université de Strasbourg: `loday@math.u-strasbg.fr`
Ieke Moerdijk, University of Utrecht: `moerdijk@math.uu.nl`
Susan Niefield, Union College: `niefiels@union.edu`
Robert Paré, Dalhousie University: `pare@mathstat.dal.ca`
Jiri Rosicky, Masaryk University: `rosicky@math.muni.cz`
Brooke Shipley, University of Illinois at Chicago: `bshipley@math.uic.edu`
James Stasheff, University of North Carolina: `jds@math.unc.edu`
Ross Street, Macquarie University: `street@math.mq.edu.au`
Walter Tholen, York University: `tholen@mathstat.yorku.ca`
Myles Tierney, Rutgers University: `tierney@math.rutgers.edu`
Robert F. C. Walters, University of Insubria: `robert.walters@uninsubria.it`
R. J. Wood, Dalhousie University: `rjwood@mathstat.dal.ca`