Записки научных
семинаров ПОМИ
Том 300, 2003 г.

E. V. Volkov

HYPERBOLIC STRUCTURE FOR THE QUADRATIC MAP

Abstract. The quadratic map of the real plane is considered. A possible construction of a hyperbolic structure on some non compact set is suggested and discussed.

§0. Introduction

Let M be a Riemannian manifold, $G : M \longrightarrow M$ be a diffeomorphism. Let $M_g \subset M$ be an invariant set with respect to the map G (i.e. $G(M_g) = M_g$). One says, that the set M_g possesses a weak uniform hyperbolic structure with respect to the map G, if for each point $x \in M_g$ the tangent space T_xM can be represented as a direct sum:

$$T_xM = E_s(x) \dot{+} E_u(x)$$

and for each constant $\lambda > 1$ there exists a constant $c > 0$ such that

$$\forall v \in E_s \ \ ||(G^n(x))' v|| \leqslant c\lambda^{-n}||v||, \ n = 1, 2, ...,$$

$$\forall v \in E_u \ \ ||(G^{-n}(x))' v|| \leqslant c\lambda^{-n}||v||, \ n = 1, 2, ...,$$

E_s, E_u are called stable and unstable subspaces (or directions if they are 1-dimensional).

We consider a generic area-preserving quadratic map of the real plane $F : \mathbb{R}^2 \longrightarrow \mathbb{R}^2$ defined as:

$$x_1 = x^2 - y + C, \tag{1}$$

$$y_1 = x, \tag{2}$$

$$F(x, y) = (x_1, y_1), \tag{3}$$

where C is a real parameter.

This work was supported by grants: INTAS 00-221; RFBR for basic research 01-01-00335; Russian Federation ministry of education grant E00-1-120.

Let M_g be an open subset of $\mathbb{R}^2$ invariant with respect to the map F. The main question we study is how to describe such a subset M_g being as wide as possible, which possesses a weak uniform hyperbolic structure with respect to the map F.

Consider an open set D consisting of points of the plane which tend to infinity under both forward and backward iterates. We prove the following results:

1. Set D is "wide enough".

2. There are stable and unstable directions at every point of the set D.

3. There exists a subset Bad of the set D, which is closed in the topology of D and of zero Lebesgue measure with the following property: the stable and unstable directions at each point of Bad coincide to each other: $E_s = E_u$. The situation in the set Bad is worse for us. But if we consider $E = D \backslash Bad$, we can construct a hyperbolic structure on E.

Here is the list of notations which might produce a confusion.

1. "$\mathbb{K}$" means an element of the set $\{\mathbb{R}, \mathbb{C}\}$,

2. "$BOUND$" means the set of all bounded subsets of the real plane $\mathbb{R}^2$.

§1. Basic facts on the dynamics of the quadratic map

To begin with we introduce some domains of $\mathbb{K}^2$ which possess important properties with respect to the quadratic map. Let A and B be real numbers. Consider a domain in $\mathbb{C}^2$ defined as:

$$\Omega^{\mathbb{C}}_{AB} = \{(x, y) \in \mathbb{C}^2 : |x| > B, |x| > A|y|\}.$$

Consider also a domain in $\mathbb{R}^2$ defined as:

$$\Omega_{AB} = \{(x, y) \in \mathbb{R}^2 : x > B, x > A|y|\}.$$

It is easy to see that under certain conditions on A and B these domains possess some properties.

Theorem 1. *Let real numbers A and B be such that*

$$B > \frac{1}{2}(\frac{1}{A} + A) + \sqrt{\frac{1}{4}(\frac{1}{A} + A)^2 + |C|},\ B > 0,\ A > 1, \tag{4}$$

then for $\Omega \in \{\Omega_{AB}, \Omega^{\mathbb{C}}_{AB}\}$ the following conditions hold:

$$G(\Omega) \subset \Omega, \tag{5}$$

$$\forall z \in \Omega \quad \lim_{n \to +\infty} ||G^n(z)|| = \infty. \tag{6}$$

Moreover for Ω_{AB} one can point out an additional property:

$$\lim_{n\to+\infty} ||G^n(z)|| = \infty \Longrightarrow \exists\, m \in \mathbb{N} :\ G^m(z) \in \Omega_{AB}. \tag{7}$$

Consider the following sets invariant with respect to the quadratic map:

$$\Gamma_t = \{(x,y) \in \mathbb{R}^2 : \lim_{n\to+\infty} ||(x_n, y_n)|| = \infty\},$$

$$\Gamma_b = \{(x,y) \in \mathbb{R}^2 : \lim_{n\to+\infty} ||(x_{-n}, y_{-n})|| = \infty\},$$

$$B_0 = \Gamma_t \cup \Gamma_b,$$

$$A_t = \{(x,y) \in \mathbb{R}^2 : \{(x_n, y_n)\}_{n\in\mathbb{Z}_+} \in BOUND\},$$

$$A_b = \{(x,y) \in \mathbb{R}^2 : \{(x_n, y_n)\}_{n\in\mathbb{Z}_-} \in BOUND\},$$

$$A_0 = A_t \cap A_b, \quad D = \Gamma_t \cap \Gamma_b,$$

$$B_t = \Gamma_t \cap A_b \subset B_0, \quad B_b = \Gamma_b \cap A_t \subset B_0.$$

First of all, it is easy to observe the following topological properties of the defined above sets.

Theorem 2. *The sets B_0, D are open, A_t, A_b are closed. The set A_0 is compact. The sets B_t, B_b are closed in the topology of B_0.*

The next theorem is an important fact about these sets which is needed in the sequel.

Theorem 3. *Let mes be the Lebesgue measure in $\mathbb{R}^2$ then the following equalities are true:*

$$mesB_t = 0, \quad mesB_b = 0. \tag{9}$$

§2. Stable and unstable vector fields on the set D. Weak uniform hyperbolic structure

Let N_g be an open subset of $\mathbb{K}^d$ and let $G : N_g \longrightarrow N_g$ be a diffeomorphism.

Definition 1. *We say that an open set N_g possesses a stable g^s_λ and an unstable g^s_λ vector fields with respect to the map G with a parameter $\lambda > 1$ if the following conditions hold:*

$$g_\lambda^{s(u)} : N_g \longrightarrow \mathbb{K}^d, \tag{10}$$

$$\forall z \in N_g \ ||g_\lambda^{s(u)}|| = 1, \tag{11}$$

$$\forall z \in N_g \ \exists c > 0 : \ \forall n \in \mathbb{N} \ ||(G^n(z))' g^s_\lambda(z)|| \leqslant c\lambda^{-n},$$

$$||(G^{-n}(z))' g^u_\lambda(z)|| \leqslant c\lambda^{-n}. \tag{12}$$

Here is the basic statement of this section.

Theorem 4. *The set D possesses analytic vector fields k^s and k^u which are stable and unstable vector fields with respect to the map F for each parameter $\lambda > 1$.*

Proof. We construct the stable vector field, the unstable one can be constructed analogously. Consider an arbitrary $\lambda > 1$. We need an auxiliary complex-valued function $\Phi^{\mathbb{C}}_s$ defined on the domain $\Omega^{\mathbb{C}}_{AB}$, where $B = \lambda + \frac{3}{4}e_1$ and A is taken such that the condition (4) holds. Here e_1 is an arbitrary number from the interval $(0, 1)$. In order to construct the function $\Phi^{\mathbb{C}}_s$ we take the function Φ defined on $\mathbb{C}^2 \backslash (\{0\} \times \mathbb{C})$ giving us the ratio of the second component to the first one:

$$\Phi(t_1, t_2) = \frac{t_2}{t_1}.$$

Let $(x_0, y_0) \in \Omega^{\mathbb{C}}_{AB}$ be a point and $\{(x_n, y_n)\}_{n\in\mathbb{Z}}$ be its trajectory. Consider a sequence $\{l_n\}_{n\in\mathbb{N}}$ of vectors from $\mathbb{C}^2$ such that $(F^n(x, y))' l_n = (2\lambda + 2e_1, 1)$. It is easy to see that $\Phi(l_n)$ is an n-continuous fraction:

$$\Phi(l_n) = [2x_0; -2x_1, 2x_2, ..., (-1)^{n-1}2x_{n-1}, (-1)^n(2\lambda + 2e_1)].$$

As it follows from the continuous fractions theory, the sequence $\{\Phi(l_n)\}$ converges uniformly relatively $(x_0, y_0) \in \Omega^{\mathbb{C}}_{AB}$ to a limit $\Phi^{\mathbb{C}}_s(x_0, y_0)$, which does not depend on λ and on e_1. Moreover the following formula holds:

$$\Phi^{\mathbb{C}}_s(x_0, y_0) = [2x_0; -2x_1, 2x_2, ..., (-1)^n 2x_n, ...]. \tag{13}$$

So the function $\Phi^{\mathbb{C}}_s$ is constructed. Notice that $\Phi(l_n)$ depends analytically on a point $(x_0, y_0) \in \Omega^{\mathbb{C}}_{AB}$. Since the convergence is uniform, $\Phi^{\mathbb{C}}_s$ depends analytically on $(x_0, y_0) \in \Omega^{\mathbb{C}}_{AB}$. We construct the restriction Φ_s

of the function $\Phi_s^{\mathbb{C}}$ to the real domain Ω_{AB} by setting $\Phi_s = \Phi_s^{\mathbb{C}}|_{\Omega_{AB}}$. Obviously, the function Φ_s is analytic on Ω_{AB}. This implies that the vector k^s defined as $k^s = (\frac{1}{\sqrt{1+\Phi_s^2}}, \frac{\Phi_s}{\sqrt{1+\Phi_s^2}})$ is analytic on Ω_{AB}. Notice that the number Φ_s is the angle coefficient of the vector k^s. Moreover the construction implies that the 1-distribution defined by the field k^s is invariant under the map F, hence the field k^s can be continued analytically on the set D.

We show that the constructed field k^s is a stable vector field with respect to the map F with the parameter λ. It is clear that the conditions (10), (11) are true. We check now the condition (12). Take an arbitrary $z_0 \in \Omega_{AB}$, set $c = 1$. We show that the required inequality holds. Either $\lim_{n\to\infty} \frac{l_n}{||l_n||} = k^s$ or $\lim_{n\to\infty} \frac{l_n}{||l_n||} = -k^s$ holds. We consider the first case only, the second one can be considered in the same way. Take arbitrary $\varepsilon > 0$ and $n \in \mathbb{N}$. There exists a natural number N greater than n such that the following is true: $||(F^n(z_0))' k^s|| = ||(F^n(z_0))'(\frac{l_N}{||l_N||} + (k^s - \frac{l_N}{||l_N||}))|| \leqslant ||(F^n(z_0))' \frac{l_N}{||l_N||}|| + \varepsilon = ||v||/||u|| + \varepsilon$, where $v = [(F^{N-n})'(F^n(z_0))]^{-1}(2\lambda + 2e_1, 1), u = [F^n(z_0)]^{-1} v = l_N$. Thus we get:

$$||(F^n(z_0))' k^s|| \leqslant ||v||/||u|| + \varepsilon. \tag{14}$$

By some simple algebraic calculation one sees that $|v_1| > (\lambda + \frac{e_1}{4})|v_2|$ and then $||u||/||v|| > \lambda^n$. Substituting this into the inequality (14) one obtains:

$$||(F^n(z_0))' k^s|| \leqslant \lambda^{-n} + \varepsilon.$$

Since ε is arbitrary, we get:

$$||(F^n(z_0))' k^s|| \leqslant \lambda^{-n}.$$

Thus for $z_0 \in \Omega_{AB}$ the condition (12) holds. Suppose that $z_0 \in D$, consider $k = min\{k_1 \in \mathbb{Z}_+ : F^{k_1}(z) \in \Omega_{AB}\}$. Since the 1-distribution defined by the field k^s is invariant under the map F, one gets:

$$\exists \alpha \in \mathbb{R}_+ \backslash \{0\} : (F^k)'(z) k^s(z) = \alpha k_s(F^k(z)).$$

Set $c = \alpha \lambda^k$. Then the condition (12) holds, indeed: $||(F^n(z))' k^s(z)|| = ||(F^{n-k})'(F^k(z))(F^k)'(z) k_s(z)|| = |\alpha| ||(F^{n-k})'(F^k(z)) k^s(F^k(z))|| \leqslant |\alpha| \lambda^{-n} \lambda^k = c\lambda^{-n}$. Q.E.D.

We denote by Φ_s and Φ_u angle coefficients of the vector fields k^s and k^u respectively, where k^s and k^u are as in the last theorem. Notice that the numbers Φ_s and Φ_u can be infinite in some points of the set D.

Consider a real-analytic function S, being the sinus between the stable k^s and the unstable k^u vector fields, defined on the set D as

$$S = k^s_x k^u_y - k^s_y k^u_x. \tag{15}$$

Consider the following set:

$$Bad = \{z \in D : S(z) = 0\}.$$

We formulate an important results about it, which are basic components in the main theorem.

Theorem 5. *The following statements are true:*

$$Cl_D(Bad) = Bad, \tag{16}$$

$$F(Bad) = Bad, \tag{17}$$

$$mesBad = 0. \tag{18}$$

Proof. The statement (16) follows from the continuity of the map S. The statement (17) follows from the invariance of the 1-distributions defined by the fields k^s, k^u under the map F. Using the formula (13), it is easy to see that, there are regions where $S < 0$ and there are regions where $S > 0$, so the function S analytic on the set D is not identically zero, hence it vanishes on a set of Lebesgue measure zero. This shows the statement (18). Q.E.D.

The following theorem is the main result of this article.

Theorem 6. *There exist an open subset E and a compact subset H of the real plane $\mathbb{R}^2$, invariant with respect to the map F and such that the following statements hold:*

(i) *the set E is contained in $\mathbb{R}^2\backslash H$ and is of the full Lebesgue measure in $\mathbb{R}2\backslash H$,*

(ii) *The set E possesses a weak uniform hyperbolic structure with respect to the map F.*

Proof. Set $H = A_0$, $E = D\backslash Bad$. The equalities (16) and (17) and the fact that the set D is open imply that the set E is open and invariant with respect to the map F. The definition of the set E, the theorem 3 and the formula (18) imply the statement (i). The statement (ii) follows from the theorem 4 and the definition of the set Bad. This completes the proof of the theorem. Q.E.D.

Acknowledgments

The author is grateful to Prof. V. F. Lazutkin who formulated the problem and suggested a lot of ideas.

References

1 A. Ja. Hinchin, *Continuous fractions.* — Moscow, Science 1978.

St.Petersburg State University Поступило 30 ноября 2002 г.