Stability of the iterative process $x_{n+1} = Tx_n$ for Kannan maps in metric spaces

Arnoldo Bezanilla López
Pedro García Juárez
Facultad de Ciencias Físico Matemáticas
B. U. A. P. MEXICO.

Abstract. We prove the stability of an iterative process for a Kannan type map

1 Introduction

Let (X, d) be a metric space, $T : X \to X$ a map, $x_0 \in X$ and $x_{n+1} = Tx_n$, $n \in \mathbf{Z}_+$. Assume that T has one and only one fixed point p and that the sequence (x_n) converges to p. Let (y_n) be another sequence in X and $E_n = d(y_{n+1}, Ty_n)$, $n \in \mathbf{Z}_+$. We say that the iterative process $x_{n+1} = Tx_n$ is *stable* if $\lim_{n\to\infty} y_n = p$ whenever $\lim_{n\to\infty} E_n = 0$.

The stability of the iterative process $x_{n+1} = Tx_n$, $n \in \mathbf{Z}_+$, when T is a contraction, was obtained by V. A. Ostrovski in 1968 ([2]). It also was obtained by A. Harder and T. Hicks in 1988 when the application T satisfies the Zamfirescu's condition or similar ones ([1]).

Here, our purpose is to prove the stability of this iterative process when the application T satisfies certain condition of Kannan type.

Let (X, d) be a metric space. A map $T : X \to X$ is said to be of Kannan type if for all $x, y \in X$

$$d(Tx, Ty) \leq b(x, y)d(x, Tx) + b(y, x)d(y, Ty)$$

where $b : X \times X \to \mathbf{R}_+$ satisfies the following conditions

i) $b(x, y) + b(y, x) < 1, \quad x, y \in X$

The existence of fixed points for Kannan type maps has been studied by R. Kannan, when $b(x,y) = b(y,x) = k < \frac{1}{2}$ ([3]), S. Reich, when $b(x,y) = b(y,x) = \frac{1}{2}$ ([5]), S. Massa ([4]) and others.

The above definition of Kannan type maps was introduced by D. Roux and E. Maluta in 1978 (cf [6]).

In this paper, however, we need a stronger condition. So, in the sequel, we say that T in of Kannan type if

ii) $\sup_{x,y \in X} [b(x,y) + b(y,x)] = \lambda < 1$

A Kannan type map can not have more than one fixed point. From theorem 2 of [7], we can easily obtain that a Kannan type map on a complete metric space (X,d) has a fixed point p and that any sequence (x_n), where $x_{n+1} = Tx_n$, $n \in \mathbf{Z}_+$ and $x_0 \in X$, converges to p

Harder and Hicks ([1], theorem 2) proved the stability of the iterative process $x_{n+1} = Tx_n$, $n \in \mathbf{Z}_+$ for a Kannan type map T such what $b(x,y) \leq k < \frac{1}{2}$ uniformly in $x, y \in X$.

The objetive of this papers is to show the stability of this iterative process when T is a Kannan type map.

2 Stability

2.1 Lemma ([1] lemma 1) *Let c be a real number, $|c| < 1$, and (b_n) a real sequence such that $\lim_{k\to\infty} b_k = 0$, then*

$$\lim_{n\to\infty} \left(\sum_{k=0}^{n} c^{n-k} b_k \right) = 0.$$

2.2 Remark *Let X be a set, $b : X \times X \to \mathbf{R}$ a map that satisfies ii) and $c \in [\lambda, 1)$. Then, for all $x, y \in X$*

$$\frac{b(y,x)}{1 - b(x,y)} \leq c.$$

2.3 Theorem *Let (X,d) be a complete metric space and $T : X \longrightarrow X$ a Kannan type map. Fix $x_0 \in X$ and set $x_{n+1} = Tx_n$, $n \in \mathbf{Z}_+$ and $p = \lim_{n\to\infty} x_n$. If (y_n) is any sequence in X, $E_n = d(y_{n+1}, Ty_n)$ and $\lambda < c < 1$. Then,*

A)

$$d(y_{n+1},p) \leq E_n + b(x_n,y_n)d(x_n,Tx_n) + d(x_{n+1},p)$$

$$+\tfrac{E_{n-1}}{1-b(y_n,y_{n-1})} + \sum_{k=0}^{n-2} c^{n-k-1}\tfrac{E_k}{1-b(y_{k+1},y_k)} + c^n d(y_0,Ty_0)$$

B) If $\lim_{n\to\infty} E_n = 0$, then $\lim_{n\to\infty} y_n = p$.

Proof

A) For any possitive integer n,

$$\begin{aligned} d(y_{n+1},p) &\leq d(y_{n+1},Ty_n) + d(Ty_n,Tx_n) + d(Tx_{n+1},p) \\ &= E_n + d(Ty_n,Tx_n) + d(x_{n+1},p), \end{aligned}$$

Then, since T is a Kannan type map,

$$\begin{aligned} d(y_{n+1},p) &\leq E_n + b(y_n,x_n)d(y_n,Ty_n) + b(x_n,y_n)d(x_n,Tx_n) \\ &\quad + d(Tx_{n+1},p). \end{aligned} \tag{1}$$

On the other hand,

$$\begin{aligned} d(y_n,Ty_n) &\leq d(y_n,Ty_{n-1}) + d(Ty_{n-1},Ty_n) \\ &= E_{n-1} + d(Ty_{n-1},Ty_n) \\ &\leq E_{n-1} + b(y_{n-1},y_n)d(y_{n-1},Ty_{n-1}) + b(y_n,y_{n-1})d(y_n,Ty_n). \end{aligned}$$

Thus,

$$\{1 - b(y_n,y_{n-1})\}d(y_n,Ty_n) \leq E_{n-1} + b(y_{n-1},y_n)d(y_{n-1},Ty_{n-1})$$

and then

$$d(y_n, Ty_n) \leq \frac{E_{n-1}}{1 - b(y_n, y_{n-1})} + \frac{b(y_{n-1}, y_n)}{1 - b(y_n, y_{n-1})} d(y_{n-1}, Ty_{n-1}), \quad n > 1. \quad (2)$$

Now, using (2), we deduce

$$\begin{aligned} d(y_n, Ty_n) \quad &\leq \tfrac{E_{n-1}}{1-b(y_n, y_{n-1})} + \sum_{i=0}^{n-2} \left[\prod_{j=i+1}^{n-1} \tfrac{b(y_j, y_{j+1})}{1-b(y_{j+1}, y_j)} \right] \tfrac{E_i}{1-b(y_{i+1}, y_i)} \\ &+ \prod_{j=0}^{n-1} \tfrac{b(y_j, y_{j+1})}{1-b(y_{j+1}, y_j)} d(y_0, Ty_0). \end{aligned} \quad (3)$$

From the remark 2.2 we have that

$$\frac{b(y_j, y_{j+1})}{1 - b(y_{j+1}, y_j)} \leq c \qquad \text{for all} \quad j = 1, 2, \cdots, n.$$

and then it follows from (3) that

$$d(y_n, Ty_n) \leq \frac{E_{n-1}}{1 - b(y_n, y_{n-1})} + \sum_{i=0}^{n-2} c^{n-1-i} \frac{E_i}{1 - b(y_{i+1}, y_i)} + c^n d(y_0, Ty_0). \quad (4)$$

Taking into account that $b(y_n, x_n) < 1$, $n \in \mathbf{Z}_+$,we finally obtain from (1) and (4) that

$$\begin{aligned} d(y_{n+1}, p) \quad &\leq E_n + b(x_n, y_n) d(x_n, Tx_n) + d(x_{n+1}, p) \\ &+ \tfrac{E_{n-1}}{1-b(y_n, y_{n-1})} + \sum_{k=0}^{n-2} c^{n-k-1} \tfrac{E_k}{1-b(y_{k+1}, y_k)} + c^n d(y_0, Ty_0) \end{aligned}$$

B) Suppose that $\lim_{n \to \infty} E_n = 0$. Take c such that $\lambda < c < 1$. Since $b(x, y) \leq \lambda < 1$ for all $x, y \in X$, we obtain from A) that

$$\begin{aligned} 0 \leq d(y_{n+1}, p) \qquad &\leq E_n + \lambda d(x_n, Tx_n) \\ &+ d(x_{n+1}, p) + \tfrac{E_{n-1}}{1-\lambda} \\ + \sum_{k=0}^{n-2} c^{n-k-1} \tfrac{E_k}{1-\lambda} & \\ + c^n d(y_0, Ty_0). & \end{aligned}$$

According to lemma 2.1, $\lim_{n\to\infty}\left(\sum_{k=0}^{n-2} c^{n-k-1}\frac{E_k}{1-\lambda}\right) = 0$. Since $x_n \to p$, we deduce that $d(x_{n+1},p) \to 0$ and $d(x_n,Tx_n) \to 0$ if $n \to \infty$. Therefore $\lim_{n\to\infty} d(y_{n+1},p) = 0$ ■

References

[1] Harder, A.M and Hicks T.L. *Stability results for fixed point iteration procedures*. Math. Jap. 33, No. 5, 1988, 693-706.

[2] Istratescu, V.I., *Fixed point Theory*, Dordrecht, Holland D. Reidel Publishing Company, 1981.

[3] Kannan, R. *Some results on fixed points*, Bull. Calcutta Math. Soc., 60, 1968, 71-76.

[4] Massa, S. , *Generalized contractions in metric spaces*. Boll. Un. Mat. Ital., (4) 10, 1974, 689-694.

[5] Reich, S., *Kannan's fixed point theorem*. Boll. Un. Mat. Ital., (4) 4, 1971, 1-11.

[6] Roux, D. and Maluta, E., *Contractive Kannan maps in compact spaces*, Milano - Istituto matemático "F. Enriques", Quaderno n. 17/p, 1978.

[7] Tiberio Bianchini, R. M., *Su un problema di S. Reich riguardante la teoria dei punti fissi*, Boll. Un. Mat. Ital., 5, 1972), 103-108.